




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : C07D 403/04, 401/14, A61K 31/505	A1	(11) International Publication Number: WO 97/42187 (43) International Publication Date: 13 November 1997 (13.11.97)
(21) International Application Number: PCT/GB97/01211 (22) International Filing Date: 2 May 1997 (02.05.97) (30) Priority Data: 96400956.7 6 May 1996 (06.05.96) EP (34) Countries for which the regional or international application was filed: AT et al. 96400957.5 6 May 1996 (06.05.96) EP (34) Countries for which the regional or international application was filed: AT et al. 96402762.7 17 December 1996 (17.12.96) EP (34) Countries for which the regional or international application was filed: AT et al. 96402763.5 17 December 1996 (17.12.96) EP (34) Countries for which the regional or international application was filed: AT et al. (71) Applicants (for all designated States except US): ZENECA LIMITED [GB/GB]; 15 Stanhope Gate, London W1Y 6LN (GB). ZENECA PHARMA S.A. [FR/FR]; 1, rue des Chauffours, Boite postale 127, F-95022 Cergy Cedex (FR). (72) Inventors; and (75) Inventors/Applicants (for US only): THOMAS, Andrew, Peter [GB/GB]; Alderley Park, Macclesfield, Cheshire SK10 4TG		(GB). HENNEQUIN, Laurent, François, André [FR/FR]; Zone Industrielle Sud-est, Boite postale 401, F-51064 Reims Cedex (FR). LOHMANN, Jean-Jacques, Marcel [FR/FR]; Zone Industrielle Sud-est, Boite postale 401, F-51064 Reims Cedex (FR). PLE, Patrick [FR/FR]; Zone Industrielle Sud-est, Boite postale 401, F-51064 Reims Cedex (FR). (74) Agent: MACK, John, Richard; Zeneca Pharmaceuticals, Intellectual Property Dept., Mereside, Alderley Park, Macclesfield, Cheshire SK10 4TG (GB). (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published With international search report.
(54) Title: OXINDOLE DERIVATIVES (57) Abstract <p>The invention relates to compounds of formula (I), wherein: R² represents hydroxy, halogeno, C₁-3alkyl, C₁-3alkoxy, C₁-3alkanoyloxy, trifluoromethyl, cyano, amino, nitro, C₂-4alkanoyl, C₁-4alkanoylamino, C₁-4alkoxycarbonyl, C₁-4alkylthio, C₁-4alkylsulphinyl, C₁-4alkylsulphonyl, carbamoyl, N-C₁-4alkylcarbamoyl, N,N-di(C₁-4alkyl)carbamoyl, aminosulphonyl, N-C₁-4alkylaminosulphonyl, N,N-di(C₁-4alkyl)aminosulphonyl, C₁-4alkylsulphonylamino, or a group R⁴X¹ wherein X¹ represents a direct bond, C₂-4alkanoyl, -CONR⁵R⁶-, -SO₂NR⁷R⁸-, or -SO₂R⁹- (wherein R⁵ and R⁷, each independently represents hydrogen or C₁-2alkyl and R⁶, R⁸ and R⁹ each independently represents C₁-4alkyl and wherein R⁴ is linked to R⁶, R⁸ or R⁹) and R⁴ represents an optionally substituted group selected from phenyl and a 5 or 6-membered heterocyclic group; n is an integer from 0 to 4; R¹ represents hydrogen, C₁-4alkyl, C₁-4alkoxymethyl, di(C₁-4alkoxy)methyl or C₁-4alkanoyl; m is an integer from 0 to 4; and R³ represents hydroxy, halogeno, nitro, trifluoromethyl, C₁-3alkyl, cyano, amino or R¹⁰X² (wherein X² represents a direct bond, -CH₂-, or a single or double heteroatom linker group including -S-, -SO- and -NR¹⁵- (wherein R¹⁵ represents hydrogen, C₁-3alkyl or C₁-3alkoxyC₂-3alkyl), and R¹⁰ is an alkenyl or alkynyl chain optionally substituted by for example hydroxy, amino, nitro, alkyl, cycloalkyl, alkoxyalkyl, or an optionally substituted group selected from pyridone, phenyl and a heterocyclic ring, which alkyl, alkenyl or alkynyl chain may have a heteroatom linker group, or R¹⁰ is an optionally substituted group selected from pyridone, phenyl and a heterocyclic ring. The compounds of formula (I) and the pharmaceutically acceptable salts thereof inhibit the effects of VEGF and FGF, properties of value in the treatment of a number of disease states including cancer and rheumatoid arthritis.</p>		



(I)

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TC	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

OXINDOLE DERIVATIVES

The present invention relates to oxindole derivatives, processes for their preparation, pharmaceutical compositions containing them as active ingredient, methods for the treatment of disease states associated with angiogenesis and/or increased vascular permeability, to their use as medicaments and to their use in the manufacture of medicaments for use in the production of antiangiogenic and/or vascular permeability reducing effects in warm-blooded animals such as humans.

Normal angiogenesis plays an important role in a variety of processes including embryonic development, wound healing and several components of female reproductive function. Undesirable or pathological angiogenesis has been associated with disease states including diabetic retinopathy, psoriasis, cancer, rheumatoid arthritis, atheroma, Kaposi's sarcoma and haemangioma (Fan et al, 1995, Trends Pharmacol. Sci. 16: 57-66; Folkman, 1995, Nature Medicine 1: 27-31). Alteration of vascular permeability is thought to play a role in both normal and pathological physiological processes (Cullinan-Bove et al, 1993, Endocrinology 133: 829-837; Senger et al, 1993, Cancer and Metastasis Reviews, 12: 303-324). Several polypeptides with in vitro endothelial cell growth promoting activity have been identified including, acidic and basic fibroblast growth factors (aFGF & bFGF) and vascular endothelial growth factor (VEGF). By virtue of the restricted expression of its receptors, the growth factor activity of VEGF, in contrast to that of the FGFs, is relatively specific towards endothelial cells. Recent evidence indicates that VEGF is an important stimulator of both normal and pathological angiogenesis (Jakeman et al, 1993, Endocrinology, 133: 848-859; Kolch et al, 1995, Breast Cancer Research and Treatment, 36:139-155) and vascular permeability (Connolly et al, 1989, J. Biol. Chem. 264: 20017-20024). Antagonism of VEGF action by sequestration of VEGF with antibody can result in inhibition of tumour growth (Kim et al, 1993, Nature 362: 841-844). Basic FGF (bFGF) is a potent stimulator of angiogenesis (e.g. Hayek et al, 1987, Biochem. Biophys. Res. Commun. 147: 876-880) and raised levels of FGFs have been found in the serum (Fujimoto et al, 1991, Biochem. Biophys. Res. Commun. 180: 386-392) and urine (Nguyen et al, 1993, J. Natl. Cancer. Inst. 85: 241-242) of patients with cancer.


Receptor tyrosine kinases (RTKs) are important in the transmission of biochemical signals across the plasma membrane of cells. These transmembrane molecules characteristically consist of an extracellular ligand-binding domain connected through a segment in the plasma membrane to an intracellular tyrosine kinase domain. Binding of
5 ligand to the receptor results in stimulation of the receptor-associated tyrosine kinase activity which leads to phosphorylation of tyrosine residues on both the receptor and other intracellular molecules. These changes in tyrosine phosphorylation initiate a signalling cascade leading to a variety of cellular responses. To date, at least nineteen distinct RTK subfamilies, defined by amino acid sequence homology, have been identified. One of these
10 subfamilies is presently comprised by the *fms*-like tyrosine kinase receptor, Flt or Flt1, the kinase insert domain-containing receptor, KDR (also referred to as Flk-1), and another *fms*-like tyrosine kinase receptor, Flt4. Two of these related RTKs, Flt and KDR, have been shown to bind VEGF with high affinity (De Vries et al, 1992, Science 255: 989-991; Terman et al, 1992, Biochem. Biophys. Res. Comm. 1992, 187: 1579-1586). Binding of VEGF to
15 these receptors expressed in heterologous cells has been associated with changes in the tyrosine phosphorylation status of cellular proteins and calcium fluxes.

The present invention is based on the discovery of compounds that surprisingly inhibit the effects of VEGF and FGF, properties of value in the treatment of disease states associated with angiogenesis and/or increased vascular permeability such as cancer, diabetes, psoriasis,
20 rheumatoid arthritis, Kaposi's sarcoma, haemangioma, acute and chronic nephropathies, atheroma, arterial restenosis, autoimmune diseases, acute inflammation and ocular diseases with retinal vessel proliferation. Compounds of the present invention possess higher potency against VEGF receptor tyrosine kinase and against FGF R1 receptor tyrosine kinase than against epidermal growth factor (EGF) receptor tyrosine kinase. Furthermore, compounds of
25 the present invention, possess substantially higher potency against VEGF receptor tyrosine kinase and against FGF R1 receptor tyrosine kinase than against EGF receptor tyrosine kinase. Compounds of the invention which have been tested possess activity against VEGF receptor tyrosine kinase and against FGF R1 receptor tyrosine kinase such that they may be used in an amount sufficient to inhibit VEGF receptor tyrosine kinase and FGF R1 receptor
30 tyrosine kinase whilst demonstrating no significant activity against EGF receptor tyrosine kinase. Thus compounds of the present invention possess good VEGF receptor tyrosine

- 3 -

kinase activity and good FGF R1 receptor tyrosine kinase activity. Compounds with both VEGF receptor tyrosine kinase activity and FGF R1 receptor tyrosine kinase activity are believed to be of particular value in the treatment of disease states associated with angiogenesis and/or increased vascular permeability.

5 According to one aspect of the present invention there are provided compounds of the formula I:



(I)

15 {wherein:

R² represents hydroxy, halogeno, C₁₋₃alkyl, C₁₋₃alkoxy, C₁₋₃alkanoyloxy, trifluoromethyl, cyano, amino or nitro, and additional values for R² are C₂₋₄alkanoyl, C₁₋₄alkanoylamino, C₁₋₄alkoxycarbonyl, C₁₋₄alkylthio, C₁₋₄alkylsulphinyl, C₁₋₄alkylsulphonyl, carbamoyl, N-C₁₋₄alkylcarbamoyl, N,N-di(C₁₋₄alkyl)carbamoyl, aminosulphonyl, N-C₁₋₄alkylaminosulphonyl, N,N-di(C₁₋₄alkyl)aminosulphonyl, C₁₋₄alkylsulphonylamino, or a group R⁴X¹ wherein X¹ represents a direct bond, C₂₋₄alkanoyl, -CONR⁵R⁶-, -SO₂NR⁷R⁸- or -SO₂R⁹- (wherein R⁵ and R⁷, each independently represents hydrogen or C₁₋₂alkyl and R⁶, R⁸ and R⁹ each independently represents C₁₋₄alkyl and wherein R⁴ is linked to R⁶, R⁸ or R⁹) and R⁴ represents phenyl or a 5 or 6-membered heterocyclic group with one or two heteroatoms, selected independently from O, S and N, which heterocyclic group may be saturated or unsaturated and which phenyl or heterocyclic group may bear one or two substituents selected from hydroxy, halogeno, C₁₋₃alkyl, C₁₋₃alkoxy, C₁₋₃alkanoyloxy, trifluoromethyl, cyano, amino, nitro and C₁₋₄alkoxycarbonyl;

n is an integer from 0 to 4;

30 R¹ represents hydrogen, C₁₋₄alkyl, C₁₋₄alkoxymethyl, di(C₁₋₄alkoxy)methyl or C₁₋₄alkanoyl;

m is an integer from 0 to 4; and

R³ represents hydroxy, halogeno, nitro, trifluoromethyl, C₁₋₃alkyl, cyano, amino or R¹⁰X² (wherein X² represents -O-, -CH₂-, -S-, -SO-, -SO₂-, -NR¹¹CO-, -CONR¹²-, -SO₂NR¹³-, -NR¹⁴SO₂- or -NR¹⁵- (wherein R¹¹, R¹², R¹³, R¹⁴ and R¹⁵ each independently represents

5 hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl), or X² represents a direct bond, and R¹⁰ is selected from one of the following fifteen groups:

1) hydrogen or C₁₋₃alkyl which may be unsubstituted or which may be substituted with one or more groups selected from hydroxy, fluoro and amino;

2) C₁₋₃alkylX³COR¹⁶ (wherein X³ represents -O- or -NR¹⁷- (in which R¹⁷ represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R¹⁶ represents C₁₋₃alkyl, -NR¹⁸R¹⁹- or -OR²⁰- (wherein R¹⁸, R¹⁹ and R²⁰ which may be the same or different each represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl));

3) C₁₋₃alkylX⁴R²¹ (wherein X⁴ represents -O-, -S-, -SO-, -SO₂-, -OCO-, -NR²²CO-, -CONR²³-, -SO₂NR²⁴-, -NR²⁵SO₂- or -NR²⁶- (wherein R²², R²³, R²⁴, R²⁵ and R²⁶ each independently represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R²¹ represents hydrogen, C₁₋₃alkyl, cyclopentyl, cyclohexyl or a 5 or 6 membered saturated heterocyclic group with one or two heteroatoms, selected independently from O, S and N, which C₁₋₃alkyl group may bear one or two substituents selected from oxo, hydroxy, halogeno and C₁₋₄alkoxy and which cyclic group may bear one or two substituents selected from oxo, hydroxy, halogeno, C₁₋₄alkyl, C₁₋₄hydroxyalkyl and C₁₋₄alkoxy);

4) C₁₋₃alkylX⁵C₁₋₃alkylX⁶R²⁷ (wherein X⁵ and X⁶ which may be the same or different are each -O-, -S-, -SO-, -SO₂-, -NR²⁸CO-, -CONR²⁹-, -SO₂NR³⁰-, -NR³¹SO₂- or -NR³²- (wherein R²⁸, R²⁹, R³⁰, R³¹ and R³² each independently represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R²⁷ represents hydrogen or C₁₋₃alkyl);

5) C₁₋₃alkylR³³ (wherein R³³ is a 5 or 6 membered saturated heterocyclic group with one or two heteroatoms, selected independently from O, S and N, which heterocyclic group may bear one or two substituents selected from oxo, hydroxy, halogeno, C₁₋₄alkyl, C₁₋₄hydroxyalkyl and C₁₋₄alkoxy);

6) C₂₋₃alkenylR³³ (wherein R³³ is as defined hereinbefore);

7) C₂₋₃alkynylR³³ (wherein R³³ is as defined hereinbefore);

- 8) R^{34} (wherein R^{34} represents a pyridone group, a phenyl group or a 5 or 6-membered aromatic heterocyclic group with 1 to 3 heteroatoms selected from O, N and S, which pyridone, phenyl or heterocyclic group may carry up to 5 substituents selected from halogeno, amino, C_{1-4} alkyl, C_{1-4} alkoxy, C_{1-4} hydroxyalkyl, C_{1-4} aminoalkyl, C_{1-4} alkylamino, C_{1-4} hydroxyalkoxy, carboxy, cyano, $-\text{CONR}^{35}\text{R}^{36}$ and $-\text{NR}^{37}\text{COR}^{38}$ (wherein R^{35} , R^{36} , R^{37} and R^{38} , which may be the same or different, each represents hydrogen, C_{1-4} alkyl or C_{1-3} alkoxy C_{2-3} alkyl)) and an additional possible substituent of R^{34} is hydroxy;
- 9) C_{1-3} alkyl R^{34} (wherein R^{34} is as defined hereinbefore);
- 10) C_{2-3} alkenyl R^{34} (wherein R^{34} is as defined hereinbefore);
- 10 11) C_{2-3} alkynyl R^{34} (wherein R^{34} is as defined hereinbefore);
- 12) C_{1-3} alkyl X^7R^{34} (wherein X^7 represents $-\text{O}-$, $-\text{S}-$, $-\text{SO}-$, $-\text{SO}_2-$, $-\text{NR}^{39}\text{CO}-$, $-\text{CONR}^{40}-$, $-\text{SO}_2\text{NR}^{41}-$, $-\text{NR}^{42}\text{SO}_2-$ or $-\text{NR}^{43}-$ (wherein R^{39} , R^{40} , R^{41} , R^{42} and R^{43} each independently represents hydrogen, C_{1-3} alkyl or C_{1-3} alkoxy C_{2-3} alkyl) and R^{34} is as defined hereinbefore);
- 13) C_{2-3} alkenyl X^8R^{34} (wherein X^8 represents $-\text{O}-$, $-\text{S}-$, $-\text{SO}-$, $-\text{SO}_2-$, $-\text{NR}^{44}\text{CO}-$, $-\text{CONR}^{45}-$, $-\text{SO}_2\text{NR}^{46}-$, $-\text{NR}^{47}\text{SO}_2-$ or $-\text{NR}^{48}-$ (wherein R^{44} , R^{45} , R^{46} , R^{47} and R^{48} each independently represents hydrogen, C_{1-3} alkyl or C_{1-3} alkoxy C_{2-3} alkyl) and R^{34} is as defined hereinbefore);
- 15 14) C_{2-3} alkynyl X^9R^{34} (wherein X^9 represents $-\text{O}-$, $-\text{S}-$, $-\text{SO}-$, $-\text{SO}_2-$, $-\text{NR}^{49}\text{CO}-$, $-\text{CONR}^{50}-$, $-\text{SO}_2\text{NR}^{51}-$, $-\text{NR}^{52}\text{SO}_2-$ or $-\text{NR}^{53}-$ (wherein R^{49} , R^{50} , R^{51} , R^{52} and R^{53} each independently represents hydrogen, C_{1-3} alkyl or C_{1-3} alkoxy C_{2-3} alkyl) and R^{34} is as defined hereinbefore); and
- 20 15) C_{1-3} alkyl $X^{10}C_{1-3}$ alkyl R^{34} (wherein X^{10} represents $-\text{O}-$, $-\text{S}-$, $-\text{SO}-$, $-\text{SO}_2-$, $-\text{NR}^{54}\text{CO}-$, $-\text{CONR}^{55}-$, $-\text{SO}_2\text{NR}^{56}-$, $-\text{NR}^{57}\text{SO}_2-$ or $-\text{NR}^{58}-$ (wherein R^{54} , R^{55} , R^{56} , R^{57} and R^{58} each independently represents hydrogen, C_{1-3} alkyl or C_{1-3} alkoxy C_{2-3} alkyl) and R^{34} is as defined hereinbefore); and additional groups of values of R^{10} are:
- 16) R^{33} (wherein R^{33} is as defined hereinbefore); and
- 25 17) C_{1-3} alkyl $X^{10}C_{1-3}$ alkyl R^{33} (wherein X^{10} and R^{33} are as defined hereinbefore)); and salts thereof.

Advantageously X^1 represents a direct bond C_{2-4} alkanoyl, $-\text{CONR}^5\text{R}^6-$, $-\text{SO}_2\text{NR}^7\text{R}^8-$ or $-\text{SO}_2\text{R}^9-$ (wherein R^5 and R^7 , each independently represents hydrogen or C_{1-2} alkyl and R^6 , R^8 and R^9 each independently represents C_{1-3} alkyl and wherein R^4 is linked to R^6 , R^8 or R^9).

Preferably X^1 represents a direct bond C_{2-4} alkanoyl, $-CONR^5R^6-$, $-SO_2NR^7R^8-$ or $-SO_2R^9-$ (wherein R^5 and R^7 , each independently represents hydrogen and R^6 , R^8 and R^9 each independently represents C_{1-3} alkyl and wherein R^4 is linked to R^6 , R^8 or R^9).

Advantageously R^2 represents hydroxy, halogeno, C_{1-2} alkyl, C_{1-2} alkoxy, trifluoromethyl, cyano, nitro, C_{2-3} alkanoyl, C_{1-3} alkanoylamino, C_{1-3} alkoxycarbonyl, C_{1-3} alkylthio, C_{1-3} alkylsulphinyl, C_{1-3} alkylsulphonyl, carbamoyl, N - C_{1-3} alkylcarbamoyl, N,N -di(C_{1-3} alkyl)carbamoyl, aminosulphonyl, N - C_{1-3} alkylaminosulphonyl, N,N -di(C_{1-3} alkyl)aminosulphonyl, C_{1-3} alkylsulphonylamino, or a group R^4X^1 (wherein X^1 is as defined hereinbefore and R^4 represents phenyl or a 5 or 6-membered heterocyclic group with one or two heteroatoms, selected independently from O, S and N, which heterocyclic group may be saturated or unsaturated and which phenyl or heterocyclic group may bear one or two substituents selected from hydroxy, halogeno, C_{1-2} alkyl, C_{1-2} alkoxy, C_{1-2} alkanoyloxy, trifluoromethyl, cyano, amino, nitro and C_{1-4} alkoxycarbonyl).

Preferably R^2 represents halogeno, trifluoromethyl, cyano, nitro, C_{2-3} alkanoyl, C_{1-3} alkoxycarbonyl, C_{1-3} alkylsulphinyl, C_{1-3} alkylsulphonyl, carbamoyl, N - C_{1-3} alkylcarbamoyl, N,N -di(C_{1-3} alkyl)carbamoyl, aminosulphonyl, N - C_{1-3} alkylaminosulphonyl, N,N -di(C_{1-3} alkyl)aminosulphonyl, or a group R^4X^1 (wherein X^1 is as defined hereinbefore and R^4 represents phenyl or a 5 or 6-membered heterocyclic group with one or two heteroatoms, selected independently from O, S and N, of which at least one is N, which heterocyclic group may be saturated or unsaturated and which phenyl or heterocyclic group may bear one or two substituents selected from hydroxy, halogeno, C_{1-2} alkyl, C_{1-2} alkoxy, C_{1-2} alkanoyloxy, trifluoromethyl, cyano, amino, nitro and C_{1-4} alkoxycarbonyl).

More preferably R^2 represents halogeno, trifluoromethyl, cyano, nitro, C_{2-3} alkanoyl, C_{1-3} alkoxycarbonyl, C_{1-3} alkylsulphinyl, C_{1-3} alkylsulphonyl, carbamoyl, N - C_{1-3} alkylcarbamoyl, N,N -di(C_{1-3} alkyl)carbamoyl, aminosulphonyl, N - C_{1-3} alkylaminosulphonyl, N,N -di(C_{1-3} alkyl)aminosulphonyl, or a group R^4X^1 (wherein X^1 is as defined hereinbefore and R^4 represents pyrrolidinyl, piperazinyl, piperidinyl, morpholino, thiomorpholino, phenyl, pyridyl, imidazolyl, thiazolyl, thienyl, triazolyl or pyridazinyl which phenyl or heterocyclic group may bear one or two substituents selected from hydroxy, halogeno, C_{1-2} alkyl, C_{1-2} alkoxy, C_{1-2} alkanoyloxy, trifluoromethyl, cyano, amino, nitro and C_{1-4} alkoxycarbonyl).

- Especially preferred values of R^2 are halogeno, trifluoromethyl, cyano, nitro, $C_{2,3}$ alkanoyl, $C_{1,2}$ alkoxycarbonyl, $C_{1,3}$ alkylsulphinyl, $C_{1,3}$ alkylsulphonyl, carbamoyl, \underline{N} - $C_{1,3}$ alkylcarbamoyl, $\underline{N,N}$ -di($C_{1,3}$ alkyl)carbamoyl, aminosulphonyl, \underline{N} - $C_{1,3}$ alkylaminosulphonyl, $\underline{N,N}$ -di($C_{1,3}$ alkyl)aminosulphonyl, or a group R^4X^1 (wherein X^1 is as defined hereinbefore and R^4 represents pyrrolidinyl, piperazinyl, piperidinyl, morpholino, thiomorpholino, imidazolyl or triazolyl which heterocyclic group is linked to X^1 via a nitrogen atom and which heterocyclic group may bear one or two substituents selected from hydroxy, halogeno, $C_{1,2}$ alkyl, $C_{1,2}$ alkoxy, $C_{1,2}$ alkanoyloxy, trifluoromethyl, cyano, amino, nitro and $C_{1,4}$ alkoxycarbonyl). In another aspect of the present invention R^2 represents hydroxy, halogeno, $C_{1,3}$ alkyl, $C_{1,3}$ alkoxy, trifluoromethyl, cyano, amino or nitro.
- In another aspect of the present invention R^2 represents hydroxy, halogeno, $C_{1,3}$ alkyl, $C_{1,3}$ alkoxy or trifluoromethyl, especially hydroxy, chloro, fluoro, methyl or methoxy.
- Preferably n is an integer from 0 to 2.
- Preferably R^1 represents hydrogen, $C_{1,4}$ alkoxymethyl, di($C_{1,4}$ alkoxy)methyl or $C_{1,4}$ alkanoyl, especially hydrogen.
- Preferably m is an integer from 0-2, most preferably 2.
- Advantageously X^2 represents -O-, -S-, -NR¹¹CO-, -NR¹⁴SO₂- or -NR¹⁵- (wherein R^{11} , R^{14} and R^{15} each independently represents hydrogen, $C_{1,2}$ alkyl or $C_{1,2}$ alkoxyethyl).
- Preferably X^2 represents -O-, -S-, -NR¹¹CO-, -NR¹⁴SO₂- (wherein R^{11} and R^{14} each independently represents hydrogen or $C_{1,2}$ alkyl) or NH.
- More preferably X^2 represents -O-, -S-, -NR¹¹CO- (wherein R^{11} represents hydrogen or $C_{1,2}$ alkyl) or NH.
- Particularly X^2 represents -O- or -NR¹¹CO- (wherein R^{11} represents hydrogen or $C_{1,2}$ alkyl), more particularly -O- or -NHCO-, especially -O-.
- Advantageously X^3 represents -O- or NR¹⁷ (wherein R^{17} represents hydrogen, $C_{1,3}$ alkyl or $C_{1,2}$ alkoxyethyl).
- Advantageously X^4 represents -O-, -S-, -SO-, -SO₂-, -NR²²CO-, -NR²⁵SO₂- or -NR²⁶- (wherein R^{22} , R^{25} and R^{26} each independently represents hydrogen, $C_{1,2}$ alkyl or $C_{1,2}$ alkoxyethyl).
- Preferably X^4 represents -O-, -S-, -SO-, -SO₂- or -NR²⁶- (wherein R^{26} represents hydrogen, $C_{1,2}$ alkyl or $C_{1,2}$ alkoxyethyl).
- More preferably X^4 represents -O- or -NR²⁶- (wherein R^{26} represents hydrogen or $C_{1,2}$ alkyl).

Advantageously X^5 and X^6 which may be the same or different each represents -O-, -S-, -SO-, -SO₂- or -NR³²- (wherein R³² represents hydrogen, C₁₋₃alkyl or C₁₋₂alkoxyethyl).

Preferably X^5 and X^6 which may be the same or different each represents -O-, -S- or -NR³²- (wherein R³² represents hydrogen, C₁₋₂alkyl or C₁₋₂alkoxyethyl).

- 5 More preferably X^5 and X^6 which may be the same or different each represents -O- or -NH-. Advantageously X^7 represents -O-, -S- or -NR⁴³- (wherein R⁴³ represents hydrogen, C₁₋₂alkyl or C₁₋₂alkoxyethyl).

Preferably X^7 represents -O- or -NR⁴³- (wherein R⁴³ represents hydrogen or C₁₋₂alkyl).

- Advantageously X^8 represents -O-, -S- or -NR⁴⁸- (wherein R⁴⁸ represents hydrogen, C₁₋₂alkyl or C₁₋₂alkoxyethyl).

Preferably X^8 represents -O- or -NR⁴⁸- (wherein R⁴⁸ represents hydrogen or C₁₋₂alkyl).

Advantageously X^9 represents -O-, -S- or -NR⁵³- (wherein R⁵³ represents hydrogen, C₁₋₂alkyl or C₁₋₂alkoxyethyl).

Preferably X^9 represents -O- or -NR⁵³- (wherein R⁵³ represents hydrogen or C₁₋₂alkyl).

- 15 Advantageously X^{10} represents -O-, -S- or -NR⁵⁸- (wherein R⁵⁸ represents hydrogen, C₁₋₂alkyl or C₁₋₂alkoxyethyl).

Preferably X^{10} represents -O- or -NR⁵⁸- (wherein R⁵⁸ represents hydrogen or C₁₋₂alkyl).

- R³³ is preferably pyrrolidinyl, piperazinyl, piperidinyl, morpholino or thiomorpholino which group may carry one or two substituents selected from oxo, hydroxy, halogeno, C₁₋₂alkyl, C₁₋₂hydroxyalkyl and C₁₋₂alkoxy.

R³⁴ preferably represents a pyridone group or a 5 or 6-membered aromatic heterocyclic group with 1 to 3 heteroatoms selected from O, N and S, which pyridone group or heterocyclic group may be substituted as hereinbefore defined.

- Where R³⁴ is a 5 or 6-membered aromatic heterocyclic group, it preferably has 1 or 2 heteroatoms, selected from O, N and S, of which more preferably one is N, and may be substituted as hereinbefore defined.

- R³⁴ is particularly a pyridone, pyridyl, imidazolyl, thiazolyl, thienyl, triazolyl or pyridazinyl group which group may be substituted as hereinbefore defined, more particularly a pyridone, pyridyl, imidazolyl, thiazolyl or triazolyl group, especially a pyridone, pyridyl, imidazolyl or triazolyl group which group may be substituted as hereinbefore defined.

In one embodiment of the invention R^{34} represents a pyridone, phenyl or 5 or 6-membered aromatic heterocyclic group with 1 to 3 heteroatoms selected from O, N and S, which group may preferably carry up to 2 substituents, more preferably up to one substituent, selected from the group of substituents as hereinbefore defined.

- 5 In the definition of R^{34} , conveniently substituents are selected from halogeno, C_{1-4} alkyl, C_{1-4} alkoxy and cyano, more conveniently substituents are selected from chloro, fluoro, methyl and ethyl.

Conveniently R^3 represents hydroxy, halogeno, nitro, trifluoromethyl, C_{1-3} alkyl, cyano, amino or $R^{10}X^2$ [wherein X^2 is as hereinbefore defined and R^{10} is selected from one of the following

- 10 fifteen groups:

1) C_{1-3} alkyl which may be unsubstituted or substituted with one or more fluorine atoms, or C_{2-3} alkyl which may be unsubstituted or substituted with one or more groups selected from hydroxy and amino;

- 2) C_{2-3} alkyl X^3COR^{16} (wherein X^3 is as hereinbefore defined and R^{16} represents C_{1-3} alkyl, -
15 $NR^{18}R^{19}$ - or $-OR^{20}$ - (wherein R^{18} , R^{19} and R^{20} which may be the same or different are each C_{1-2} alkyl or C_{1-2} alkoxyethyl));

- 3) C_{2-4} alkyl X^4R^{21} (wherein X^4 is as hereinbefore defined and R^{21} represents hydrogen, C_{1-3} alkyl, cyclopentyl, cyclohexyl or a 5 or 6 membered saturated heterocyclic group with one or two heteroatoms, selected independently from O, S and N, which C_{1-3} alkyl group may bear
20 one or two substituents selected from oxo, hydroxy, halogeno and C_{1-3} alkoxy and which cyclic group may bear one or two substituents selected from oxo, hydroxy, halogeno, C_{1-4} alkyl, C_{1-4} hydroxyalkyl and C_{1-4} alkoxy);

4) C_{2-3} alkyl X^5C_{2-3} alkyl X^6R^{27} (wherein X^5 and X^6 are as hereinbefore defined and R^{27} represents hydrogen or C_{1-3} alkyl);

- 25 5) C_{1-3} alkyl R^{59} (wherein R^{59} is a 5 or 6 membered saturated heterocyclic group with one or two heteroatoms, selected independently from O, S and N, which heterocyclic group is linked to C_{1-3} alkyl through a carbon atom and which heterocyclic group may bear one or two substituents selected from oxo, hydroxy, halogeno, C_{1-4} alkyl, C_{1-4} hydroxyalkyl and C_{1-4} alkoxy) or C_{2-3} alkyl R^{60} (wherein R^{60} is a 5 or 6 membered saturated heterocyclic group with one or
30 two heteroatoms of which one is N and the other is selected independently from O, S and N, which heterocyclic group is linked to C_{2-3} alkyl through a nitrogen atom and which

heterocyclic group may bear one or two substituents selected from oxo, hydroxy, halogeno, C_{1-4} alkyl, C_{1-4} hydroxyalkyl and C_{1-4} alkoxy);

6) C_{3-4} alkenylR⁶¹ (wherein R⁶¹ represents R⁵⁹ or R⁶⁰ as defined hereinbefore);

7) C_{3-4} alkynylR⁶¹ (wherein R⁶¹ represents R⁵⁹ or R⁶⁰ as defined hereinbefore);

5 8) R³⁴ (wherein R³⁴ is as defined hereinbefore);

9) C_{1-3} alkylR³⁴ (wherein R³⁴ is as defined hereinbefore);

10) C_{3-3} alkenylR³⁴ (wherein R³⁴ is as defined hereinbefore);

11) C_{3-3} alkynylR³⁴ (wherein R³⁴ is as defined hereinbefore);

12) C_{1-3} alkylX⁷R³⁴ (wherein X⁷ and R³⁴ are as defined hereinbefore);

10 13) C_{4-3} alkenylX⁸R³⁴ (wherein X⁸ and R³⁴ are as defined hereinbefore);

14) C_{4-3} alkynylX⁹R³⁴ (wherein X⁹ and R³⁴ are as defined hereinbefore); and

15) C_{2-3} alkylX¹⁰ C_{1-2} alkylR³⁴ (wherein X¹⁰ and R³⁴ are as defined hereinbefore); and additional convenient values of R¹⁰ are:

16) R³³ (wherein R³³ is as defined hereinbefore); and

15 17) C_{2-3} alkylX¹⁰ C_{1-2} alkylR³³ (wherein X¹⁰ and R³³ are as defined hereinbefore)].

Advantageously R³ represents hydroxy, halogeno, nitro, trifluoromethyl, C_{1-3} alkyl, cyano, amino or R¹⁰X² [wherein X² is as hereinbefore defined and R¹⁰ is selected from one of the following fifteen groups:

1) C_{1-4} alkyl which may be unsubstituted or substituted with one or more fluorine atoms, or C_{2-4} alkyl which may be unsubstituted or substituted with one or two groups selected from

20 hydroxy and amino;

2) C_{2-3} alkylX³COR¹⁶ (wherein X³ is as hereinbefore defined and R¹⁶ represents -NR¹⁸R¹⁹- or -OR²⁰- (wherein R¹⁸, R¹⁹ and R²⁰ which may be the same or different are each C_{1-2} alkyl or C_{1-2} alkoxyethyl));

25 3) C_{2-4} alkylX⁴R²¹ (wherein X⁴ is as hereinbefore defined and R²¹ is a group selected from C_{1-3} alkyl, cyclopentyl, cyclohexyl, pyrrolidinyl and piperidinyl which group is linked to X⁴ through a carbon atom and which C_{1-3} alkyl group may bear one or two substituents selected from oxo, hydroxy, halogeno and C_{1-2} alkoxy and which cyclopentyl, cyclohexyl, pyrrolidinyl or piperidinyl group may carry one substituent selected from oxo, hydroxy, halogeno, C_{1-2} alkyl, C_{1-2} hydroxyalkyl and C_{1-2} alkoxy);

30

- 4) $C_{2,3}\text{alkyl}X^5C_{2,3}\text{alkyl}X^6R^{27}$ (wherein X^5 and X^6 are as hereinbefore defined and R^{27} represents hydrogen or $C_{1,3}\text{alkyl}$);
- 5) $C_{1,4}\text{alkyl}R^{62}$ (wherein R^{62} is a group selected from pyrrolidinyl, piperazinyl, piperidinyl, 1,3-dioxolan-2-yl, 1,3-dioxan-2-yl, 1,3-dithiolan-2-yl and 1,3-dithian-2-yl, which group is linked to $C_{1,4}\text{alkyl}$ through a carbon atom and which group may carry one or two substituents selected from oxo, hydroxy, halogeno, $C_{1,3}\text{alkyl}$, $C_{1,2}\text{hydroxyalkyl}$ and $C_{1,2}\text{alkoxy}$) or $C_{2,4}\text{alkyl}R^{63}$ (wherein R^{63} is a group selected from morpholino, thiomorpholino, pyrrolidin-1-yl, piperazin-1-yl and piperidino which group may carry one or two substituents selected from oxo, hydroxy, halogeno, $C_{1,2}\text{alkyl}$, $C_{1,2}\text{hydroxyalkyl}$ and $C_{1,2}\text{alkoxy}$);
- 10 6) $C_{3,4}\text{alkenyl}R^{64}$ (wherein R^{64} represents R^{62} or R^{63} as defined hereinbefore);
- 7) $C_{3,4}\text{alkynyl}R^{64}$ (wherein R^{64} represents R^{62} or R^{63} as defined hereinbefore);
- 8) R^{34} (wherein R^{34} is as defined hereinbefore);
- 9) $C_{1,4}\text{alkyl}R^{34}$ (wherein R^{34} is as defined hereinbefore);
- 10) $1-R^{34}\text{prop-1-en-3-yl}$ or $1-R^{34}\text{but-2-en-4-yl}$ (wherein R^{34} is as defined hereinbefore with the proviso that when R^{10} is $1-R^{34}\text{prop-1-en-3-yl}$, R^{34} is linked to the alkenyl group via a carbon atom);
- 15 11) $1-R^{34}\text{prop-1-yn-3-yl}$ or $1-R^{34}\text{but-2-yn-4-yl}$ (wherein R^{34} is as defined hereinbefore with the proviso that when R^{10} is $1-R^{34}\text{prop-1-yn-3-yl}$, R^{34} is linked to the alkynyl group via a carbon atom);
- 20 12) $C_{1,3}\text{alkyl}X^7R^{34}$ (wherein X^7 and R^{34} are as defined hereinbefore);
- 13) $1-(R^{34}X^8)\text{but-2-en-4-yl}$ (wherein X^8 and R^{34} are as defined hereinbefore);
- 14) $1-(R^{34}X^9)\text{but-2-yn-4-yl}$ (wherein X^9 and R^{34} are as defined hereinbefore); and
- 15) $C_{2,3}\text{alkyl}X^{10}C_{1,2}\text{alkyl}R^{34}$ (wherein X^{10} and R^{34} are as defined hereinbefore); and additional advantageous values of R^{10} are:
- 25 16) R^{33} (wherein R^{33} is as defined hereinbefore); and
- 17) $C_{2,3}\text{alkyl}X^{10}C_{1,2}\text{alkyl}R^{33}$ (wherein X^{10} and R^{33} are as defined hereinbefore)].

Preferably R^3 represents hydroxy, halogeno, nitro, trifluoromethyl, $C_{1,3}\text{alkyl}$, cyano, amino or $R^{10}X^2$ [wherein X^2 is as hereinbefore defined and R^{10} is selected from one of the following thirteen groups:

- 1) $C_{1,3}$ alkyl which may be unsubstituted or substituted with one or more fluorine atoms, or $C_{2,3}$ alkyl which may be unsubstituted or substituted with one or two groups selected from hydroxy and amino;
- 2) 2-(3,3-dimethylureido)ethyl, 3-(3,3-dimethylureido)propyl, 2-(3-methylureido)ethyl, 3-(3-methylureido)propyl, 2-ureidoethyl, 3-ureidopropyl, 2-(N,N-dimethylcarbamoyloxy)ethyl, 3-(N,N-dimethylcarbamoyloxy)propyl, 2-(N-methylcarbamoyloxy)ethyl, 3-(N-methylcarbamoyloxy)propyl, 2-(carbamoyloxy)ethyl, 3-(carbamoyloxy)propyl;
- 3) $C_{2,3}$ alkyl X^4R^{21} (wherein X^4 is as defined hereinbefore and R^{21} is a group selected from $C_{1,2}$ alkyl, cyclopentyl, cyclohexyl, pyrrolidinyl and piperidinyl which group is linked to X^4 through a carbon atom and which $C_{1,2}$ alkyl group may bear one or two substituents selected from hydroxy, halogeno and $C_{1,2}$ alkoxy and which cyclopentyl, cyclohexyl, pyrrolidinyl or piperidinyl group may carry one substituent selected from oxo, hydroxy, halogeno, $C_{1,2}$ alkyl, $C_{1,2}$ hydroxyalkyl and $C_{1,2}$ alkoxy);
- 4) $C_{2,3}$ alkyl $X^5C_{2,3}$ alkyl X^6R^{27} (wherein X^5 and X^6 are as hereinbefore defined and R^{27} represents hydrogen or $C_{1,2}$ alkyl);
- 5) $C_{1,2}$ alkyl R^{62} (wherein R^{62} is a group selected from pyrrolidinyl, piperazinyl, piperidinyl, 1,3-dioxolan-2-yl, 1,3-dioxan-2-yl, 1,3-dithiolan-2-yl and 1,3-dithian-2-yl, which group is linked to $C_{1,2}$ alkyl through a carbon atom and which group may carry one substituent selected from oxo, hydroxy, halogeno, $C_{1,2}$ alkyl, $C_{1,2}$ hydroxyalkyl and $C_{1,2}$ alkoxy) or $C_{2,3}$ alkyl R^{63} (wherein R^{63} is a group selected from morpholino, thiomorpholino, piperidino, piperazin-1-yl and pyrrolidin-1-yl which group may carry one substituent selected from oxo, hydroxy, halogeno, $C_{1,2}$ alkyl, $C_{1,2}$ hydroxyalkyl and $C_{1,2}$ alkoxy);
- 6) R^{34} (wherein R^{34} is as defined hereinbefore);
- 7) $C_{1,4}$ alkyl R^{34} (wherein R^{34} is as defined hereinbefore);
- 8) 1- R^{34} but-2-en-4-yl (wherein R^{34} is as defined hereinbefore);
- 9) 1- R^{34} but-2-yn-4-yl (wherein R^{34} is as defined hereinbefore);
- 10) $C_{1,3}$ alkyl X^7R^{34} (wherein X^7 and R^{34} are as defined hereinbefore);
- 11) 1-($R^{34}X^8$)but-2-en-4-yl (wherein X^8 and R^{34} are as defined hereinbefore);
- 12) 1-($R^{34}X^9$)but-2-yn-4-yl (wherein X^9 and R^{34} are as defined hereinbefore); and
- 13) ethyl X^{10} methyl R^{34} (wherein X^{10} and R^{34} are as defined hereinbefore); and additional preferred values of R^{10} are:


14) R^{33} (wherein R^{33} is as defined hereinbefore); and

15) ethyl X^{10} methyl R^{33} (wherein X^{10} and R^{33} are as defined hereinbefore)].

More preferably R^3 represents hydroxy, C_{1-3} alkyl, amino or $R^{10}X^2$ [wherein X^2 is as hereinbefore defined and R^{10} represents methyl, ethyl, trifluoromethyl, 2,2,2-trifluoroethyl, 2-hydroxyethyl, 3-hydroxypropyl, 2-methoxyethyl, 3-methoxypropyl, 2-(methylsulphinyl)ethyl, 2-(methylsulphonyl)ethyl, 2-(N,N-dimethylsulphamoyl)ethyl, 2-(N-methylsulphamoyl)ethyl, 2-sulphamoyl, 2-(N,N-dimethylamino)ethyl, 3-(N,N-dimethylamino)propyl, 2-morpholinoethyl, 3-morpholinopropyl, 2-piperidinoethyl, 3-piperidinopropyl, 2-(piperazin-1-yl)ethyl, 3-(piperazin-1-yl)propyl, 2-(pyrrolidin-1-yl)ethyl, 3-(pyrrolidin-1-yl)propyl, (1,3-dioxolan-2-yl)methyl, 2-(1,3-dioxolan-2-yl)ethyl, 2-(2-methoxyethylamino)ethyl, 2-(2-hydroxyethylamino)ethyl, 3-(2-methoxyethylamino)propyl, 3-(2-hydroxyethylamino)propyl, 2-methylthiazol-4-ylmethyl, 2-acetamidothiazol-4-ylmethyl, 1-methylimidazol-2-ylmethyl, 2-(imidazol-1-yl)ethyl, 2-(1,2,3-triazol-1-yl)ethyl, 2-(1,2,3-triazol-2-yl)ethyl, 2-(1,2,4-triazol-1-yl)ethyl, 2-(1,2,4-triazol-4-yl)ethyl, 4-pyridylmethyl, 2-(4-pyridyl)ethyl, 3-(4-pyridyl)propyl, 2-(4-pyridyloxy)ethyl, 2-(4-pyridylamino)ethyl, 2-(4-oxo-1,4-dihydro-1-pyridyl)ethyl, 2-thiomorpholinoethyl, 3-thiomorpholinopropyl, 2-(2-methoxyethoxy)ethyl, 2-(4-methylpiperazin-1-yl)ethyl or 3-(4-methylpiperazin-1-yl)propyl and additional more preferred values of R^{10} are 3-(methylsulphinyl)propyl, 3-(methylsulphonyl)propyl, 2-(2-methylimidazol-1-yl)ethyl, 2-(5-methyl-1,2,4-triazol-1-yl)ethyl, morpholino, 2-(N-(1-methylimidazol-4-ylsulphonyl)-N-methyl)amino)ethyl, 2-(N-(3-morpholinopropylsulphonyl)-N-methyl)amino)ethyl, 2-(N-methyl-N-4-pyridyl)amino)ethyl and 3-(4-oxidomorpholino)propyl].

Where one of the R^3 substituents is $R^{10}X^2$ the substituent $R^{10}X^2$ is preferably at the 6 or 7-position of the quinazoline ring, more preferably at the 7-position of the quinazoline ring.

25 In a particular aspect of the current invention there are provided compounds of the formula Ia:



(Ia)

[wherein:

R^1 , R^2 and n are as defined hereinbefore;

R^{3a} represents hydrogen, hydroxy, methoxy, amino, nitro or halogeno;

R^{4a} represents hydrogen, hydroxy, halogeno, cyano, nitro, trifluoromethyl, C_{1-3} alkyl, C_{1-3} alkoxy, C_{1-3} alkylthio, or $-NR^{6a}R^{7a}$ - (wherein R^{6a} and R^{7a} , which may be the same or different,

each represents hydrogen or C_{1-3} alkyl), and an additional value of R^{4a} is $R^{8a}(CH_2)_aX^{2a}$

(wherein R^{8a} is a 5 or 6 membered saturated heterocyclic group with one or two heteroatoms, selected independently from O, S and N, which heterocyclic group may bear one or two substituents selected from oxo, hydroxy, halogeno, C_{1-4} alkyl, C_{1-4} hydroxyalkyl and C_{1-4} alkoxy,

z_a is an integer from 0 to 4 and X^{2a} represents a direct bond, $-O-$, $-CH_2-$, $-S-$, $-SO-$, $-SO_2-$, $-NR^{9a}CO-$, $-CONR^{10a}-$, $-SO_2NR^{11a}-$, $-NR^{12a}SO_2-$ or $-NR^{13a}$ (wherein R^{9a} , R^{10a} , R^{11a} , R^{12a} and R^{13a} each independently represents hydrogen, C_{1-3} alkyl or C_{1-3} alkoxy C_{2-3} alkyl));

X^{1a} represents $-O-$, $-CH_2-$, $-S-$, $-SO-$, $-SO_2-$, $-NR^{14a}CO-$, $-CONR^{15a}-$, $-SO_2NR^{16a}-$, $-NR^{17a}SO_2-$ or $-NR^{18a}$ (wherein R^{14a} , R^{15a} , R^{16a} , R^{17a} and R^{18a} each independently represents hydrogen, C_{1-3} alkyl or C_{1-3} alkoxy C_{2-3} alkyl) or X^{1a} represents a direct bond;

R^{5a} is selected from one of the following fifteen groups:

1) hydrogen or C_{1-3} alkyl which may be unsubstituted or which may be substituted with one or more groups selected from hydroxy, fluoro and amino;

2) C_{1-3} alkyl $X^{3a}COR^{19a}$ (wherein X^{3a} represents $-O-$ or $-NR^{20a}$ (in which R^{20a} represents

hydrogen, C_{1-3} alkyl or C_{1-3} alkoxy C_{2-3} alkyl) and R^{19a} represents C_{1-3} alkyl, $-NR^{21a}R^{22a}$ or -

- OR^{23a}- (wherein R^{21a}, R^{22a} and R^{23a} which may be the same or different each represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl));
- 3) C₁₋₃alkylX^{4a}R^{24a} (wherein X^{4a} represents -O-, -S-, -SO-, -SO₂-, -OCO-, -NR^{25a}CO-, -CONR^{26a}-, -SO₂NR^{27a}-, -NR^{28a}SO₂- or -NR^{29a}- (wherein R^{25a}, R^{26a}, R^{27a}, R^{28a} and R^{29a} each independently represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R^{24a} represents hydrogen, C₁₋₃alkyl, cyclopentyl, cyclohexyl or a 5 or 6 membered saturated heterocyclic group with one or two heteroatoms, selected independently from O, S and N, which C₁₋₃alkyl group may bear one or two substituents selected from oxo, hydroxy, halogeno and C₁₋₄alkoxy and which cyclic group may bear one or two substituents selected from oxo, hydroxy, halogeno, C₁₋₄alkyl, C₁₋₄hydroxyalkyl and C₁₋₄alkoxy);
- 4) C₁₋₃alkylX^{5a}C₁₋₃alkylX^{6a}R^{30a} (wherein X^{5a} and X^{6a} which may be the same or different are each -O-, -S-, -SO-, -SO₂-, -NR^{31a}CO-, -CONR^{32a}-, -SO₂NR^{33a}-, -NR^{34a}SO₂- or -NR^{35a}- (wherein R^{31a}, R^{32a}, R^{33a}, R^{34a} and R^{35a} each independently represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R^{30a} represents hydrogen or C₁₋₃alkyl);
- 5) C₁₋₃alkylR^{36a} (wherein R^{36a} is a 5 or 6 membered saturated heterocyclic group with one or two heteroatoms, selected independently from O, S and N, which heterocyclic group may bear one or two substituents selected from oxo, hydroxy, halogeno, C₁₋₄alkyl, C₁₋₄hydroxyalkyl and C₁₋₄alkoxy);
- 6) C₂₋₃alkenylR^{36a} (wherein R^{36a} is as defined hereinbefore);
- 7) C₂₋₃alkynylR^{36a} (wherein R^{36a} is as defined hereinbefore);
- 8) R^{37a} (wherein R^{37a} represents a pyridone group, a phenyl group or a 5 or 6-membered aromatic heterocyclic group with 1 to 3 heteroatoms selected from O, N and S, which pyridone, phenyl or heterocyclic group may carry up to 5 substituents selected from halogeno, amino, C₁₋₄alkyl, C₁₋₄alkoxy, C₁₋₄hydroxyalkyl, C₁₋₄aminoalkyl, C₁₋₄alkylamino, C₁₋₄hydroxyalkoxy, carboxy, cyano, -CONR^{38a}R^{39a} and -NR^{40a}COR^{41a} (wherein R^{38a}, R^{39a}, R^{40a} and R^{41a}, which may be the same or different, each represents hydrogen, C₁₋₄alkyl or C₁₋₃alkoxyC₂₋₃alkyl)) and an additional possible substituent of R^{37a} is hydroxy;
- 9) C₁₋₃alkylR^{37a} (wherein R^{37a} is as defined hereinbefore);
- 10) C₂₋₃alkenylR^{37a} (wherein R^{37a} is as defined hereinbefore);
- 11) C₂₋₃alkynylR^{37a} (wherein R^{37a} is as defined hereinbefore);

- 12) $C_{1,3}\text{alkyl}X^{7a}R^{37a}$ (wherein X^{7a} represents -O-, -S-, -SO-, -SO₂-, -NR^{42a}CO-, -CONR^{43a}-, -SO₂NR^{44a}-, -NR^{45a}SO₂- or -NR^{46a}- (wherein R^{42a}, R^{43a}, R^{44a}, R^{45a} and R^{46a} each independently represents hydrogen, C_{1,3}alkyl or C_{1,3}alkoxyC_{2,3}alkyl) and R^{37a} is as defined hereinbefore);
- 13) $C_{2,3}\text{alkenyl}X^{8a}R^{37a}$ (wherein X^{8a} represents -O-, -S-, -SO-, -SO₂-, -NR^{47a}CO-, -CONR^{48a}-, -SO₂NR^{49a}-, -NR^{50a}SO₂- or -NR^{51a}- (wherein R^{47a}, R^{48a}, R^{49a}, R^{50a} and R^{51a} each independently represents hydrogen, C_{1,3}alkyl or C_{1,3}alkoxyC_{2,3}alkyl) and R^{37a} is as defined hereinbefore);
- 14) $C_{2,3}\text{alkynyl}X^{9a}R^{37a}$ (wherein X^{9a} represents -O-, -S-, -SO-, -SO₂-, -NR^{52a}CO-, -CONR^{53a}-, -SO₂NR^{54a}-, -NR^{55a}SO₂- or -NR^{56a}- (wherein R^{52a}, R^{53a}, R^{54a}, R^{55a} and R^{56a} each independently represents hydrogen, C_{1,3}alkyl or C_{1,3}alkoxyC_{2,3}alkyl) and R^{37a} is as defined hereinbefore); and
- 15) $C_{1,3}\text{alkyl}X^{10a}C_{1,3}\text{alkyl}R^{37a}$ (wherein X^{10a} represents -O-, -S-, -SO-, -SO₂-, -NR^{57a}CO-, -CONR^{58a}-, -SO₂NR^{59a}-, -NR^{60a}SO₂- or -NR^{61a}- (wherein R^{57a}, R^{58a}, R^{59a}, R^{60a} and R^{61a} each independently represents hydrogen, C_{1,3}alkyl or C_{1,3}alkoxyC_{2,3}alkyl) and R^{37a} is as defined hereinbefore) and additional groups of values of R^{5a} are:
- 16) R^{36a} (wherein R^{36a} is as defined hereinbefore); and
- 17) $C_{1,3}\text{alkyl}X^{10a}C_{1,3}\text{alkyl}R^{36a}$ (wherein X^{10a} and R^{36a} are as defined hereinbefore)];
- and salts thereof.

Preferably R^{3a} represents hydrogen, amino, nitro or halogeno, but especially hydrogen.

Advantageously X^{2a} represents -O-, -S-, -NR^{9a}CO-, -NR^{12a}SO₂- or -NR^{13a}- (wherein R^{9a}, R^{12a} and R^{13a} each independently represents hydrogen, C_{1,2}alkyl or C_{1,2}alkoxyethyl).

- 20 Preferably X^{2a} represents -O-, -S-, -NR^{9a}CO-, -NR^{12a}SO₂- (wherein R^{9a} and R^{12a} each independently represents hydrogen or C_{1,2}alkyl) or NH.

More preferably X^{2a} represents -O-, -S-, -NR^{9a}CO- (wherein R^{9a} represents hydrogen or C_{1,2}alkyl) or NH.

- 25 Particularly X^{2a} represents -O- or -NR^{9a}CO- (wherein R^{9a} represents hydrogen or C_{1,2}alkyl), more particularly -O- or -NHCO-, especially -O-.

Preferably za is an integer from 1 to 3.

Preferably R^{8a} is a group selected from pyrrolidinyl, piperazinyl, piperidinyl, morpholino and thiomorpholino which group may carry one or two substituents selected from oxo, hydroxy, halogeno, C_{1,2}alkyl, C_{1,2}hydroxyalkyl and C_{1,2}alkoxy.

Advantageously R^{4a} represents hydrogen, hydroxy, cyano, nitro, trifluoromethyl, C_{1-3} alkyl, C_{1-3} alkoxy or amino and an additional advantageous value of R^{4a} is $R^{8a}(CH_2)_{za}X^{2a}$ (wherein R^{8a} , X^{2a} and za are as defined hereinbefore).

Preferably R^{4a} is hydrogen, hydroxy, cyano, nitro, trifluoromethyl, methyl, ethyl, methoxy or ethoxy, and an additional preferred value of R^{4a} is $R^{8a}(CH_2)_{za}X^{2a}$ (wherein R^{8a} , X^{2a} and za are as defined hereinbefore).

More preferably R^{4a} is hydrogen, hydroxy, cyano, nitro, trifluoromethyl, methyl, ethyl, methoxy, ethoxy or $R^{8a}(CH_2)_{za}X^{2a}$ (wherein R^{8a} is a group selected from pyrrolidinyl, piperazinyl, piperidinyl, morpholino and thiomorpholino which group may carry one or two substituents selected from oxo, hydroxy, halogeno, C_{1-2} alkyl, C_{1-2} hydroxyalkyl and C_{1-2} alkoxy, X^{2a} is -O-, -S-, -NR^{9a}CO-, -NR^{12a}SO₂- (wherein R^{9a} and R^{12a} each independently represents hydrogen or C_{1-2} alkyl) or NH and za is an integer from 1 to 3).

Particularly R^{4a} represents hydrogen, hydroxy, cyano, nitro, trifluoromethyl, methyl or methoxy and an additional more preferred value of R^{4a} is $R^{8a}(CH_2)_{za}X^{2a}$ (wherein R^{8a} , X^{2a} and za are as defined hereinbefore).

More particularly R^{4a} represents hydrogen or methoxy.

R^{4a} especially represents methoxy.

Advantageously X^{1a} represents -O-, -S-, -NR^{14a}CO-, -NR^{17a}SO₂- or -NR^{18a}- (wherein R^{14a} , R^{17a} and R^{18a} each independently represents hydrogen, C_{1-2} alkyl or C_{1-2} alkoxyethyl).

Preferably X^{1a} represents -O-, -S-, -NR^{14a}CO-, -NR^{17a}SO₂- (wherein R^{14a} and R^{17a} each independently represents hydrogen or C_{1-2} alkyl) or NH.

More preferably X^{1a} represents -O-, -S-, -NR^{14a}CO- (wherein R^{14a} represents hydrogen or C_{1-2} alkyl) or NH.

Particularly X^{1a} represents -O- or -NR^{14a}CO- (wherein R^{14a} represents hydrogen or C_{1-2} alkyl), more particularly -O- or -NHCO-, especially -O-.

Advantageously X^{3a} represents -O- or NR^{20a} (wherein R^{20a} represents hydrogen, C_{1-3} alkyl or C_{1-2} alkoxyethyl).

Advantageously X^{4a} represents -O-, -S-, -SO-, -SO₂-, -NR^{25a}CO-, -NR^{28a}SO₂- or -NR^{29a}- (wherein R^{25a} , R^{28a} and R^{29a} each independently represents hydrogen, C_{1-2} alkyl or C_{1-2} alkoxyethyl).

- Preferably X^{4a} represents -O-, -S-, -SO-, -SO₂- or -NR^{29a}- (wherein R^{29a} represents hydrogen, C₁₋₂alkyl or C₁₋₂alkoxyethyl).
- More preferably X^{4a} represents -O- or -NR^{29a}- (wherein R^{29a} represents hydrogen or C₁₋₂alkyl).
- Advantageously X^{5a} and X^{6a} which may be the same or different each represents -O-, -S-, -SO-, -SO₂- or -NR^{35a}- (wherein R^{35a} represents hydrogen, C₁₋₃alkyl or C₁₋₂alkoxyethyl).
- 5 Preferably X^{5a} and X^{6a} which may be the same or different each represents -O-, -S- or -NR^{35a}- (wherein R^{35a} represents hydrogen, C₁₋₂alkyl or C₁₋₂alkoxyethyl).
- More preferably X^{5a} and X^{6a} which may be the same or different each represents -O- or -NH-.
- Advantageously X^{7a} represents -O-, -S- or -NR^{46a}- (wherein R^{46a} represents hydrogen, C₁₋₂alkyl or C₁₋₂alkoxyethyl).
- 10 Preferably X^{7a} represents -O- or -NR^{46a}- (wherein R^{46a} represents hydrogen or C₁₋₂alkyl).
- Advantageously X^{8a} represents -O-, -S- or -NR^{51a}- (wherein R^{51a} represents hydrogen, C₁₋₂alkyl or C₁₋₂alkoxyethyl).
- Preferably X^{8a} represents -O- or -NR^{51a}- (wherein R^{51a} represents hydrogen or C₁₋₂alkyl).
- 15 Advantageously X^{9a} represents -O-, -S- or -NR^{56a}- (wherein R^{56a} represents hydrogen, C₁₋₂alkyl or C₁₋₂alkoxyethyl).
- Preferably X^{9a} represents -O- or -NR^{56a}- (wherein R^{56a} represents hydrogen or C₁₋₂alkyl).
- Advantageously X^{10a} represents -O-, -S- or -NR^{61a}- (wherein R^{61a} represents hydrogen, C₁₋₂alkyl or C₁₋₂alkoxyethyl).
- 20 Preferably X^{10a} represents -O- or -NR^{61a}- (wherein R^{61a} represents hydrogen or C₁₋₂alkyl).
- R^{36a} is preferably pyrrolidinyl, piperazinyl, piperidinyl, morpholino or thiomorpholino which group may carry one or two substituents selected from oxo, hydroxy, halogeno, C₁₋₂alkyl, C₁₋₂hydroxyalkyl and C₁₋₂alkoxy.
- R^{37a} preferably represents a pyridone group or a 5 or 6-membered aromatic heterocyclic group with 1 to 3 heteroatoms selected from O, N and S, which pyridone group or heterocyclic group may be substituted as hereinbefore defined.
- 25 Where R^{37a} is a 5 or 6-membered aromatic heterocyclic group, it preferably has 1 or 2 heteroatoms, selected from O, N and S, of which more preferably one is N, and may be substituted as hereinbefore defined.
- 30 R^{37a} is particularly a pyridone, pyridyl, imidazolyl, thiazolyl, thienyl, triazolyl or pyridazinyl group which group may be substituted as hereinbefore defined, more particularly a pyridone,

pyridyl, imidazolyl, thiazolyl or triazolyl group, especially a pyridone, pyridyl, imidazolyl or triazolyl group which group may be substituted as hereinbefore defined.

In one embodiment of the invention R^{37a} represents a pyridone, phenyl or 5 or 6-membered aromatic heterocyclic group with 1 to 3 heteroatoms selected from O, N and S, which group
5 may preferably carry up to 2 substituents, more preferably up to one substituent, selected from the group of substituents as hereinbefore defined.

In the definition of R^{37a} , conveniently substituents are selected from halogeno, C_{1-4} alkyl, C_{1-4} alkoxy and cyano, more conveniently substituents are selected from chloro, fluoro, methyl and ethyl.

10 Conveniently R^{5a} is selected from one of the following fifteen groups:

1) C_{1-3} alkyl which may be unsubstituted or substituted with one or more fluorine atoms, or C_{2-3} alkyl which may be unsubstituted or substituted with one or more groups selected from hydroxy and amino;

2) C_{2-3} alkyl X^{3a} COR^{19a} (wherein X^{3a} is as hereinbefore defined and R^{19a} represents C_{1-3} alkyl, -
15 NR^{21a}R^{22a}- or -OR^{23a}- (wherein R^{21a} , R^{22a} and R^{23a} which may be the same or different are each C_{1-2} alkyl or C_{1-2} alkoxyethyl));

3) C_{2-4} alkyl X^{4a} R^{24a} (wherein X^{4a} is as hereinbefore defined and R^{24a} represents hydrogen, C_{1-3} alkyl, cyclopentyl, cyclohexyl or a 5 or 6 membered saturated heterocyclic group with one or two heteroatoms, selected independently from O, S and N, which C_{1-3} alkyl group may bear
20 one or two substituents selected from oxo, hydroxy, halogeno and C_{1-3} alkoxy and which cyclic group may bear one or two substituents selected from oxo, hydroxy, halogeno, C_{1-4} alkyl, C_{1-4} hydroxyalkyl and C_{1-4} alkoxy);

4) C_{2-3} alkyl X^{5a} C_{2-3} alkyl X^{6a} R^{30a} (wherein X^{5a} and X^{6a} are as hereinbefore defined and R^{30a} represents hydrogen or C_{1-3} alkyl);

25 5) C_{1-3} alkylR^{62a} (wherein R^{62a} is a 5 or 6 membered saturated heterocyclic group with one or two heteroatoms, selected independently from O, S and N, which heterocyclic group is linked to C_{1-3} alkyl through a carbon atom and which heterocyclic group may bear one or two substituents selected from halogeno, C_{1-4} alkyl, C_{1-4} hydroxyalkyl and C_{1-4} alkoxy) or C_{2-3} alkylR^{63a} (wherein R^{63a} is a 5 or 6 membered saturated heterocyclic group with one or two
30 heteroatoms of which one is N and the other is selected independently from O, S and N, which heterocyclic group is linked to C_{2-3} alkyl through a nitrogen atom and which heterocyclic group

may bear one or two substituents selected from halogeno, C₁₋₄alkyl, C₁₋₄hydroxyalkyl and C₁₋₄alkoxy);

- 6) C₃₋₄alkenylR^{64a} (wherein R^{64a} represents R^{62a} or R^{63a} as defined hereinbefore);
 - 7) C₃₋₄alkynylR^{64a} (wherein R^{64a} represents R^{62a} or R^{63a} as defined hereinbefore);
 - 5 8) R^{37a} (wherein R^{37a} is as defined hereinbefore);
 - 9) C₁₋₃alkylR^{37a} (wherein R^{37a} is as defined hereinbefore);
 - 10) C₃₋₅alkenylR^{37a} (wherein R^{37a} is as defined hereinbefore);
 - 11) C₃₋₅alkynylR^{37a} (wherein R^{37a} is as defined hereinbefore);
 - 12) C₁₋₅alkylX^{7a}X^{37a} (wherein X^{7a} and R^{37a} are as defined hereinbefore);
 - 10 13) C₄₋₅alkenylX^{8a}R^{37a} (wherein X^{8a} and R^{37a} are as defined hereinbefore);
 - 14) C₄₋₅alkynylX^{9a}R^{37a} (wherein X^{9a} and R^{37a} are as defined hereinbefore); and
 - 15) C₂₋₃alkylX^{10a}C₁₋₂alkylR^{37a} (wherein X^{10a} and R^{37a} are as defined hereinbefore)
- and additional convenient groups of values of R^{5a} are:
- 16) R^{36a} (wherein R^{36a} is as defined hereinbefore); and
 - 15 17) C₂₋₃alkylX^{10a}C₁₋₂alkylR^{36a} (wherein X^{10a} and R^{36a} are as defined hereinbefore).

Advantageously R^{5a} is selected from one of the following fifteen groups:

- 1) C₁₋₄alkyl which may be unsubstituted or substituted with one or more fluorine atoms, or C₂₋₄alkyl which may be unsubstituted or substituted with one or two groups selected from hydroxy and amino;
- 20 2) C₂₋₃alkylX^{3a}COR^{19a} (wherein X^{3a} is as hereinbefore defined and R^{19a} represents -NR^{21a}R^{22a}- or -OR^{23a}- (wherein R^{21a}, R^{22a} and R^{23a} which may be the same or different are each C₁₋₂alkyl or C₁₋₂alkoxyethyl));
- 3) C₂₋₄alkylX^{4a}R^{24a} (wherein X^{4a} is as hereinbefore defined and R^{24a} is a group selected from C₁₋₃alkyl, cyclopentyl, cyclohexyl, pyrrolidinyl and piperidinyl which group is linked to X^{4a}
- 25 through a carbon atom and which C₁₋₃alkyl group may bear one or two substituents selected from oxo, hydroxy, halogeno and C₁₋₂alkoxy and which cyclopentyl, cyclohexyl, pyrrolidinyl or piperidinyl group may carry one substituent selected from oxo, hydroxy, halogeno, C₁₋₂alkyl, C₁₋₂hydroxyalkyl and C₁₋₂alkoxy);
- 4) C₂₋₃alkylX^{5a}C₂₋₃alkylX^{6a}R^{30a} (wherein X^{5a} and X^{6a} are as hereinbefore defined and R^{30a}
- 30 represents hydrogen or C₁₋₃alkyl);


- 5) C_{1-4} alkylR^{65a} wherein R^{65a} is a group selected from pyrrolidinyl, piperazinyl, piperidinyl, 1,3-dioxolan-2-yl, 1,3-dioxan-2-yl, 1,3-dithiolan-2-yl and 1,3-dithian-2-yl, which group is linked to C_{1-4} alkyl through a carbon atom and which group may carry one or two substituents selected from oxo, hydroxy, halogeno, C_{1-2} alkyl, C_{1-2} hydroxyalkyl and C_{1-2} alkoxy) or C_2 .
- 5 C_{1-4} alkylR^{66a} (wherein R^{66a} is a group selected from morpholino, thiomorpholino, pyrrolidin-1-yl, piperazin-1-yl and piperidino which group may carry one or two substituents selected from oxo, hydroxy, halogeno, C_{1-2} alkyl, C_{1-2} hydroxyalkyl and C_{1-2} alkoxy);
- 6) C_{3-4} alkenylR^{67a} (wherein R^{67a} represents R^{65a} or R^{66a} as defined hereinbefore);
- 7) C_{3-4} alkynylR^{67a} (wherein R^{67a} represents R^{65a} or R^{66a} as defined hereinbefore);
- 10 8) R^{37a} (wherein R^{37a} is as defined hereinbefore);
- 9) C_{1-4} alkylR^{37a} (wherein R^{37a} is as defined hereinbefore);
- 10) 1-R^{37a}prop-1-en-3-yl or 1-R^{37a}but-2-en-4-yl (wherein R^{37a} is as defined hereinbefore with the proviso that when R^{5a} is 1-R^{37a}prop-1-en-3-yl, R^{37a} is linked to the alkenyl group via a carbon atom);
- 15 11) 1-R^{37a}prop-1-yn-3-yl or 1-R^{37a}but-2-yn-4-yl (wherein R^{37a} is as defined hereinbefore with the proviso that when R^{5a} is 1-R^{37a}prop-1-yn-3-yl, R^{37a} is linked to the alkynyl group via a carbon atom);
- 12) C_{1-3} alkylX^{7a}R^{37a} (wherein X^{7a} and R^{37a} are as defined hereinbefore);
- 13) 1-(R^{37a}X^{8a})but-2-en-4-yl (wherein X^{8a} and R^{37a} are as defined hereinbefore);
- 20 14) 1-(R^{37a}X^{9a})but-2-yn-4-yl (wherein X^{9a} and R^{37a} are as defined hereinbefore); and
- 15) C_{2-3} alkylX^{10a} C_{1-2} alkylR^{37a} (wherein X^{10a} and R^{37a} are as defined hereinbefore)
- and additional advantageous groups of values of R^{5a} are:
- 16) R^{36a} (wherein R^{36a} is as defined hereinbefore); and
- 17) C_{2-3} alkylX^{10a} C_{1-2} alkylR^{36a} (wherein X^{10a} and R^{36a} are as defined hereinbefore).
- 25 Preferably R^{5a} is selected from one of the following thirteen groups:
- 1) C_{1-3} alkyl which may be unsubstituted or substituted with one or more fluorine atoms, or C_{2-3} alkyl which may be unsubstituted or substituted with one or two groups selected from hydroxy and amino;
- 2) 2-(3,3-dimethylureido)ethyl, 3-(3,3-dimethylureido)propyl, 2-(3-methylureido)ethyl, 3-(3-methylureido)propyl, 2-ureidoethyl, 3-ureidopropyl, 2-(N,N-dimethylcarbamoyloxy)ethyl, 3-
- 30

(N,N-dimethylcarbamoyloxy)propyl, 2-(N-methylcarbamoyloxy)ethyl, 3-(N-methylcarbamoyloxy)propyl, 2-(carbamoyloxy)ethyl, 3-(carbamoyloxy)propyl;

- 3) $C_{2,3}alkylX^{4a}R^{24a}$ (wherein X^{4a} is as defined hereinbefore and R^{24a} is a group selected from $C_{1,2}alkyl$, cyclopentyl, cyclohexyl, pyrrolidinyl and piperidinyl which group is linked to X^{4a} through a carbon atom and which $C_{1,2}alkyl$ group may bear one or two substituents selected from hydroxy, halogeno and $C_{1,2}alkoxy$ and which cyclopentyl, cyclohexyl, pyrrolidinyl or piperidinyl group may carry one substituent selected from oxo, hydroxy, halogeno, $C_{1,2}alkyl$, $C_{1,2}hydroxyalkyl$ and $C_{1,2}alkoxy$);
 - 4) $C_{2,3}alkylX^{5a}C_{2,3}alkylX^{6a}R^{30a}$ (wherein X^{5a} and X^{6a} are as hereinbefore defined and R^{30a} represents hydrogen or $C_{1,2}alkyl$);
 - 5) $C_{1,2}alkylR^{65a}$ (wherein R^{65a} is a group selected from pyrrolidinyl, piperazinyl, piperidinyl, 1,3-dioxolan-2-yl, 1,3-dioxan-2-yl, 1,3-dithiolan-2-yl and 1,3-dithian-2-yl, which group is linked to $C_{1,2}alkyl$ through a carbon atom and which group may carry one substituent selected from oxo, hydroxy, halogeno, $C_{1,2}alkyl$, $C_{1,2}hydroxyalkyl$ and $C_{1,2}alkoxy$) or $C_{2,3}alkylR^{66a}$ (wherein R^{66a} is a group selected from morpholino, thiomorpholino, piperidino, piperazin-1-yl and pyrrolidin-1-yl which group may carry one substituent selected from oxo, hydroxy, halogeno, $C_{1,2}alkyl$, $C_{1,2}hydroxyalkyl$ and $C_{1,2}alkoxy$);
 - 6) R^{37a} (wherein R^{37a} is as defined hereinbefore);
 - 7) $C_{1,4}alkylR^{37a}$ (wherein R^{37a} is as defined hereinbefore);
 - 8) $1-R^{37a}but-2-en-4-yl$ (wherein R^{37a} is as defined hereinbefore);
 - 9) $1-R^{37a}but-2-yn-4-yl$ (wherein R^{37a} is as defined hereinbefore);
 - 10) $C_{1,3}alkylX^{7a}R^{37a}$ (wherein X^{7a} and R^{37a} are as defined hereinbefore);
 - 11) $1-(R^{37a}X^{8a})but-2-en-4-yl$ (wherein X^{8a} and R^{37a} are as defined hereinbefore);
 - 12) $1-(R^{37a}X^{9a})but-2-yn-4-yl$ (wherein X^{9a} and R^{37a} are as defined hereinbefore); and
 - 13) $ethylX^{10a}methylR^{37a}$ (wherein X^{10a} and R^{37a} are as defined hereinbefore)
- and additional preferred groups of values of R^{5a} are:
- 14) R^{36a} (wherein R^{36a} is as defined hereinbefore); and
 - 15) $ethylX^{10a}methylR^{36a}$ (wherein X^{10a} and R^{36a} are as defined hereinbefore).
- More preferably R^{5a} represents methyl, ethyl, trifluoromethyl, 2,2,2-trifluoroethyl, 2-hydroxyethyl, 3-hydroxypropyl, 2-methoxyethyl, 3-methoxypropyl, 2-(methylsulphinyl)ethyl, 2-(methylsulphonyl)ethyl, 2-(N,N-dimethylsulphamoyl)ethyl, 2-(N-methylsulphamoyl)ethyl,

2-sulphamoyl ethyl, 2-(N,N-dimethylamino)ethyl, 3-(N,N-dimethylamino)propyl, 2-morpholinoethyl, 3-morpholinopropyl, 2-piperidinoethyl, 3-piperidinopropyl, 2-(piperazin-1-yl)ethyl, 3-(piperazin-1-yl)propyl, 2-(pyrrolidin-1-yl)ethyl, 3-(pyrrolidin-1-yl)propyl, (1,3-dioxolan-2-yl)methyl, 2-(1,3-dioxolan-2-yl)ethyl, 2-(2-methoxyethylamino)ethyl, 2-(2-hydroxyethylamino)ethyl, 3-(2-methoxyethylamino)propyl, 3-(2-hydroxyethylamino)propyl, 2-methylthiazol-4-ylmethyl, 2-acetamidothiazol-4-ylmethyl, 1-methylimidazol-2-ylmethyl, 2-(imidazol-1-yl)ethyl, 2-(1,2,3-triazol-1-yl)ethyl, 2-(1,2,3-triazol-2-yl)ethyl, 2-(1,2,4-triazol-1-yl)ethyl, 2-(1,2,4-triazol-4-yl)ethyl, 4-pyridylmethyl, 2-(4-pyridyl)ethyl, 3-(4-pyridyl)propyl 2-(4-pyridyloxy)ethyl, 2-(4-pyridylamino)ethyl, 2-(4-oxo-1,4-dihydro-1-pyridyl)ethyl, 2-thiomorpholinoethyl, 3-thiomorpholinopropyl, 2-(2-methoxyethoxy)ethyl, 2-(4-methylpiperazin-1-yl)ethyl or 3-(4-methylpiperazin-1-yl)propyl and additional more preferred values of R^{5a} are 3-(methylsulphinyl)propyl, 3-(methylsulphonyl)propyl, 2-(2-methylimidazol-1-yl)ethyl, 2-(5-methyl-1,2,4-triazol-1-yl)ethyl, morpholino, 2-(N-(1-methylimidazol-4-ylsulphonyl)-N-methyl)amino)ethyl, 2-(N-(3-morpholinopropylsulphonyl)-N-methyl)amino)ethyl, 2-(N-methyl-N-4-pyridyl)amino)ethyl and 3-(4-oxidomorpholino)propyl.

In a further aspect of the current invention there are provided compounds of the formula Ib:



(Ib)

[wherein:

R^{1b} represents hydrogen;

R^{2b} represents halogeno, trifluoromethyl, cyano, nitro, C₂₋₃alkanoyl, C₁₋₃alkoxycarbonyl, C₁₋₃alkylsulphinyl, C₁₋₃alkylsulphonyl, carbamoyl, N-C₁₋₃alkylcarbamoyl, N,N-di(C₁₋

- ,alkyl)carbamoyl, aminosulphonyl, N-C₁₋₃alkylaminosulphonyl, N,N-di(C₁₋₃alkyl)aminosulphonyl, or a group R^{6b}X^{2b} (wherein X^{2b} represents a direct bond, C₂₋₄alkanoyl, -CONR^{7b}R^{8b}-, -SO₂NR^{9b}R^{10b}- or -SO₂R^{11b}- (wherein R^{7b} and R^{9b}, each independently represents hydrogen and R^{8b}, R^{10b} and R^{11b} each independently represents C₁₋₃alkyl and wherein R^{6b} is linked to R^{8b}, R^{10b} or R^{11b}) and R^{6b} represents pyrrolidinyl, piperazinyl, piperidinyl, morpholino, thiomorpholino, imidazolyl or triazolyl which heterocyclic group is linked to X^{2b} via a nitrogen atom and which heterocyclic group may bear one or two substituents selected from hydroxy, halogeno, C₁₋₂alkyl, C₁₋₂alkoxy, C₁₋₂alkanoyloxy, trifluoromethyl, cyano, amino, nitro and C₁₋₄alkoxycarbonyl);
- 5 nb is an integer from 0 to 2;
 R^{3b} represents hydrogen;
 R^{4b} represents represents hydrogen, hydroxy, cyano, nitro, trifluoromethyl, methyl, methoxy or a group R^{12b}(CH₂)_{zb}X^{3b} (wherein R^{12b} represents a group selected from pyrrolidinyl, piperazinyl, piperidinyl, morpholino and thiomorpholino which group may carry one or two substituents selected from oxo, hydroxy, halogeno, C₁₋₂alkyl, C₁₋₂hydroxyalkyl and C₁₋₂alkoxy, zb is an integer from 1 to 3 and X^{3b} represents -O- or -NR^{13b}CO- (wherein R^{13b} represents hydrogen or C₁₋₂alkyl));
- 15 X^{1b} represents -O- or -NR^{14b}CO- (wherein R^{14b} represents hydrogen or C₁₋₂alkyl); and
 R^{5b} represents methyl, ethyl, trifluoromethyl, 2,2,2-trifluoroethyl, 2-hydroxyethyl, 3-hydroxypropyl, 2-methoxyethyl, 3-methoxypropyl, 2-(methylsulphinyl)ethyl, 2-(methylsulphonyl)ethyl, 2-(N,N-dimethylsulphamoyl)ethyl, 2-(N-methylsulphamoyl)ethyl, 2-sulphamoyl ethyl, 2-(N,N-dimethylamino)ethyl, 3-(N,N-dimethylamino)propyl, 2-morpholinoethyl, 3-morpholinopropyl, 2-piperidinoethyl, 3-piperidinopropyl, 2-(piperazin-1-yl)ethyl, 3-(piperazin-1-yl)propyl, 2-(pyrrolidin-1-yl)ethyl, 3-(pyrrolidin-1-yl)propyl, (1,3-dioxolan-2-yl)methyl, 2-(1,3-dioxolan-2-yl)ethyl, 2-(2-methoxyethylamino)ethyl, 2-(2-hydroxyethylamino)ethyl, 3-(2-methoxyethylamino)propyl, 3-(2-hydroxyethylamino)propyl, 2-methylthiazol-4-ylmethyl, 2-acetamidothiazol-4-ylmethyl, 1-methylimidazol-2-ylmethyl, 2-(imidazol-1-yl)ethyl, 2-(1,2,3-triazol-1-yl)ethyl, 2-(1,2,3-triazol-2-yl)ethyl, 2-(1,2,4-triazol-1-yl)ethyl, 2-(1,2,4-triazol-4-yl)ethyl, 4-pyridylmethyl, 2-(4-pyridyl)ethyl, 3-(4-pyridyl)propyl
- 25 2-(4-pyridyloxy)ethyl, 2-(4-pyridylamino)ethyl, 2-(4-oxo-1,4-dihydro-1-pyridyl)ethyl, 2-thiomorpholinoethyl, 3-thiomorpholinopropyl, 2-(2-methoxyethoxy)ethyl, 2-(4-
- 30

methylpiperazin-1-yl)ethyl, 3-(4-methylpiperazin-1-yl)propyl, 3-(methylsulphinyl)propyl, 3-(methylsulphonyl)propyl, 2-(2-methylimidazol-1-yl)ethyl, 2-(5-methyl-1,2,4-triazol-1-yl)ethyl, morpholino, 2-((N-(1-methylimidazol-4-ylsulphonyl)-N-methyl)amino)ethyl, 2-((N-(3-morpholinopropylsulphonyl)-N-methyl)amino)ethyl, 2-((N-methyl-N-4-pyridyl)amino)ethyl or 3-(4-oxidomorpholino)propyl;]

5 and salts thereof.

Preferred compounds are

- 7-methoxy-6-(3-morpholinopropoxy)-4-(oxindol-3-yl)quinazoline,
- 4-(5-cyanooxindol-3-yl)-6-methoxy-7-(2-methoxyethoxy)quinazoline,
- 10 7-(2-(imidazol-1-yl)ethoxy)-6-methoxy-4-(oxindol-3-yl)quinazoline,
- 4-(5-cyanooxindol-3-yl)-6-methoxy-7-(3-morpholinopropoxy)quinazoline,
- 4-(5-cyanooxindol-3-yl)-6-methoxy-7-(2-((N-methyl-N-(4-pyridyl)]amino)ethoxy)quinazoline,
- 6-methoxy-4-(5-methoxycarbonyloxindol-3-yl)-7-(3-morpholinopropoxy)quinazoline,
- 15 4-(6-fluorooxindol-3-yl)-7-(2-(imidazol-1-yl)ethoxy)-6-methoxyquinazoline,
- 4-(7-hydroxyoxindol-3-yl)-6-methoxy-7-(3-morpholinopropoxy)quinazoline,
- 6-methoxy-7-(2-methoxyethoxy)-4-(5-nitrooxindol-3-yl)-quinazoline,
- 6-methoxy-7-(3-morpholinopropoxy)-4-(5-nitrooxindol-3-yl)-quinazoline,
- 4-(5-acetamidooxindol-3-yl)-6-methoxy-7-(2-methoxyethoxy)quinazoline,
- 20 4-(5-acetyloxindol-3-yl)-6-methoxy-7-(2-methoxyethoxy)quinazoline,
- 6-methoxy-4-(5-methylsulphonyloxindol-3-yl)-7-(3-morpholinopropoxy)quinazoline,
- 6-methoxy-4-(5-methylsulphinylloxindol-3-yl)-7-(3-morpholinopropoxy)quinazoline,
- 7-(2-(imidazol-1-yl)ethoxy)-6-methoxy-4-(5-(2-morpholinoethylaminosulphonyl)oxindol-3-yl)quinazoline,
- 25 7-(2-methoxyethoxy)-4-(5-nitrooxindol-3-yl)quinazoline,
- 6-methoxy-7-(2-methoxyethoxy)-4-(6-trifluoromethyloxindol-3-yl)quinazoline,
- 4-(6-cyanooxindol-3-yl)-7-(3-morpholinopropoxy)quinazoline,
- 4-(6-cyanooxindol-3-yl)-6-methoxy-7-(3-morpholinopropoxy)quinazoline,
- 4-(6-bromooxindol-3-yl)-7-(3-morpholinopropoxy)quinazoline,
- 30 4-(5-aminosulphonyloxindol-3-yl)-6-methoxy-7-(3-morpholinopropoxy)quinazoline,
- 4-(oxindol-3-yl)-6-morpholinoquinazoline,

- 4-(5-cyanooxindol-3-yl)-6-methoxy-7-(2-(pyrrolidin-1-yl)ethoxy)quinazoline,
6-methoxy-4-(5-methylthiooxindol-3-yl)-7-(3-morpholinopropoxy)quinazoline,
6-methoxy-7-(2-methoxyethoxy)-4-(5-(2-morpholinoethylaminosulphonyl)oxindol-3-yl)quinazoline,
- 5 6,7-dimethoxy-4-(5-(2-morpholinoethylaminosulphonyl)oxindol-3-yl)quinazoline,
4-(5-cyanooxindol-3-yl)-7-(2-methoxyethoxy)quinazoline,
7-(3-morpholinopropoxy)-4-(5-nitrooxindol-3-yl)quinazoline,
4-(6-cyanooxindol-3-yl)-6-methoxy-7-(2-methoxyethoxy)quinazoline,
6-methoxy-7-(2-methoxyethoxy)-4-(5-methylaminosulphonyloxindol-3-yl)quinazoline,
- 10 4-(5-N,N-dimethylaminosulphonyloxindol-3-yl)-6-methoxy-7-(2-methoxyethoxy)quinazoline,
6-methoxy-4-(oxindol-3-yl)-7-(2-(pyrrolidin-1-yl)ethoxy)quinazoline,
4-(5-bromooxindol-3-yl)-6-methoxy-7-(3-morpholinopropoxy)quinazoline,
6-methoxy-4-(oxindol-3-yl)-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline,
7-(3-morpholinopropoxy)-4-(oxindol-3-yl)quinazoline,
- 15 6-methoxy-7-(2-([N-methyl-N-(3-morpholinopropylsulphonyl)]amino)ethoxy)-4-(oxindol-3-yl)quinazoline and
6-methoxy-7-(2-([N-methyl-N-(1-methyl-1H-imidazol-4-ylsulphonyl)]amino)ethoxy)-4-(oxindol-3-yl)quinazoline
and salts thereof, particularly hydrochloride salts thereof.
- 20 More preferred compounds are
6,7-dimethoxy-4-(6-fluorooxindol-3-yl)quinazoline,
6,7-dimethoxy-4-(oxindol-3-yl)quinazoline,
7-(2-methoxyethoxy)-4-(oxindol-3-yl)quinazoline,
4-(5-bromooxindol-3-yl)-6-methoxy-7-(2-methoxyethoxy)quinazoline,
- 25 4-(5-hydroxyoxindol-3-yl)-6-methoxy-7-(3-morpholinopropoxy)quinazoline,
6-methoxy-7-(3-morpholinopropoxy)-4-(6-trifluoromethyloxindol-3-yl)quinazoline,
4-(5-aminosulphonyloxindol-3-yl)-6-methoxy-7-(2-methoxyethoxy)quinazoline,
4-(5-cyanooxindol-3-yl)-7-(3-morpholinopropoxy)quinazoline,
6-methoxy-7-(2-(2-methoxyethoxy)ethoxy)-4-(oxindol-3-yl)quinazoline and
- 30 4-(5-aminosulphonyloxindol-3-yl)-6-methoxy-7-(2-(2-methoxyethoxy)ethoxy)quinazoline
and salts thereof, particularly hydrochloride salts thereof.

Especially preferred compounds are

- 4-(5-fluorooxindol-3-yl)-6-methoxy-7-(3-morpholinopropoxy)quinazoline,
4-(5-fluorooxindol-3-yl)-6-methoxy-7-(2-methoxyethoxy)quinazoline,
4-(5-cyanooxindol-3-yl)-7-(2-(imidazol-1-yl)ethoxy)-6-methoxyquinazoline and
5 7-(3-morpholinopropoxy)-4-(6-trifluoromethyloxindol-3-yl)quinazoline
and salts thereof, particularly hydrochloride salts thereof.

More especially preferred compounds are

- 4-(6-fluorooxindol-3-yl)-6-methoxy-7-(3-morpholinopropoxy)quinazoline,
6-methoxy-7-(2-methoxyethoxy)-4-(oxindol-3-yl)quinazoline and
10 6-methoxy-7-(3-morpholinopropoxy)-4-(oxindol-3-yl)quinazoline
and salts thereof, particularly hydrochloride salts thereof.

In another aspect of the present invention particularly preferred compounds by virtue of their good activity against VEGF receptor tyrosine kinase activity and their lack of significant activity against epidermal growth factor (EGF) receptor tyrosine kinase include:

- 15 6-methoxy-7-(2-methoxyethoxy)-4-(oxindol-3-yl)quinazoline,
6-methoxy-7-(3-morpholinopropoxy)-4-(oxindol-3-yl)quinazoline,
4-(5-fluorooxindol-3-yl)-6-methoxy-7-(3-morpholinopropoxy)quinazoline,
4-(5-bromooxindol-3-yl)-6-methoxy-7-(2-methoxyethoxy)quinazoline,
4-(5-fluorooxindol-3-yl)-6-methoxy-7-(2-methoxyethoxy)quinazoline,
20 4-(5-hydroxyoxindol-3-yl)-6-methoxy-7-(3-morpholinopropoxy)quinazoline,
6,7-dimethoxy-4-(6-fluorooxindol-3-yl)quinazoline,
4-(6-fluorooxindol-3-yl)-6-methoxy-7-(3-morpholinopropoxy)quinazoline,
7-(2-methoxyethoxy)-4-(oxindol-3-yl)quinazoline,
6,7-dimethoxy-4-(oxindol-3-yl)quinazoline,
25 and salts thereof, particularly hydrochloride salts thereof.

For the avoidance of doubt it is to be understood that where in this specification a group is qualified by 'hereinbefore defined' or 'defined hereinbefore' the said group encompasses the first occurring and broadest definition as well as each and all of the preferred definitions for that group.

- 30 In this specification unless stated otherwise the term "alkyl" includes both straight and branched chain alkyl groups but references to individual alkyl groups such as "propyl" are

specific for the straight chain version only. An analogous convention applies to other generic terms. Unless otherwise stated the term "alkyl" advantageously refers to chains with 1-6 carbon atoms, preferably 1-4 carbon atoms. The term "alkoxy" as used herein, unless stated otherwise includes "alkyl"-O- groups in which "alkyl" is as hereinbefore defined. The term

5 "aryl" as used herein unless stated otherwise includes reference to a C₆₋₁₀ aryl group which may, if desired, carry one or more substituents selected from halogeno, alkyl, alkoxy, nitro, trifluoromethyl and cyano, (wherein alkyl and alkoxy are as hereinbefore defined). The term "aryloxy" as used herein unless otherwise stated includes "aryl"-O-groups in which "aryl" is as hereinbefore defined. The term "sulphonyloxy" as used herein refers to alkylsulphonyloxy


10 and arylsulphonyloxy groups in which "alkyl" and "aryl" are as hereinbefore defined. The term "alkanoyl" as used herein unless otherwise stated includes formyl and alkylC=O groups in which "alkyl" is as defined hereinbefore, for example C₂alkanoyl is ethanoyl and refers to CH₃C=O, C₁alkanoyl is formyl and refers to CHO. In this specification unless stated otherwise the term "alkenyl" includes both straight and branched chain alkenyl groups but

15 references to individual alkenyl groups such as 2-butenyl are specific for the straight chain version only. Unless otherwise stated the term "alkenyl" advantageously refers to chains with 2-5 carbon atoms, preferably 3-4 carbon atoms. In this specification unless stated otherwise the term "alkynyl" includes both straight and branched chain alkynyl groups but references to individual alkynyl groups such as 2-butylnyl are specific for the straight chain version only.


20 Unless otherwise stated the term "alkynyl" advantageously refers to chains with 2-5 carbon atoms, preferably 3-4 carbon atoms.

Within the present invention it is to be understood that a compound of the formula I or a salt thereof may exhibit the phenomenon of tautomerism and that the formulae drawings within this specification can represent only one of the possible tautomeric forms. It is to be

25 understood that the invention encompasses any tautomeric form which inhibits VEGF receptor tyrosine kinase activity and which inhibits FGF R1 receptor tyrosine kinase activity and is not to be limited merely to any one tautomeric form utilised within the formulae drawings. In particular the oxindole group can exist in at least two forms giving compounds of formulae I and II:



(I)



(II)

(wherein R^1 , R^2 , R^3 , m and n are as hereinbefore defined). The formulae drawings within this specification can represent only one of the possible tautomeric forms and it is to be

5 understood that the specification encompasses all possible tautomeric forms of the compounds drawn not just those forms which it has been possible to show graphically herein.

It is also to be understood that certain compounds of the formula I and salts thereof can exist in solvated as well as unsolvated forms such as, for example, hydrated forms. It is to be understood that the invention encompasses all such solvated forms which inhibit VEGF
10 receptor tyrosine kinase activity and inhibit FGF R1 receptor tyrosine kinase activity.

For the avoidance of any doubt, it is to be understood that when X^2 is, for example, a group of formula $-NR^{11}CO-$, it is the nitrogen atom bearing the R^{11} group which is attached to the quinazoline ring and the carbonyl (CO) group is attached to R^{10} , whereas when X^2 is, for example, a group of formula $-CONR^{12}-$, it is the carbonyl group which is attached to the
15 quinazoline ring and the nitrogen atom bearing the R^{12} group is attached to R^{10} . A similar convention applies to the other two atom X^2 linking groups such as $-NR^{14}SO_2-$ and $-SO_2NR^{13}-$. When X^2 is $-NR^{15}-$ it is the nitrogen atom bearing the R^{15} group which is linked to the quinazoline ring and to R^{10} . An analogous convention applies to other groups. It is further to be understood that when X^2 represents $-NR^{15}-$ and R^{15} is $C_{1,3}alkoxyC_{2,3}alkyl$ it is the $C_{2,3}alkyl$
20 moiety which is linked to the nitrogen atom of X^2 and an analogous convention applies to other groups.

For the avoidance of any doubt, it is to be understood that in a compound of the formula I when R^{10} is, for example, a group of formula $C_{1,3}alkylX^{10}C_{1,3}alkylR^{34}$, it is the terminal $C_{1,3}alkyl$ moiety which is linked to X^2 , similarly when R^{10} is, for example, a group of
25 formula $C_{2,3}alkenylR^{34}$ it is the $C_{2,3}alkenyl$ moiety which is linked to X^2 and an analogous convention applies to other groups. When R^{10} is a group 1- $R^{34}prop-1-en-3-yl$ it is the first

carbon to which the group R^{34} is attached and it is the third carbon which is linked to X^2 and an analogous convention applies to other groups.

For the avoidance of any doubt, it is to be understood that when R^{34} carries a C_{1-4} aminoalkyl substituent it is the C_{1-4} alkyl moiety which is attached to R^{34} whereas when R^{34} carries a C_{1-4} alkylamino substituent it is the amino moiety which is attached to R^{34} and an analogous convention applies to other groups.

For the avoidance of any doubt when X^1 is C_{2-4} alkanoyl it is the carbonyl moiety which is linked to the benz ring of the oxindole group and it is the alkyl moiety which is linked to R^4 and an analogous convention applies to other groups.

10 The present invention relates to the compounds of formula I as hereinbefore defined as well as to the salts thereof. Salts for use in pharmaceutical compositions will be pharmaceutically acceptable salts, but other salts may be useful in the production of the compounds of formula I and their pharmaceutically acceptable salts. Pharmaceutically acceptable salts of the invention may, for example, include acid addition salts of the
15 compounds of formula I as hereinbefore defined which are sufficiently basic to form such salts. Such acid addition salts include for example salts with inorganic or organic acids affording pharmaceutically acceptable anions such as with hydrogen halides (especially hydrochloric or hydrobromic acid of which hydrochloric acid is particularly preferred) or with sulphuric or phosphoric acid, or with trifluoroacetic, citric or maleic acid. In addition where
20 the compounds of formula I are sufficiently acidic, pharmaceutically acceptable salts may be formed with an inorganic or organic base which affords a pharmaceutically acceptable cation. Such salts with inorganic or organic bases include for example an alkali metal salt, such as a sodium or potassium salt, an alkaline earth metal salt such as a calcium or magnesium salt, an ammonium salt or for example a salt with methylamine, dimethylamine, trimethylamine,
25 piperidine, morpholine or tris-(2-hydroxyethyl)amine.


A compound of the formula I, or salt thereof, and other compounds of the invention (as hereinafter defined) may be prepared by any process known to be applicable to the preparation of chemically-related compounds. Such processes include, for example, those illustrated in European Patent Applications Publication Nos. 0520722, 0566226, 0602851,
30 0635498 and 0636608. Such processes, are provided as a further feature of the invention and are as described hereinafter. Necessary starting materials may be obtained by standard

procedures of organic chemistry. The preparation of such starting materials is described within the accompanying non-limiting Examples. Alternatively necessary starting materials are obtainable by analogous procedures to those illustrated which are within the ordinary skill of an organic chemist.

5 Thus the following processes (a) to (h) and (i) to (viii) constitute further features of the present invention.


Synthesis of Compounds of Formula I

(a) Compounds of the formula I and salts thereof may be prepared by the reaction of a compound of the formula III:



(III)

(wherein R³ and m are as defined hereinbefore and L¹ is a displaceable moiety), with a
20 compound of the formula IV:




(IV)

(wherein R¹, R² and n are as defined hereinbefore) whereby to obtain compounds of the formula I and salts thereof. A convenient displaceable moiety L¹ is, for example, a halogeno, alkoxy (preferably C₁₋₄alkoxy), aryloxy or sulphonyloxy group, for example a chloro, bromo, methoxy, phenoxy, methanesulphonyloxy or toluene-4-sulphonyloxy group.
30

The reaction is advantageously effected in the presence of a base. Such a base is, for example, an organic amine base such as, for example, pyridine, 2,6-lutidine, collidine, 4-dimethylaminopyridine, triethylamine, morpholine, N-methylmorpholine or diazabicyclo[5.4.0]undec-7-ene, tetramethylguanidine or for example, an alkali metal or
5 alkaline earth metal carbonate or hydroxide, for example sodium carbonate, potassium carbonate, calcium carbonate, sodium hydroxide or potassium hydroxide. Alternatively such a base is, for example, an alkali metal hydride, for example sodium hydride, or an alkali metal or alkaline earth metal amide, for example sodium amide, sodium bis(trimethylsilyl)amide, potassium amide or potassium bis(trimethylsilyl)amide. The reaction is preferably effected in
10 the presence of an inert solvent or diluent, for example an ether such as tetrahydrofuran or 1,4-dioxan, an aromatic hydrocarbon solvent such as toluene, or a dipolar aprotic solvent such as N,N-dimethylformamide, N,N-dimethylacetamide, N-methylpyrrolidin-2-one or dimethylsulphoxide. The reaction is conveniently effected at a temperature in the range, for example, 10 to 150°C, preferably in the range 20 to 90°C.

15 When it is desired to obtain the acid salt, the free base may be treated with an acid such as a hydrogen halide, for example hydrogen chloride, sulphuric acid, a sulphonic acid, for example methane sulphonic acid, or a carboxylic acid, for example acetic or citric acid, using a conventional procedure.

(b) A compound of the formula I and salts thereof can be prepared by the deprotection of
20 a compound of formula V:



(V)

5

(wherein R^2 , R^3 , n and m are as hereinbefore defined and P^1 represents a protecting group).

The choice of indole protecting group P^1 is within the standard knowledge of an organic chemist, for example those included in standard texts such as "Protective Groups in Organic Synthesis" T.W. Greene and R.G.M. Wuts, 2nd Ed. Wiley 1991, including N-sulphonyl derivatives (for example, p-toluenesulphonyl), carbamates (for example, t-butyl carbonyl), N-alkyl derivatives (for example, 2-chloroethyl, benzyl) and particularly amino acetal derivatives (for example, diethoxymethyl and benzyloxymethyl). The removal of such a protecting group may be effected by any of the procedures known for such a transformation, including those reaction conditions indicated in standard texts such as that indicated

10

15

hereinbefore, or by a related procedure. The reaction conditions preferably being such that the unprotected derivative is produced without unwanted reactions at other sites within the starting or product compounds. For example, where the protecting group P^1 is diethoxymethyl, the transformation may conveniently be effected by treatment of the oxindole derivative with an acid (as defined hereinbefore in process (a)) preferably in the presence of a

20


protic solvent or co-solvent such as water or an alcohol, for example methanol or ethanol. Such a reaction can be effected in the presence of an additional inert solvent or diluent (as defined hereinbefore in process (a)) advantageously at a temperature in the range 0 to 100°C, conveniently in the range 20 to 90°C.

(c) Compounds of the formula I and salts thereof wherein R^1 is hydrogen may also be

25

prepared by the reduction and cyclisation of a compound of formula VI:


- 34 -



(VI)

(wherein R^2 , R^3 , m and n are as defined hereinbefore and Y represents cyano, carboxy or C_{1-4} alkoxycarbonyl). The reduction of the nitro group may conveniently be effected as described in process (i) hereinafter. Where Y represents carboxy or C_{1-4} alkoxycarbonyl, the cyclisation to a compound of formula I occurs spontaneously after reduction of the nitro group or may be promoted, if necessary, by heating in an inert solvent or diluent such as xylene, toluene or *N,N*-dimethylformamide, optionally in the presence of an acid (as described hereinbefore in process (a)), advantageously at a temperature in the range of 20 to 150°C preferably in the range 20 to 100°C. Where Y represents cyano, the cyclisation to a compound of formula I can be effected in the presence of an acid (as described hereinbefore in process (a)) or a Lewis acid, such as a haloborane derivative for example boron trifluoride, and an alcohol or thiol, such as methanol, ethanol or *t*-butanethiol, in an inert solvent or diluent (as defined hereinbefore in process (a)) and at a temperature in the range -20 to 50°C preferably -5 to 10°C, followed by treatment with an aqueous acid, (as defined hereinbefore in process (a)), at a temperature in the range 20 to 150°C preferably in the range 20 to 100°C.

(d) Compounds of formula I and salts thereof wherein at least one of the R^2 groups is hydroxy may also be prepared by the deprotection of a compound of formula VII:




(VII)

(wherein R^1 , R^2 , R^3 , n and m are as defined hereinbefore, P^2 represents a phenolic hydroxy protecting group and p_1 is an integer from 1 to 4 equal to the number of protected hydroxy groups and such that $n-p_1$ is equal to the number of R^2 substituents which are not protected hydroxy). The choice of phenolic hydroxy protecting group P^2 is within the standard knowledge of an organic chemist, for example those included in standard texts such as "Protective Groups in Organic Synthesis" T.W. Greene and R.G.M. Wuts, 2nd Ed. Wiley 1991, including ethers (for example, methyl, methoxymethyl, allyl, benzyl and benzyl substituted with up to 2 substituents selected from C_{1-4} alkoxy and nitro), silyl ethers (for example, t-butyldiphenylsilyl and t-butyldimethylsilyl), esters (for example, acetate and benzoate) and carbonates (for example, methyl, benzyl and benzyl substituted with up to 2 substituents selected from C_{1-4} alkoxy and nitro). The removal of such a phenolic hydroxy protecting group may be effected by any of the procedures known for such a transformation, including those reaction conditions indicated in standard texts such as that indicated hereinbefore, or by a related procedure. The reaction conditions preferably being such that the hydroxy derivative is produced without unwanted reactions at other sites within the starting or product compounds. For example, where the protecting group P^2 is benzyl, the transformation may conveniently be effected by treatment of the oxindole derivative with an acid (as defined hereinbefore in process (a)) particularly trifluoroacetic acid, preferably in the presence of an ether or thioether such as thioanisole. Such a reaction can be effected in the presence of an additional inert solvent or diluent (as defined hereinbefore in process (a)) advantageously at a temperature in the range 0 to 80°C, conveniently at about 40°C.

(e) Production of those compounds of formula I and salts thereof wherein at least one R^3 is $R^{10}X^2$ wherein R^{10} is as defined hereinbefore and X^2 is -O-, -S-, -SO₂-, -CONR¹²-, -

SO₂NR¹³- or -NR¹⁵- (wherein R¹², R¹³ and R¹⁵ each independently represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) can be achieved by the reaction, conveniently in the presence of a base (as defined hereinbefore in process (a)) of a compound of the formula VIII:



(VIII)


(wherein R¹, R², R³, n and X² are as hereinbefore defined and s is an integer from 0 to 3) with a compound of formula IX:



(wherein R¹⁰ and L¹ are as hereinbefore defined), L¹ is a displaceable moiety for example a halogeno or sulphonyloxy group such as a bromo or methanesulphonyloxy group. The reaction is preferably effected in the presence of a base (as defined hereinbefore in process (a)) and advantageously in the presence of an inert solvent or diluent (as defined hereinbefore in process (a)), advantageously at a temperature in the range, for example 10 to 150°C, conveniently at about 50°C.

(f) Compounds of the formula I and salts thereof wherein at least one R³ is R¹⁰X² wherein R¹⁰ is as defined hereinbefore and X² is -O-, -S-, or -NR¹⁵- (wherein R¹⁵ represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) may be prepared by the reaction of a compound of the formula X:


- 37 -



(X)

with a compound of the formula XI:

5



(XI)

(wherein L^1 , R^1 , R^2 , R^3 , R^{10} , n , s and X^2 are all as hereinbefore defined). The reaction may conveniently be effected in the presence of a base (as defined hereinbefore in process (a)) and advantageously in the presence of an inert solvent or diluent (as defined hereinbefore in process (a)), advantageously at a temperature in the range, for example 10 to 150°C, conveniently at about 100°C.


- (g) Compounds of the formula I and salts thereof wherein at least one R^3 is $R^{10}X^2$ wherein X^2 is as defined hereinbefore and R^{10} is $C_{1-3}alkylR^{65}$, [wherein R^{65} is selected from one of the following six groups:
- 1) $X^{11}C_{1-3}alkyl$ (wherein X^{11} represents -O-, -S-, -SO₂-, -NR⁶⁶CO- or -NR⁶⁷SO₂- (wherein R^{66} and R^{67} which may be the same or different are each hydrogen, $C_{1-3}alkyl$ or $C_{1-3}alkoxyC_{2-3}alkyl$);
 - 2) $NR^{68}R^{69}$ (wherein R^{68} and R^{69} which may be the same or different are each hydrogen, $C_{1-3}alkyl$ or $C_{1-3}alkoxyC_{2-3}alkyl$);
 - 3) $X^{12}C_{1-3}alkylX^6R^{27}$ (wherein X^{12} represents -O-, -S-, -SO₂-, -NR⁷⁰CO-, -NR⁷¹SO₂- or NR⁷² (wherein R^{70} , R^{71} , and R^{72} which may be the same or different are each hydrogen, $C_{1-3}alkyl$ or $C_{1-3}alkoxyC_{2-3}alkyl$) and X^6 and R^{27} are as defined hereinbefore);
 - 4) R^{60} (wherein R^{60} is as defined hereinbefore);

5) $X^{13}R^{34}$ (wherein X^{13} represents -O-, -S-, -SO₂-, -NR⁷³CO-, -NR⁷⁴SO₂-, or -NR⁷⁵- (wherein R⁷³, R⁷⁴, and R⁷⁵ which may be the same or different are each hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R³⁴ is as defined hereinbefore); and

6) $X^{14}C_{1-3}alkylR^{34}$ (wherein X^{14} represents -O-, -S-, -SO₂-, -NR⁷⁶CO-, -NR⁷⁷SO₂- or -NR⁷⁸-

(wherein R⁷⁶, R⁷⁷ and R⁷⁸ each independently represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R³⁴ is as defined hereinbefore);]

may be prepared by reacting a compound of the formula XII:



10

(XII)

(wherein L¹, X², R¹, R², R³, n and s are as hereinbefore defined) with a compound of the formula XIII:

15

R⁶⁵-H

(XIII)

(wherein R⁶⁵ is as defined hereinbefore) to give a compound of the formula I or salt thereof.

The reaction may conveniently be effected in the presence of a base (as defined hereinbefore in process (a)) and advantageously in the presence of an inert solvent or diluent (as defined hereinbefore in process (a)), and at a temperature in the range, for example 0 to 150°C, conveniently at about 50°C.

20

Processes (a)-(d) and (g) are preferred over processes (e) and (f).

(h) The production of those compounds of the formula I and salts thereof wherein one or more of the substituents (R³)_m is represented by -NR⁷⁹R⁸⁰-, where one or both of R⁷⁹


25

and R⁸⁰ are C₁₋₃alkyl, may be effected by the reaction of compounds of formula I wherein the

substituent $(R^3)_m$ is an amino group and an alkylating agent, preferably in the presence of a base as defined hereinbefore. Such alkylating agents are C_{1-3} alkyl moieties bearing a displaceable moiety as defined hereinbefore such as C_{1-3} alkyl halides for example C_{1-3} alkyl chloride, bromide or iodide. The reaction is preferably effected in the presence of an inert solvent or diluent (as defined hereinbefore in process (a)) and at a temperature in the range, for example, 10 to 100°C, conveniently at about ambient temperature. The production of compounds of formula I and salts thereof wherein one or more of the substituents R^2 or R^3 is an amino group may be effected by the reduction of a corresponding compound of formula I wherein the substituent(s) at the corresponding position(s) of the quinazoline and/or benz ring of the oxindole group is/are a nitro group(s). The reduction may conveniently be effected as described in process (i) hereinafter. The production of a compound of formula I and salts thereof wherein the substituent(s) at the corresponding position(s) of the quinazoline and/or benz ring of the oxindole group is/are a nitro group(s) may be effected by the processes described hereinbefore and hereinafter in processes (a-g) and (i-viii) using a compound selected from the compounds of the formulae (I-XXXIII) in which the substituent(s) at the corresponding position(s) of the quinazoline and/or benz ring of the oxindole group is/are a nitro group(s).

Synthesis of Intermediates

(i) The compounds of formula III and salts thereof, constitute a further feature of the present invention. Such compounds in which L^1 is halogeno may for example be prepared by halogenating a compound of the formula XIV:



(XIV)


(wherein R^3 and m are as hereinbefore defined).

Convenient halogenating agents include inorganic acid halides, for example thionyl chloride, phosphorus(III)chloride, phosphorus(V)oxychloride and phosphorus(V)chloride. The halogenation reaction is conveniently effected in the presence of an inert solvent or

diluent such as for example a halogenated solvent such as methylene chloride, trichloromethane or carbon tetrachloride, or an aromatic hydrocarbon solvent such as benzene or toluene. The reaction is conveniently effected at a temperature in the range, for example 10 to 150°C, preferably in the range 40 to 100°C.

- 5 The compounds of formula XIV and salts thereof which constitute a further feature of the present invention may for example be prepared by reacting a compound of the formula XV:

10




(XV)

- 15 (wherein R^3 , s and L^1 are as hereinbefore defined) with a compound of the formula XI as hereinbefore defined. The reaction may conveniently be effected in the presence of a base (as defined hereinbefore in process (a)) and advantageously in the presence of an inert solvent or diluent (as defined hereinbefore in process (a)), advantageously at a temperature in the range, for example 10 to 150°C, conveniently at about 100°C.

- 20 The compounds of formula XIV and salts thereof may also be prepared by cyclising a compound of the formula XVI:

25




(XVI)

- 30 (wherein R^3 and m , are as hereinbefore defined, and A^1 is an hydroxy, alkoxy (preferably C_1 , alkoxy) or amino group) whereby to form a compound of formula XIV or salt thereof. The

cyclisation may be effected by reacting a compound of the formula XVI, where A¹ is an hydroxy or alkoxy group, with formamide or an equivalent thereof effective to cause cyclisation whereby a compound of formula XIV or salt thereof is obtained, such as [3-(dimethylamino)-2-azaprop-2-enylidene]dimethylammonium chloride. The cyclisation is
 5 conveniently effected in the presence of formamide as solvent or in the presence of an inert solvent or diluent such as an ether for example 1,4-dioxan. The cyclisation is conveniently effected at an elevated temperature, preferably in the range 80 to 200°C. The compounds of formula XIV may also be prepared by cyclising a compound of the formula XVI, where A¹ is an amino group, with formic acid or an equivalent thereof effective to cause cyclisation
 10 whereby a compound of formula XIV or salt thereof is obtained. Equivalents of formic acid effective to cause cyclisation include for example a tri-C₁₋₄alkoxymethane, for example triethoxymethane and trimethoxymethane. The cyclisation is conveniently effected in the presence of a catalytic amount of an anhydrous acid, such as a sulphonic acid for example p-toluenesulphonic acid, and in the presence of an inert solvent or diluent such as for example a
 15 halogenated solvent such as methylene chloride, trichloromethane or carbon tetrachloride, an ether such as diethylether or tetrahydrofuran, or an aromatic hydrocarbon solvent such as toluene. The cyclisation is conveniently effected at a temperature in the range, for example 10 to 100°C, preferably in the range 20 to 50°C.

Compounds of formula XVI and salts thereof, which constitute a further feature of
 20 the present invention, may for example be prepared by the reduction of the nitro group in a compound of the formula XVII:




(XVII)

25

(wherein R³, m and A¹ are as hereinbefore defined) to yield a compound of formula XVI as hereinbefore defined. The reduction of the nitro group may conveniently be effected by any of the procedures known for such a transformation. The reduction may be carried out, for

example, by the hydrogenation of a solution of the nitro compound in the presence of an inert solvent or diluent as defined hereinbefore in the presence of a metal effective to catalyse hydrogenation reactions such as palladium or platinum. A further reducing agent is, for example, an activated metal such as activated iron (produced for example by washing iron powder with a dilute solution of an acid such as hydrochloric acid). Thus, for example, the reduction may be effected by heating the nitro compound and the activated metal in the presence of a solvent or diluent such as a mixture of water and alcohol, for example methanol or ethanol, to a temperature in the range, for example 50 to 150°C, conveniently at about 70°C.

Compounds of the formula XVII and salts thereof which constitute a further feature of the present invention, may for example be prepared by the reaction of a compound of the formula XVIII:




(XVIII)

(wherein R^3 , s , L^1 and A^1 are as hereinbefore defined) with a compound of the formula XI as hereinbefore defined to give a compound of the formula XVII. The reaction of the compounds of formulae XVIII and XI is conveniently effected under conditions as described for process (f) hereinbefore.

Compounds of formula XVII and salts thereof wherein at least one R^3 is $R^{10}X^2$ and wherein X^2 is $-O-$, $-S-$, $-SO_2-$, $-CONR^{12}-$, $-SO_2NR^{13}-$ or $-NR^{15}-$ (wherein R^{12} , R^{13} and R^{15} each independently represents hydrogen, C_{1-3} alkyl or C_{1-3} alkoxy C_{2-3} alkyl), may for example also be prepared by the reaction of a compound of the formula XIX:


- 43 -



(XIX)

(wherein R^3 , s , A^1 and X^2 are as hereinbefore defined) with a compound of the formula IX as
 5 hereinbefore defined to yield a compound of formula XVII as hereinbefore defined. The
 reaction of the compounds of formulae XIX and IX is conveniently effected under conditions
 as described for process (e) hereinbefore.

The compounds of formula III and salts thereof wherein at least one R^3 is $R^{10}X^2$ and
 wherein X^2 is -O-, -S-, -SO₂-, -CONR¹²-, -SO₂NR¹³- or -NR¹⁵- (wherein R^{12} , R^{13} and R^{15} each
 10 independently represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl), may also be prepared for
 example by reacting a compound of the formula XX:




(XX)

(wherein R^3 , X^2 and s are as hereinbefore defined, and L^2 represents a displaceable protecting
 20 moiety) with a compound of the formula IX as hereinbefore defined, whereby to obtain a
 compound of formula III in which L^1 is represented by L^2 .

A compound of formula XX is conveniently used in which L^2 represents a phenoxy
 group which may if desired carry up to 5 substituents, preferably up to 2 substituents, selected
 25 from halogeno, nitro and cyano. The reaction may be conveniently effected under conditions
 as described for process (e) hereinbefore.

The compounds of formula XX and salts thereof which constitute a further feature of the present invention may for example be prepared by deprotecting a compound of the formula XXI:




(XXI)

(wherein R^3 , P^2 , X^2 , s and L^2 are as hereinbefore defined). Deprotection may be effected by techniques well known in the literature, for example where P^2 represents a benzyl group deprotection may be effected by hydrogenolysis or by treatment with trifluoroacetic acid.

One compound of formula III may if desired be converted into another compound of formula III in which the moiety L^1 is different. Thus for example a compound of formula III in which L^1 is other than halogeno, for example optionally substituted phenoxy, may be converted to a compound of formula III in which L^1 is halogeno by hydrolysis of a compound of formula III (in which L^1 is other than halogeno) to yield a compound of formula XIV as hereinbefore defined, followed by introduction of halide to the compound of formula XIV, thus obtained as hereinbefore defined, to yield a compound of formula III in which L^1 represents halogen.

(ii) Compounds of formula IV and salts thereof may be prepared by any of the known procedures, references include: "The Chemistry of Indoles" R. J. Sundberg page 341, 1970 Academic, New York, Gassman and Bergen, 1974, Jnl. Am. Chem. Soc., 96, 5508 and 5512, Quallich and Morrissey, 1993, Synthesis, 51, Cherest et al., 1989, Tetrahedron Letters, 30, 715, Marfat et al., 1987, Tetrahedron Letters, 28, 4027-4030.

(iii) The compounds of formula V and salts thereof, constitute a further feature of the present invention, and may for example be prepared by the reaction of a compound of formula III as hereinbefore defined with a compound of the formula XXII:




(XXII)

- 5 (wherein R^2 , n and P^I are as hereinbefore defined). The reaction may for example be effected as described for process (a) hereinbefore.

The compounds of formula V and salts thereof may also be prepared by reacting a compound of formula XXIII:

10




(XXIII)

- 15 (wherein R^2 , R^3 , L^I , n , s and P^I are as hereinbefore defined) with a compound of formula XI as hereinbefore defined. The reaction may for example be effected as described for process (f) above.

- The compounds of formula V and salts thereof wherein at least one R^3 is $R^{10}X^2$ and wherein X^2 is -O-, -S-, -SO₂-, -CONR¹²-, -SO₂NR¹³- or -NR¹⁵- (wherein R^{12} , R^{13} and R^{15} each
 20 independently represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl), may also be prepared by reacting a compound of formula XXIV:


- 46 -



(XXIV)

5 (wherein R^2 , R^3 , X^2 , n , s and P^1 are as hereinbefore defined) with a compound of the formula IX as hereinbefore defined. The reaction may for example be effected as described for process (e) hereinbefore.

The compounds of formula XXIII and salts thereof constitute a further feature of the present invention and may for example be prepared by reaction of a compound of formula
 10 XXV:




(XXV)

15 (wherein R^3 , s and each L^1 are as hereinbefore defined, and the L^1 in the 4-position and the other L^1 in a further position on the quinazoline ring may be the same or different) with a compound of the formula XXII as hereinbefore defined. The reaction may be effected for example by a process as described in (a) above.

20 Compounds of the formula XXIV and salts thereof which constitute a further feature of the present invention may be made by reacting compounds of the formulae XXI and XXII as hereinbefore defined, under conditions described in (a) hereinbefore, to give a compound of formula XXVI:


- 47 -



(XXVI)

- 5 (wherein R^2 , R^3 , P^1 , P^2 , X^2 , s and n are as hereinbefore defined with the proviso that X^2 is not $-CH_2-$) and then deprotecting the compound of formula XXVI for example as described in (i) above.


- (iv) Compounds of the formula VI and salts thereof constitute a further feature of the present invention and may be made by the reaction of a compound of formula III as
 10 hereinbefore defined with a compound of the formula XXVII:



(XXVII)

(wherein R^2 , n , and Y are as hereinbefore defined). The reaction may for example be effected as described for the process (a) hereinbefore.

- 15 (v) The compounds of formula VII and salts thereof, constitute a further feature of the present invention, and may for example be prepared by the reaction of a compound of formula III as hereinbefore defined with a compound of the formula XXVIII:




(XXVIII)

(wherein R^1 , R^2 , n , $p1$ and P^2 are as hereinbefore defined). The reaction may for example be effected as described for process (a) hereinbefore.

The compounds of formula VII and salts thereof may also be prepared by reacting a compound of formula XXIX:

5




(XXIX)

10

(wherein R^1 , R^2 , R^3 , L^1 , n , $p1$, s and P^2 are as hereinbefore defined) with a compound of formula XI as hereinbefore defined. The reaction may for example be effected as described for process (f) above.

The compounds of formula VII and salts thereof wherein at least one R^3 is $R^{10}X^2$ and wherein X^2 is $-O-$, $-S-$, $-SO_2-$, $-CONR^{12}-$, $-SO_2NR^{13}-$ or $-NR^{15}-$ (wherein R^{12} , R^{13} and R^{15} each independently represents hydrogen, C_{1-3} alkyl or C_{1-3} alkoxy C_{2-3} alkyl), may also be prepared by reacting a compound of formula XXX:

15




20

(XXX)

(wherein R^1 , R^2 , R^3 , X^2 , n , s , $p1$ and P^2 are as hereinbefore defined) with a compound of the formula IX as hereinbefore defined. The reaction may for example be effected as described for process (e) hereinbefore.

- 5 The compounds of formula XXIX and salts thereof may for example be prepared by the reaction of a compound of formula XXV as hereinbefore defined with a compound of the formula XXVIII as hereinbefore defined. The reaction may be effected for example by a process as described in (a) above.

- Compounds of the formula XXX and salts thereof may be made by reacting compounds of the formulae XXI and XXVIII as hereinbefore defined, under conditions
10 described in (a) hereinbefore, to give a compound of formula XXXI:




(XXXI)

- 15 (wherein R^1 , R^2 , R^3 , P^2 , X^2 , $p1$, n and s are as hereinbefore defined with the proviso that X^2 is not $-CH_2-$) and then deprotecting the compound of formula XXXI for example as described in (i) above.

- (vi) Compounds of formula VIII as hereinbefore defined and salts thereof constitute a further feature of the present invention and may be made by deprotecting the compound of
20 formula XXXII:

- 50 -




(XXXII)

- 5 (wherein R¹, R², R³, P², X², s and n are as hereinbefore defined with the proviso that X² is not -CH₂-) by a process for example as described in (i) above.

Compounds of the formula XXXII and salts thereof which constitute a further feature of the present invention may be made by reacting compounds of the formulae XXI and IV as hereinbefore defined, under the conditions described in (a) hereinbefore, to give a
 10 compound of the formula XXXII or salt thereof.

(vii) Compounds of the formula X and salts thereof constitute a further feature of the present invention and may be made by reacting compounds of the formulae XXV and IV as hereinbefore defined, the reaction for example being effected by a process as described in (a) above.

- 15 (viii) Compounds of formula XII as defined hereinbefore and salts thereof constitute a further feature of the present invention and may for example be made by the reaction of compounds of formula VIII as defined hereinbefore with compounds of the formula XXXIII:



(XXXIII)

20

(wherein L¹ is as hereinbefore defined) to give compounds of formula XII or salts thereof. The reaction may be effected for example by a process as described in (e) above.

- When a pharmaceutically acceptable salt of a compound of the formula I is required,
 25 it may be obtained, for example, by reaction of said compound with, for example, an acid using a conventional procedure, the acid having a pharmaceutically acceptable anion.

Many of the intermediates defined herein are novel, for example, those of the formulae III, V, VI, VII, VIII, X, XII, XIV, XV, XVI, XVII, XVIII, XIX, XX, XXI, XXIII, XXIV, XXV, XXVI, XXIX, XXX, XXXI and XXXII and these are provided as a further feature of the invention.

5 The identification of compounds which potently inhibit the tyrosine kinase activity associated with the VEGF receptors such as Flt and/or KDR and which inhibit angiogenesis and/or increased vascular permeability is desirable and is the subject of the present invention. These properties may be assessed, for example, using one or more of the procedures set out below:

10 (a) In Vitro Receptor Tyrosine Kinase Inhibition Test

 This assay determines the ability of a test compound to inhibit tyrosine kinase activity. DNA encoding VEGF, FGF or EGF receptor cytoplasmic domains may be obtained by total gene synthesis (Edwards M, International Biotechnology Lab 5(3), 19-25, 1987) or by cloning. These may then be expressed in a suitable expression system to obtain polypeptide
15 with tyrosine kinase activity. For example VEGF, FGF and EGF receptor cytoplasmic domains, which were obtained by expression of recombinant protein in insect cells, were found to display intrinsic tyrosine kinase activity. In the case of the VEGF receptor Flt (Genbank accession number X51602), a 1.7kb DNA fragment encoding most of the cytoplasmic domain, commencing with methionine 783 and including the termination codon,
20 described by Shibuya et al (Oncogene, 1990, 5: 519-524), was isolated from cDNA and cloned into a baculovirus transplacement vector (for example pAcYM1 (see The Baculovirus Expression System: A Laboratory Guide, L.A. King and R. D. Possee, Chapman and Hall, 1992) or pAc360 or pBlueBacHis (available from Invitrogen Corporation)). This recombinant construct was co-transfected into insect cells (for example *Spodoptera frugiperda* 21(Sf21))
25 with viral DNA (eg Pharmingen BaculoGold) to prepare recombinant baculovirus. (Details of the methods for the assembly of recombinant DNA molecules and the preparation and use of recombinant baculovirus can be found in standard texts for example Sambrook et al, 1989, Molecular cloning - A Laboratory Manual, 2nd edition, Cold Spring Harbour Laboratory Press and O'Reilly et al, 1992, Baculovirus Expression Vectors - A Laboratory Manual, W. H. Freeman and Co, New York). For other tyrosine kinases for use in assays, cytoplasmic
30 fragments starting from methionine 806 (KDR, Genbank accession number L04947),

methionine 668 (EGF receptor, Genbank accession number X00588) and methionine 399 (FGF R1 receptor, Genbank accession number X51803) may be cloned and expressed in a similar manner.

For expression of cFIt tyrosine kinase activity, Sf21 cells were infected with
5 plaque-pure cFIt recombinant virus at a multiplicity of infection of 3 and harvested 48 hours later. Harvested cells were washed with ice cold phosphate buffered saline solution (PBS) (10mM sodium phosphate pH7.4, 138mM sodium chloride, 2.7mM potassium chloride) then resuspended in ice cold HNTG/PMSF (20mM Hepes pH7.5, 150mM sodium chloride, 10% v/v glycerol, 1% v/v Triton X100, 1.5mM magnesium chloride, 1mM ethylene glycol-
10 bis(β-aminoethyl ether) N,N,N',N'-tetraacetic acid (EGTA), 1mM PMSF (phenylmethylsulphonyl fluoride); the PMSF is added just before use from a freshly-prepared 100mM solution in methanol) using 1ml HNTG/PMSF per 10 million cells. The suspension was centrifuged for 10 minutes at 13,000 rpm at 4°C, the supernatant (enzyme stock) was removed and stored in aliquots at -70°C. Each new batch of stock enzyme was titrated in the
15 assay by dilution with enzyme diluent (100mM Hepes pH 7.4, 0.2mM sodium orthovanadate, 0.1% v/v Triton X100, 0.2mM dithiothreitol). For a typical batch, stock enzyme is diluted 1 in 2000 with enzyme diluent and 50μl of dilute enzyme is used for each assay well.

A stock of substrate solution was prepared from a random copolymer containing tyrosine, for example Poly (Glu, Ala, Tyr) 6:3:1 (Sigma P3899), stored as 1 mg/ml stock in
20 PBS at -20°C and diluted 1 in 500 with PBS for plate coating.

On the day before the assay 100μl of diluted substrate solution was dispensed into all wells of assay plates (Nunc maxisorp 96-well immunoplates) which were sealed and left overnight at 4°C.

On the day of the assay the substrate solution was discarded and the assay plate
25 wells were washed once with PBST (PBS containing 0.05% v/v Tween 20) and once with 50mM Hepes pH7.4.

Test compounds were diluted with 10% dimethylsulphoxide (DMSO) and 25μl of diluted compound was transferred to wells in the washed assay plates. "Total" control wells contained 10% DMSO instead of compound. Twenty five microlitres of 40mM
30 manganese(II)chloride containing 8μM adenosine-5'-triphosphate (ATP) was added to all test wells except "blank" control wells which contained manganese(II)chloride without ATP. To

start the reactions 50µl of freshly diluted enzyme was added to each well and the plates were incubated at room temperature for 20 minutes. The liquid was then discarded and the wells were washed twice with PBST. One hundred microlitres of mouse IgG anti-phosphotyrosine antibody (Upstate Biotechnology Inc. product 05-321), diluted 1 in 6000 with PBST

5 containing 0.5% w/v bovine serum albumin (BSA), was added to each well and the plates were incubated for 1 hour at room temperature before discarding the liquid and washing the wells twice with PBST. One hundred microlitres of horse radish peroxidase (HRP)-linked sheep anti-mouse Ig antibody (Amersham product NXA 931), diluted 1 in 500 with PBST

10 containing 0.5% w/v BSA, was added and the plates were incubated for 1 hour at room temperature before discarding the liquid and washing the wells twice with PBST. One hundred microlitres of 2,2'-azino-bis(3-ethylbenzthiazoline-6-sulphonic acid) (ABTS) solution, freshly prepared using one 50mg ABTS tablet (Boehringer 1204 521) in 50ml freshly prepared 50mM phosphate-citrate buffer pH5.0 + 0.03% sodium perborate (made with

15 1 phosphate citrate buffer with sodium perborate (PCSB) capsule (Sigma P4922) per 100ml distilled water), was added to each well. Plates were then incubated for 20-60 minutes at room temperature until the optical density value of the "total" control wells, measured at 405nm using a plate reading spectrophotometer, was approximately 1.0. "Blank" (no ATP) and "total" (no compound) control values were used to determine the dilution range of test compound which gave 50% inhibition of enzyme activity.

20

(b) In Vitro HUVEC Proliferation Assay

This assay determines the ability of a test compound to inhibit the growth factor-stimulated proliferation of human umbilical vein endothelial cells (HUVEC).

HUVEC cells were isolated in MCDB 131 (Gibco BRL) + 7.5% v/v foetal calf

25 serum (FCS) and were plated out (at passage 2 to 8), in MCDB 131 + 2% v/v FCS + 3µg/ml heparin + 1µg/ml hydrocortisone, at a concentration of 1000 cells/well in 96 well plates. After a minimum of 4 hours they were dosed with the appropriate growth factor (i.e. VEGF 3ng/ml, EGF 3ng/ml or b-FGF 0.3ng/ml) and compound. The cultures were then incubated for 4 days at 37°C with 7.5% CO₂. On day 4 the cultures were pulsed with 1µCi/well of

30 tritiated-thymidine (Amersham product TRA 61) and incubated for 4 hours. The cells were harvested using a 96-well plate harvester (Tomtek) and then assayed for incorporation of

tritium with a Beta plate counter. Incorporation of radioactivity into cells, expressed as cpm, was used to measure inhibition of growth factor-stimulated cell proliferation by compounds.

(c) In Vivo Rat Uterine Oedema Assay

5 This test measures the capacity of compounds to reduce the acute increase in uterine weight in rats which occurs in the first 4-6 hours following oestrogen stimulation. This early increase in uterine weight has long been known to be due to oedema caused by increased permeability of the uterine vasculature and recently Cullinan-Bove and Koos (Endocrinology, 1993,133:829-837) demonstrated a close temporal relationship with increased expression of
10 VEGF mRNA in the uterus. We have found that prior treatment of the rats with a neutralising monoclonal antibody to VEGF significantly reduces the acute increase in uterine weight, confirming that the increase in weight is substantially mediated by VEGF.

 Groups of 20 to 22-day old rats were treated with a single subcutaneous dose of oestradiol benzoate (2.5µg/rat) in a solvent, or solvent only. The latter served as unstimulated
15 controls. Test compounds were orally administered at various times prior to the administration of oestradiol benzoate. Five hours after the administration of oestradiol benzoate the rats were humanely sacrificed and their uteri were dissected, blotted and weighed. The increase in uterine weight in groups treated with test compound and oestradiol benzoate and with oestradiol benzoate alone was compared using a Student T test. Inhibition
20 of the effect of oestradiol benzoate was considered significant when $p < 0.05$.

 According to a further aspect of the invention there is provided a pharmaceutical composition which comprises a compound of the formula I as defined hereinbefore or a pharmaceutically acceptable salt thereof, in association with a pharmaceutically acceptable excipient or carrier.

25 The composition may be in a form suitable for oral administration, for example as a tablet or capsule, for parenteral injection (including intravenous, subcutaneous, intramuscular, intravascular or infusion) for example as a sterile solution, suspension or emulsion, for topical administration for example as an ointment or cream or for rectal administration for example as a suppository. In general the above compositions may be prepared in a conventional manner
30 using conventional excipients.

The compositions of the present invention are advantageously presented in unit dosage form. The compound will normally be administered to a warm-blooded animal at a unit dose within the range 5-5000mg per square metre body area of the animal, i.e. approximately 0.1-100mg/kg. A unit dose in the range, for example, 1-100mg/kg, preferably 1-50mg/kg is envisaged and this normally provides a therapeutically-effective dose. A unit dose form such as a tablet or capsule will usually contain, for example 1-250mg of active ingredient.

According to a further aspect of the present invention there is provided a compound of the formula I or a pharmaceutically acceptable salt thereof as defined hereinbefore for use in a method of treatment of the human or animal body by therapy.

We have found that compounds of the present invention inhibit VEGF receptor tyrosine kinase activity and FGF R1 receptor tyrosine kinase activity and are therefore of interest for their antiangiogenic effects and/or their ability to cause a reduction in vascular permeability.

A further feature of the present invention is a compound of formula I, or a pharmaceutically acceptable salt thereof, for use as a medicament, conveniently a compound of formula I, or a pharmaceutically acceptable salt thereof, for use as a medicament for producing an antiangiogenic and/or vascular permeability reducing effect in a warm-blooded animal such as a human being.

Thus according to a further aspect of the invention there is provided the use of a compound of the formula I, or a pharmaceutically acceptable salt thereof in the manufacture of a medicament for use in the production of an antiangiogenic and/or vascular permeability reducing effect in a warm-blooded animal such as a human being.

According to a further feature of the invention there is provided a method for producing an antiangiogenic and/or vascular permeability reducing effect in a warm-blooded animal, such as a human being, in need of such treatment which comprises administering to said animal an effective amount of a compound of formula I or a pharmaceutically acceptable salt thereof as defined hereinbefore.

As stated above the size of the dose required for the therapeutic or prophylactic treatment of a particular disease state will necessarily be varied depending on the host treated, the route of administration and the severity of the illness being treated. Preferably a daily

dose in the range of 1-50mg/kg is employed. However the daily dose will necessarily be varied depending upon the host treated, the particular route of administration, and the severity of the illness being treated. Accordingly the optimum dosage may be determined by the practitioner who is treating any particular patient.

- 5 The antiangiogenic and/or vascular permeability reducing treatment defined hereinbefore may be applied as a sole therapy or may involve, in addition to a compound of the invention, one or more other substances and/or treatments. Such conjoint treatment may be achieved by way of the simultaneous, sequential or separate administration of the individual components of the treatment. In the field of medical oncology it is normal practice
- 10 to use a combination of different forms of treatment to treat each patient with cancer. In medical oncology the other component(s) of such conjoint treatment in addition to the antiangiogenic and/or vascular permeability reducing treatment defined hereinbefore may be: surgery, radiotherapy or chemotherapy. Such chemotherapy may cover three main categories of therapeutic agent:
- 15 (i) other antiangiogenic agents that work by different mechanisms from those defined hereinbefore (for example linomide, inhibitors of integrin $\alpha v \beta 3$ function, angiostatin, razoxin, thalidomide);
- (ii) cytostatic agents such as antioestrogens (for example tamoxifen, toremifene, raloxifene, droloxifene, idoxifyfene), progestogens (for example megestrol acetate), aromatase inhibitors
- 20 (for example anastrozole, letrozole, vorazole, exemestane), antiprogestogens, antiandrogens (for example flutamide, nilutamide, bicalutamide, cyproterone acetate), LHRH agonists and antagonists (for example goserelin acetate, luprolide), inhibitors of testosterone 5α -dihydroreductase (for example finasteride), anti-invasion agents (for example metalloproteinase inhibitors like marimastat and inhibitors of urokinase plasminogen activator
- 25 receptor function) and inhibitors of growth factor function, (such growth factors include for example EGF, platelet derived growth factor and hepatocyte growth factor such inhibitors include growth factor antibodies, growth factor receptor antibodies, tyrosine kinase inhibitors and serine/threonine kinase inhibitors); and
- (iii) antiproliferative/antineoplastic drugs and combinations thereof, as used in medical
- 30 oncology, such as antimetabolites (for example antifolates like methotrexate, fluoropyrimidines like 5-fluorouracil, purine and adenosine analogues, cytosine arabinoside);

antitumour antibiotics (for example anthracyclines like doxorubicin, daunomycin, epirubicin and idarubicin, mitomycin-C, dactinomycin, mithramycin); platinum derivatives (for example cisplatin, carboplatin); alkylating agents (for example nitrogen mustard, melphalan, chlorambucil, busulphan, cyclophosphamide, ifosfamide, nitrosoureas, thiotepa); antimitotic
5 agents (for example vinca alkaloids like vincristine and taxoids like taxol, taxotere);
topoisomerase inhibitors (for example epipodophyllotoxins like etoposide and teniposide, amsacrine, topotecan).

As stated above the compounds defined in the present invention are of interest for their antiangiogenic and/or vascular permeability reducing effects. Such compounds of the
10 invention are expected to be useful in a wide range of disease states including cancer, diabetes, psoriasis, rheumatoid arthritis, Kaposi's sarcoma, haemangioma, acute and chronic nephropathies, atheroma, arterial restenosis, autoimmune diseases, acute inflammation and ocular diseases with retinal vessel proliferation. In particular such compounds of the
15 invention are expected to slow advantageously the growth of primary and recurrent solid tumours of, for example, the colon, breast, prostate, lungs and skin. More particularly such compounds of the invention are expected to inhibit the growth of those primary and recurrent solid tumours which are associated with VEGF and/or FGF, especially those tumours which are significantly dependent on VEGF and/or FGF for their growth and spread, including for example, certain tumours of the colon, breast, prostate, lung, vulva and skin.

20 In addition to their use in therapeutic medicine, the compounds of formula I and their pharmaceutically acceptable salts are also useful as pharmacological tools in the development and standardisation of in vitro and in vivo test systems for the evaluation of the effects of inhibitors of VEGF receptor tyrosine kinase activity and FGF R1 receptor tyrosine kinase activity in laboratory animals such as cats, dogs, rabbits, monkeys, rats and mice, as
25 part of the search for new therapeutic agents.

It is to be understood that where the term "ether" is used anywhere in this specification it refers to diethyl ether.

The invention will now be illustrated in the following non-limiting Examples in which, unless otherwise stated:-

- 58 -

[(i) evaporations were carried out by rotary evaporation in vacuo and work-up procedures were carried out after removal of residual solids such as drying agents by filtration;

(ii) operations were carried out at ambient temperature, that is in the range 18-25°C and under an atmosphere of an inert gas such as argon;

(iii) column chromatography (by the flash procedure) and medium pressure liquid chromatography (MPLC) were performed on Merck Kieselgel silica (Art. 9385) or Merck Lichroprep RP-18 (Art. 9303) reversed-phase silica obtained from E. Merck, Darmstadt, Germany;

(iv) yields are given for illustration only and are not necessarily the maximum attainable;

(v) melting points are uncorrected and were determined using a Mettler SP62 automatic melting point apparatus, an oil-bath apparatus or a Koffler hot plate apparatus.

(vi) the structures of the end-products of the formula I were confirmed by nuclear (generally proton) magnetic resonance (NMR) and mass spectral techniques; proton magnetic resonance chemical shift values were measured on the delta scale and peak multiplicities are shown as follows: s, singlet; d, doublet; t, triplet; m, multiplet; br, broad; q, quartet, quin, quintet;

(vii) intermediates were not generally fully characterised and purity was assessed by thin layer chromatography (TLC), high-performance liquid chromatography (HPLC), infra-red (IR) or NMR analysis;

(viii) petroleum ether refers to that fraction boiling between 40-60°C

(ix) the following abbreviations have been used:-

DMF N,N-dimethylformamide
DMSO dimethylsulphoxide
TFA trifluoroacetic acid
THF tetrahydrofuran.]

Example 1

Potassium bis(trimethylsilyl)amide (5ml of a 0.5M solution in toluene, 2.5mmol) was rapidly added at ambient temperature to a solution of 4-chloro-6,7-dimethoxyquinazoline (337mg, 1.5mmol) and 1-methyloxindole (368mg, 2.5mmol), (prepared according to the method described in J. Am. Chem. Soc 1945, 67, 1656), in THF (30ml). After 30 minutes, the precipitate was collected by filtration and washed with ether. A second precipitate was isolated from the filtrate. The two solids were mixed and purified by column flash chromatography using methylene chloride/methanol (96/4 and 94/6) as eluent. During the evaporation of the solvents, the compound crystallised. The solid was filtered off, washed with ether and dried under vacuum to give **6,7-dimethoxy-4-(1-methyloxindol-3-yl)quinazoline** (103mg, 20%) as red orange needles.

m.p. 233-235°C

¹H NMR Spectrum: (DMSO-d₆, TFA) 3.37(s, 3H); 3.85(s, 3H); 4.00(s, 3H); 7.11(t, 1H); 7.16(d, 1H); 7.26(t, 1H); 7.27(s, 1H); 7.72(d, 1H); 7.75(s, 1H); 8.90(s, 1H)

MS - ESI: 336 [MH]⁺

Elemental analysis:	Found	C 66.6	H 5.3	N 12.3
---------------------	-------	--------	-------	--------

C ₁₉ H ₁₇ N ₃ O ₃ 0.4H ₂ O	Requires	C 66.6	H 5.2	N 12.3%
---	----------	--------	-------	---------

The starting material was prepared as follows:

A mixture of 4,5-dimethoxyanthranilic acid (19.7g) and formamide (10ml) was stirred and heated at 190°C for 5 hours. The mixture was allowed to cool to approximately 80°C and water (50ml) was added. The mixture was then allowed to stand at ambient temperature for 3 hours. The precipitate was collected by filtration, washed with water and dried to give 6,7-dimethoxy-3,4-dihydroquinazolin-4-one (3.65g).

To a portion (2.06g) of the material so obtained were added thionyl chloride (20ml) and DMF (1 drop) and the mixture stirred and heated at reflux for 2 hours. Excess thionyl chloride was removed by evaporation and the residue was partitioned between ethyl acetate and a saturated aqueous sodium hydrogen carbonate solution. The organic phase was washed with water, dried (MgSO₄) and the solvent removed by evaporation. The residue was purified by column chromatography using increasingly polar mixtures of methylene chloride and ethyl acetate as eluent to give 4-chloro-6,7-dimethoxyquinazoline (0.6g, 27%).

Example 2

- Potassium bis(trimethylsilyl)amide (48ml of a 0.5M solution in toluene, 24mmol) was rapidly added to a solution of 4-chloro-6-methoxy-7-(2-methoxyethoxy)quinazoline (3.2g, 11.9mmol) and 1-diethoxymethyloxindole (4.0g, 14.9mmol), (prepared according to Synthesis 1975, 168), in THF (240ml) at ambient temperature. After 2 hours, the precipitate was filtered off and washed with ether. The filtrate was evaporated. The residue was mixed with the precipitate and partitioned between methylene chloride and water and the organic phase washed with saturated aqueous sodium chloride solution. The organic layer was dried (MgSO₄) and the solvent removed by evaporation. The residue was purified by column flash chromatography using methylene chloride/methanol (95/5) as eluent. The purified product was triturated with a mixture of ether and petroleum ether to give 4-(1-diethoxymethyloxindol-3-yl)-6-methoxy-7-(2-methoxyethoxy)quinazoline (4.03g, 74%) as a red orange solid.
- m.p. 148-150°C
- ¹H NMR Spectrum: (DMSO-d₆, TFA) 1.07(t, 6H); 3.37(s, 3H); 3.45(q, 4H); 3.77-3.79(m, 2H); 3.85(s, 3H); 4.34-4.37(m, 2H); 7.13-7.23(m, 2H); 7.28(s, 1H); 7.87(br d, 1H); 7.95(s, 1H); 8.07(d, 1H); 8.64(s, 1H); 9.44(s, 1H)
- MS - ESI: 490 [MNa]⁺
- | | | | | |
|---|----------|--------|-------|--------|
| Elemental analysis: | Found | C 63.3 | H 6.6 | N 8.9 |
| C ₂₅ H ₂₉ N ₃ O ₆ 0.4H ₂ O | Requires | C 63.3 | H 6.3 | N 8.9% |

The starting material was prepared as follows:

- A mixture of ethyl 4-hydroxy-3-methoxybenzoate (9.8g, 50mmol), 2-bromoethyl methyl ether (8.46ml, 90mmol) and potassium carbonate (12.42g, 90mmol) in acetone (60ml) was heated at reflux for 30 hours. The mixture was allowed to cool and the solids removed by filtration. The volatiles were removed from the filtrate by evaporation and the residue triturated with hexane to give ethyl 3-methoxy-4-(2-methoxyethoxy)benzoate (11.3g, 89%) as a white solid.
- m.p. 57-60°C

¹H NMR Spectrum: (DMSO-d₆) 1.31(t, 3H); 3.29(s, 3H); 3.32(s, 3H); 3.68(m, 2H); 4.16(m, 2H); 4.28(q, 2H); 7.06(d, 1H); 7.45(d, 1H); 7.56(dd, 1H)

MS-FAB: 255 [MH]⁺

5 Ethyl 3-methoxy-4-(2-methoxyethoxy)benzoate (9.5g, 37mmol) was added in portions to stirred concentrated nitric acid (75ml) at 0°C. The mixture was allowed to warm to ambient temperature and stirred for a further 90 minutes. The mixture was diluted with water and extracted with methylene chloride, dried (MgSO₄) and the solvent removed by evaporation. The residue was triturated with hexane to give ethyl 5-methoxy-4-(2-methoxyethoxy)-2-nitrobenzoate (10.6g, 95%) as an orange solid.

10 m.p. 68-69°C

¹H NMR Spectrum: (DMSO-d₆) 1.27(t, 3H); 3.30(s, 3H); 3.69(m, 2H); 3.92(s, 3H); 4.25(m, 2H); 4.29(q, 2H); 7.30(s, 1H); 7.65(s, 1H)

MS-Cl: 300 [MH]⁺

15 A mixture of ethyl 5-methoxy-4-(2-methoxyethoxy)-2-nitrobenzoate (10.24g, 34mmol), cyclohexene (30ml) and 10% palladium-on-charcoal catalyst (2.0g) in methanol (150ml) was heated at reflux for 5 hours. The reaction mixture was allowed to cool and diluted with methylene chloride. The catalyst was removed by filtration and the volatiles removed from the filtrate by evaporation. The residue was recrystallised from ethyl acetate/hexane to give ethyl 2-amino-5-methoxy-4-(2-methoxyethoxy) benzoate (8.0g) as a buff solid. Formamide (80ml) was added to this product and the mixture heated at 170°C for 20 18 hours. About half the solvent was removed by evaporation under high vacuum and the residue was left to stand overnight. The solid product was collected by filtration, washed with ether and dried to give 6-methoxy-7-(2-methoxyethoxy)-3,4-dihydroquinazolin-4-one (5.3g, 62% over two steps) as a grey solid.

25 ¹H NMR Spectrum: (DMSO-d₆) 3.35(s, 3H); 3.74(m, 2H); 3.89(s, 3H); 4.26(m, 2H); 7.15(s, 1H); 7.47(s, 1H); 7.98(s, 1H); 12.03(br s, 1 H)

MS-Cl: 251 [MH]⁺

30 DMF (0.5ml) was added to a mixture of 6-methoxy-7-(2-methoxyethoxy)-3,4-dihydroquinazolin-4-one (5.1g, 20mmol) in thionyl chloride (50ml). The mixture was stirred and heated at reflux for 3 hours, allowed to cool and the excess thionyl chloride removed by evaporation. The residue was suspended in methylene chloride and washed with aqueous

sodium hydrogen carbonate solution. The aqueous phase was extracted with methylene chloride and the combined extracts dried (MgSO_4). The crude product was recrystallised from methylene chloride/hexane to give 4-chloro-6-methoxy-7-(2-methoxyethoxy)quinazoline (2.8g, 51%) as a fine white solid.

- 5 ^1H NMR Spectrum: (DMSO-d_6) 3.37(s, 3H); 3.77(m, 2H); 4.01(s, 3H); 4.37(m, 2H); 7.40(s, 1H); 7.49(s, 1H); 8.88(s, 1H)
MS-Cl: 269 $[\text{MH}]^+$

Example 3

- 10 2M Hydrochloric acid (40ml, 80mmol) was added to a solution of 4-(1-diethoxymethyloxindol-3-yl)-6-methoxy-7-(2-methoxyethoxy)quinazoline (3.7g, 79mmol), (prepared as described in Example 2), in ethanol (300ml) at 80°C. A precipitate rapidly appeared which then went into solution followed after 15 minutes by the formation of a new precipitate. The mixture was stirred for 45 minutes at 80°C, the solid was collected by
15 filtration, washed with ethanol and then ether and dried under vacuum at 65°C to give 6-methoxy-7-(2-methoxyethoxy)-4-(oxindol-3-yl)quinazoline (2.42g, 84%) as a red orange solid.

m.p. 262-268°C

- ^1H NMR Spectrum: (DMSO-d_6 , TFA) 3.38(s, 3H); 3.78(t, 2H); 3.87(s, 3H); 4.33(t, 2H); 7.02-
20 7.07(m, 2H); 7.18(t, 1H); 7.31(s, 1H); 7.70(d, 1H); 7.75(s, 1H); 8.94(s, 1H)
MS - ESI: 366 $[\text{MH}]^+$

Elemental analysis:	Found	C 59.4	H 5.3	N 10.6	Cl 4.8
$\text{C}_{20}\text{H}_{19}\text{N}_3\text{O}_4 \cdot 0.9\text{H}_2\text{O} \cdot 0.55\text{HCl}$	Requires	C 59.8	H 5.4	N 10.5	Cl 4.9%

25 Example 4

- 2M Hydrochloric acid (0.54ml, 1.08mmol) was added to a solution of 4-(1-diethoxymethyloxindol-3-yl)-6-methoxy-7-(2-methoxyethoxy)quinazoline (250mg, 0.53mmol), (prepared as described in Example 2), in ethanol (10ml) at ambient temperature. A precipitate rapidly appeared. The mixture was stirred for 30 minutes, the solid was
30 collected by filtration washed with ethanol and then ether and dried under vacuum at 60°C to

give 4-(1-formyloxindol-3-yl)-6-methoxy-7-(2-methoxyethoxy)quinazoline (141mg, 68%) as an orange solid.

m.p. 228-230°C

¹H NMR Spectrum: (DMSO-d₆, TFA) 3.36(s, 3H); 3.77(t, 2H); 3.84(s, 3H); 4.34(t, 2H); 7.13(t, 2H); 7.20(t, 1H); 7.27(s, 1H); 7.88(d, 1H); 7.96(s, 1H); 8.06(d, 1H); 8.60(s, 1H); 9.43(s, 1H)

MS - ESI: 316 [MNa]⁺

Elemental analysis:	Found	C 62.6	H 5.1	N 10.4
C ₂₁ H ₁₉ N ₃ O ₅ · 0.4H ₂ O	Requires	C 63.0	H 5.0	N 10.5%

10 Example 5

4-Chloro-6-methoxy-7-(3-morpholinopropoxy)quinazoline (338mg, 1mmol) was dissolved in THF (20ml) at 40°C, and 1-diethoxymethyloxindole (705mg, 3mmol), (prepared according to Synthesis 1975, 168), and sodium hydride (138mg, 5.8mmol, pre-washed with petroleum ether) were added successively to the solution at ambient temperature. After stirring for 45 minutes, the volatiles were removed by evaporation and the residue dissolved in methylene chloride. The solution was washed with an aqueous saturated sodium chloride solution, dried (MgSO₄) and the solvent removed by evaporation. The residue was purified by column flash chromatography using methylene chloride/ethyl acetate (1/1) followed by an increasingly polar mixture of methylene chloride and methanol (2 to 4%) as eluent. The purified product was triturated with ether, collected by filtration, washed with a 1/1 mixture of ether/petroleum ether and dried under vacuum to give 4-(1-diethoxymethyloxindol-3-yl)-6-methoxy-7-(3-morpholinopropoxy)quinazoline (425mg, 84%) as a yellow-orange solid.

m.p. 174-176°C

¹H NMR Spectrum: (CDCl₃, CD₃CO₂D) 1.26(t, 6H); 2.33-2.41(m, 2H); 3.24-3.28(m, 6H); 3.55-3.61(m, 2H); 3.75-3.82(m, 2H); 3.88(s, 3H); 3.95-3.97(m, 4H); 4.26-4.29(m, 2H); 6.48(s, 1H); 7.07(t, 1H); 7.14(t, 1H); 7.31(s, 1H); 7.63(d, 1H); 7.81(s, 1H); 8.27(s, 1H)

MS - ESI: 537 [MH]⁺

Elemental analysis:	Found	C 65.4	H 7.1	N 10.5
C ₂₉ H ₃₆ N ₄ O ₆	Requires	C 64.9	H 6.8	N 10.4%

The starting material was prepared as follows:

A mixture of 4-hydroxy-3-methoxybenzoic acid (4.5g, 26.8mmol), 3-morpholinopropyl chloride (9.5g, 58.0mmol), (prepared according to J. Am. Chem. Soc. 1945, 67, 736), potassium carbonate (8.0g, 58mmol), potassium iodide (1.0g, 0.22mmol) and DMF (80ml) was stirred and heated at 100°C for 3 hours. The solid was removed by filtration and the filtrate evaporated. The residue was dissolved in ethanol (50ml), 2M sodium hydroxide (50ml) was added and the mixture heated at 90°C for 2 hours. After partial evaporation, the mixture was acidified with concentrated hydrochloric acid, washed with ether and then subjected to purification on a Diaion (trade mark of Mitsubishi) HP20SS resin column, eluting with water and then with a gradient of methanol (0 to 25%) in hydrochloric acid (pH2). Partial evaporation of the solvents and lyophilisation gave 3-methoxy-4-(3-morpholinopropoxy)benzoic acid (8.65g, 97%).

¹H NMR Spectrum: (DMSO-d₆, TFA) 2.17-2.24(m, 2H); 3.10-3.16(m, 2H); 3.30(t, 2H); 3.52(d, 2H); 3.71(t, 2H); 3.82(s, 3H); 4.01(br d, 2H); 4.14(t, 2H); 7.08(d, 1H); 7.48(d, 1H); 7.59(dd, 1H)

MS - ESI: 296 [MH]⁺

Fuming nitric acid (1.5ml, 36.2mmol) was added slowly at 0°C to a solution of 3-methoxy-4-(3-morpholinopropoxy)benzoic acid (7.78g, 23.5mmol) in TFA (25ml). The cooling bath was removed and the reaction mixture stirred at ambient temperature for 1 hour. The TFA was removed by evaporation and ice was added to the residue. The precipitate was collected by filtration, washed with a minimum of water followed by toluene and ether. The solid was dried under vacuum over phosphorus pentoxide to give 5-methoxy-4-(3-morpholinopropoxy)-2-nitrobenzoic acid (7.54g) which was used without further purification.

¹H NMR Spectrum: (DMSO-d₆, TFA) 2.16-2.23(m, 2H); 3.10-3.17(m, 2H); 3.30(t, 2H); 3.52(d, 2H); 3.66(t, 2H); 3.93(s, 3H); 4.02(br d, 2H); 4.23(t, 2H); 7.34(s, 1H); 7.61(s, 1H)

MS - EI: 340 [M]⁺

Thionyl chloride (15ml) and DMF (0.05ml) were added to 5-methoxy-4-(3-morpholinopropoxy)-2-nitrobenzoic acid (7.54g). The mixture was heated at 50°C for 1 hour, the excess thionyl chloride was removed by evaporation and by azeotropeing with toluene (x2). The resulting solid was suspended in THF (200ml) and ammonia was bubbled through the mixture for 30 minutes. The precipitate was removed by filtration and washed with THF. After concentration of the filtrate by evaporation, the product crystallised and was collected

by filtration to give 5-methoxy-4-(3-morpholinopropoxy)-2-nitrobenzamide (5.25g) as light yellow crystals which were used without further purification.

¹H NMR Spectrum: (DMSO-d₆, TFA) 2.17-2.24(m, 2H); 3.11-3.18(m, 2H); 3.31(t, 2H); 3.53(d, 2H); 3.67(t, 2H); 3.93(s, 3H); 4.03(br d, 2H); 4.21(t, 2H); 7.17(s, 1H); 7.62(s, 1H)

5 MS - EI: 339 [M]⁺

Concentrated hydrochloric acid (30ml) was added to a suspension of 5-methoxy-4-(3-morpholinopropoxy)-2-nitrobenzamide (5.67g) in methanol (150ml) and the mixture was heated to 60°C. When the 5-methoxy-4-(3-morpholinopropoxy)-2-nitrobenzamide had dissolved, iron powder (5.6g, 100mmol) was added in portions to the reaction mixture which was then heated for 90 minutes. After cooling, the insolubles were removed by filtration through diatomaceous earth, the volatiles were removed from the filtrate by evaporation and the residue was purified on a Diaion (trade mark of Mitsubishi) HP20SS resin column, eluting with water and then with hydrochloric acid (pH2). Concentration of the fractions by evaporation gave a precipitate which was collected by filtration and dried under vacuum over phosphorus pentoxide to give 2-amino-5-methoxy-4-(3-morpholinopropoxy)benzamide as a hydrochloride salt (4.67g, 75%) as beige crystals.

¹H NMR Spectrum: (DMSO-d₆, TFA) 2.22-2.28(m, 2H); 3.12(br t, 2H); 3.29(t, 2H); 3.51(d, 2H); 3.75(t, 2H); 3.87(s, 3H); 4.00(br d, 2H); 4.12(t, 2H); 7.06(s, 1H); 7.53(s, 1H)

MS - EI: 309 [M]⁺

20 A mixture of 2-amino-5-methoxy-4-(3-morpholinopropoxy)benzamide (4.57g, 12.25mmol) and Gold's reagent (2.6g, 15.89mmol) in dioxane (35ml) was heated at reflux for 5 hours. Acetic acid (0.55ml) and sodium acetate (1.0g) were added to the reaction mixture which was heated for a further 3 hours. The mixture was cooled to ambient temperature and the volatiles removed by evaporation. The residue was adjusted to pH7 with 2M sodium hydroxide and then purified on a Diaion (trade mark of Mitsubishi) HP20SS resin column, eluting with methanol (gradient of 0 to 60%) in water. Concentration of the fractions by evaporation gave a precipitate which was collected by filtration and dried under vacuum over phosphorus pentoxide to give 4-hydroxy-6-methoxy-7-(3-morpholinopropoxy)quinazoline (3.04g, 78%) as a white solid.

30 ¹H NMR Spectrum: (CDCl₃) 2.10(q, 2H); 2.48(m, 4H); 2.56(t, 2H); 3.72(t, 4H); 4.00(s, 3H); 4.24(t, 2H); 7.18(s, 1H); 7.60(s, 1H); 8.00(s, 1H); 10.86(br s, 1H)

MS - EI: 319 [M]⁺

A mixture of 4-hydroxy-6-methoxy-7-(3-morpholinopropoxy)quinazoline (638mg, 2mmol) and thionyl chloride (8ml) was heated at reflux for 30 minutes. Excess thionyl chloride was removed by evaporation and by azeotrope with toluene (x2). The residue was
 5 suspended in methylene chloride and 10% aqueous solution of sodium hydrogen carbonate was added to the mixture. The organic layer was separated, dried (MgSO₄) and the solvent removed by evaporation. The residue was triturated with ether, the solid was collected by filtration, washed with ether and dried under vacuum to give 4-chloro-6-methoxy-7-(3-morpholinopropoxy)quinazoline (590mg, 87%).

10 ¹H NMR Spectrum: (CDCl₃) 2.10-2.16(m, 2H); 2.48(br s, 4H); 2.57(t, 2H); 3.73(t, 4H); 4.05(s, 3H); 4.29(t, 2H); 7.36(s, 1H); 7.39(s, 1H); 8.86(s, 1H)

MS - ESI: 337 [MH]⁺

Example 6

15 2M Hydrochloric acid (3ml, 6mmol) was added to a solution of 4-(1-diethoxymethyloxindol-3-yl)-6-methoxy-7-(3-morpholinopropoxy)quinazoline (295mg, 0.55mmol), (prepared as described in Example 5), in ethanol (20ml) at 80°C. The mixture was stirred for 45 minutes at 80°C. After cooling of the reaction mixture, partial evaporation of the solvents led to the crystallisation of the compound. The solid was collected by
 20 filtration, washed with ethanol and then ether and dried under vacuum at 50°C to give 6-methoxy-7-(3-morpholinopropoxy)-4-(oxindol-3-yl)quinazoline as a hydrochloride salt (207mg, 84%) as an orange solid.

m.p. 180-183°C

¹H NMR Spectrum: (DMSO-d₆, TFA) 2.28-2.36(m, 2H); 3.13-3.20(m, 2H); 3.34-3.38(m, 2H);
 25 3.56(br d, 2H); 3.74(br t, 2H); 3.88(s, 3H); 4.04(dd, 2H); 4.31-4.34(m, 2H); 7.03-7.08(m, 2H); 7.19(t, 1H); 7.34(s, 1H); 7.69(d, 1H); 7.79(s, 1H); 8.93(s, 1H)

MS - ESI: 435 [MH]⁺

Elemental analysis:	Found	C 55.1	H 6.3	N 10.7	Cl 9.9
C ₂₄ H ₂₆ N ₄ O ₄ · 2H ₂ O · 1.45HCl	Requires	C 55.1	H 6.1	N 10.7	Cl 9.8%

Example 7

Sodium hydride (80mg, 3.4mmol, pre-washed with petroleum ether) was added to a solution of 5-fluorooxindole (423mg, 2.8mmol), (prepared according to Synthesis 1993, 51), in THF (10ml) at ambient temperature under argon. The resulting suspension was stirred for 30 minutes and a partial solution of 4-chloro-6-methoxy-7-(3-morpholinopropoxy)quinazoline (236mg, 0.7mmol), (prepared as described for the starting material in Example 5), in a 1/1 mixture of THF/DMF (14ml), was added and the reaction mixture was stirred at ambient temperature for 15 hours. The volatiles were then removed by evaporation and the residue purified by column flash chromatography using an increasingly polar mixture of methylene chloride and methanol (2 to 10%) as eluent. The purified product was dissolved in a minimum volume of methanol and a 2.9M solution of hydrogen chloride in ether was added to the solution. During the evaporation of the solvents a solid precipitated which was collected by filtration, washed with ether and dried under vacuum to give 4-(5-fluorooxindol-3-yl)-6-methoxy-7-(3-morpholinopropoxy)quinazoline as a hydrochloride salt (298mg, 81%).

m.p. 230-240°C

¹H NMR Spectrum: (DMSO-d₆, TFA) 2.28-2.33(m, 2H); 3.13-3.18(m, 2H); 3.35(t, 2H); 3.55(d, 2H); 3.74(br t, 2H); 3.89(s, 3H); 4.03(d, 2H); 4.33(t, 2H); 6.94-6.99(m, 2H); 7.34(s, 1H); 7.46(d, 1H); 7.81(s, 1H); 8.81(s, 1H)

MS - ESI: 453 [MH]⁺

Elemental analysis:	Found	C 54.4	H 5.5	N 10.3	Cl 11.5
C ₂₄ H ₂₃ N ₄ O ₄ F 0.6H ₂ O 1.75HCl	Requires	C 54.7	H 5.3	N 10.6	Cl 11.8%

Example 8

2M Hydrochloric acid (2.75ml, 5.5mmol) was added to a solution of 4-(5-bromo-1-diethoxymethyloxindol-3-yl)-6-methoxy-7-(2-methoxyethoxy)quinazoline (300mg, 0.55mmol) in ethanol (20ml) at 90°C. The mixture was heated for 1 hour, the solid product was collected by filtration and purified on a Diaion (trade mark of Mitsubishi) HP20SS resin column, eluting with acetonitrile (using a gradient of 0 to 100%) in hydrochloric acid (pH2). Concentration of the fractions by evaporation gave a precipitate which was collected by filtration, washed with water to give a solid which was purified again by column flash chromatography using an increasingly polar mixture of methylene chloride/methanol (95/5 to

80/20) as eluent. After evaporation of the solvent, the resulting solid was dissolved in a minimum volume of methanol and a 2.9M solution of hydrogen chloride in ether was added to the solution. The solid was collected by filtration, washed with ether and dried under vacuum to give 4-(5-bromooxindol-3-yl)-6-methoxy-7-(2-methoxyethoxy)quinazoline as a

5 hydrochloride salt (74mg, 30%) as an orange solid.

¹H NMR Spectrum: (DMSO-d₆, TFA) 3.36(s, 3H); 3.78(t, 2H); 3.93(s, 3H); 4.34(t, 2H); 6.96(d, 1H); 7.26(dd, 1H); 7.31(s, 1H); 7.77(s, 1H); 7.79(d, 1H); 8.80(s, 1H)

MS - ESI: 444 [MH]⁺

Elemental analysis: Found C 49.4 H 4.1 N 8.6 Cl 5.0

10 C₂₀H₁₈N₃O₄Br 0.7H₂O 0.7HCl Requires C 49.8 H 4.2 N 8.7 Cl 5.1%

The starting material was prepared as follows:

A mixture of 5-bromooxindole (6.36g, 30 mmol), (prepared according to J. Am. Chem. Soc. 1975, 67, 1656), and triethyl orthoformate (100ml) was heated and stirred at
15 reflux for 8 hours. The volatiles were removed by evaporation and the residue purified by column flash chromatography using petroleum ether/ether (7/3) as eluent. Evaporation of the solvents lead to a crystalline material which was collected, washed with a minimum of petroleum ether and dried under vacuum to give 5-bromo-1-diethoxymethyloxindole (6.43g, 68%).

20 ¹H NMR Spectrum: (CDCl₃) 1.23(t, 6H); 3.50-3.57(m, 2H); 3.58(s, 2H); 3.69-3.77(m, 2H); 6.22(s, 1H); 7.29-7.38(m, 3H)

5-Bromo-1-diethoxymethyloxindole (942mg, 3mmol) and sodium hydride (138mg, 5.8mmol, pre-washed with petroleum ether) were added successively to a solution of 4-chloro-6-methoxy-7-(2-methoxyethoxy)quinazoline (269mg, 1mmol), (prepared as described
25 for the starting material in Example 2), in THF (25ml) at ambient temperature. After stirring for 2 hours at ambient temperature, the volatiles were removed by evaporation and the residue dissolved in methylene chloride. The solution was washed with an aqueous saturated sodium chloride solution, dried (MgSO₄) and the solvent evaporated. The residue was purified by column flash chromatography using an increasingly polar mixture of methylene chloride and
30 ethyl acetate followed by a 7/3 mixture of methylene chloride/acetonitrile as eluent. The purified product was triturated with ether, collected by filtration, washed with a 1/1 mixture of

ether/petroleum ether and dried under vacuum to give 4-(5-bromo-1-diethoxymethyloxindol-3-yl)-6-methoxy-7-(2-methoxyethoxy)quinazoline (400mg, 73%) as a yellow-orange solid.

¹H NMR Spectrum: (CDCl₃) 1.25(t, 6H); 3.50(s, 3H); 3.52-3.60(m, 2H); 3.73-3.80(m, 2H); 3.89-3.91(m, 2H); 4.00(s, 3H); 4.35-4.37(m, 2H); 6.47(s, 1H); 7.19(d, 1H); 7.21(s, 1H);

5 7.48(s, 1H); 7.75(s, 1H); 7.81(dd, 1H); 8.11(s, 1H)

MS - ESI: 568 [MNa]⁺

Example 9

Sodium hydride (95mg of a 60% suspension in paraffin oil, 2.4mmol) was added to
10 a solution of 5-fluoro-1-diethoxymethyloxindole (520mg, 2.05mmol) and 4-chloro-6-methoxy-7-(2-methoxyethoxy)quinazoline (220mg, 0.82mmol), (prepared as described for the starting material in Example 2), in THF (20ml). After stirring for 2 hours at ambient temperature, the volatiles were removed by evaporation. The residue was partitioned between methylene chloride and water, the organic layer was washed with water and then brine, dried
15 (MgSO₄) and the solvent evaporated. The residue was purified by flash chromatography using an increasingly polar mixture of methylene chloride/ethyl acetate (100/0 to 0/100) as eluent to give **4-(5-fluoro-1-diethoxymethyloxindol-3-yl)-6-methoxy-7-(2-methoxyethoxy)quinazoline** (300mg, 68%) as a yellow solid.

m.p. 245-255°C

20 ¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 1.06(t, 6H); 3.36(s, 3H); 3.44(q, 4H); 3.78(t, 2H); 3.87(s, 3H); 4.33(t, 2H); 6.87-6.89(m, 1H); 7.20(s, 1H); 7.85-7.95(m, 1H); 8.0-8.05(dd, 1H); 8.20-8.25(m, 1H); 8.64(s, 1H); 9.38(s, 1H)

The starting material was prepared as follows:

25 A solution of 5-fluorooxindole (980mg, 6.5mmol), (prepared according to Synthesis 1993, 51), in triethyl orthoformate (25ml) was stirred and heated at reflux for 8 hours. The volatiles were removed by evaporation and the crude product purified by flash chromatography using petroleum ether/ether 7/3 as eluent to give 5-fluoro-1-diethoxymethyloxindole (637mg, 40%).

¹H NMR Spectrum: (CDCl₃) 1.23(t, 6H); 3.51-3.62(m, 4H); 3.69-3.77(m, 2H); 6.23(s, 1H); 6.92-6.98(m, 2H); 7.30-7.36(m, 1H)

Example 10

5 A solution of 4-(5-fluoro-1-diethoxymethyloxindol-3-yl)-6-methoxy-7-(2-methoxyethoxy)quinazoline (255mg, 0.52mmol), (prepared as described in Example 9), in ethanol (25ml) containing 2M hydrochloric acid (2.6ml) was heated at 80°C for 1 hour. The solid formed was collected by filtration, washed with ethanol and dried under vacuum. The solid was dissolved in hot DMF, the solution was diluted with ethyl acetate and thoroughly
10 washed with water. The organic layer was dried (MgSO₄), the volatiles removed by evaporation and the resulting solid was dried under vacuum at 70°C to give 4-(5-fluorooxindol-3-yl)-6-methoxy-7-(2-methoxyethoxy)quinazoline (49mg, 25%).

m.p. 277-280°C

¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 3.36(s, 3H); 3.78(t, 2H); 3.88(s, 3H); 4.34(t, 2H);
15 6.94-6.99(m, 2H); 7.30(s, 1H); 7.48(dd, 1H); 7.78(s, 1H); 8.83(s, 1H)

MS - ESI: 383 [MH]⁺

Elemental analysis: Found C 61.9 H 5.0 N 10.5

C₂₀H₁₈N₃O₄F 0.1 ethyl acetate Requires C 62.5 H 4.8 N 10.7%

20 Example 11

A solution of oxindole (1.2g, 9mmol) in THF (10ml) was added dropwise to a suspension of sodium hydride (360mg of a 60% suspension in paraffin oil, 9mmol) in THF (15ml) over 15 minutes under nitrogen. The resulting white suspension was stirred for 15 minutes at ambient temperature and a solution of 7-benzyloxy-4-chloro-6-methoxyquinazoline (900mg, 3mmol) in THF (10 ml) was added followed by DMF (5ml).
25 The resulting reddish solution was stirred for 30 minutes at ambient temperature followed by 1 hour at 65°C. The mixture was concentrated to half its volume by evaporation and was partitioned between ethyl acetate and water. The aqueous layer was separated, adjusted to pH7 with 2M hydrochloric acid and the resulting precipitate was collected by filtration,

washed with water, followed by ethyl acetate and ether and dried under vacuum to give 7-benzyloxy-6-methoxy-4-(oxindol-3-yl)quinazoline (1g, 83%) as a yellow solid.

m.p. 280-285°C

¹H NMR Spectrum: (DMSO-d₆, NaOD) 3.82(s, 3H); 5.23(s, 2H); 6.6-6.75(m, 3H); 7.05(s, 1H);
5 7.3-7.45(m, 3H); 7.52(d, 2H); 8.1(d, 1H); 8.5(s, 1H); 8.8(s, 1H)

MS - ESI m/z: 398 [MH]⁺

Elemental analysis:	Found	C 71.8	H 5.0	N 10.6
C ₂₄ H ₁₉ N ₃ O ₃ · 0.2H ₂ O	Requires	C 71.9	H 4.9	N 10.5%

10 The starting material was prepared as follows:

A mixture of 2-amino-4-benzyloxy-5-methoxybenzamide (10g, 0.04mol), (prepared according to J. Med. Chem. 1977, vol 20, 146-149), and Gold's reagent (7.4g, 0.05mol) in dioxane (100ml) was stirred and heated to reflux for 24 hours. Sodium acetate (3.02g, 0.037mol) and acetic acid (1.65ml, 0.029mol) were added to the reaction mixture and
15 it was heated for a further 3 hours. The volatiles were removed by evaporation, water was added to the residue, the solid was collected by filtration, washed with water and dried. Recrystallisation from acetic acid gave 7-benzyloxy-6-methoxy-3,4-dihydroquinazolin-4-one (8.7g, 84%).

A mixture of 7-benzyloxy-6-methoxy-3,4-dihydroquinazolin-4-one (2.82g, 0.01mol), thionyl chloride (40ml) and DMF (0.28ml) was stirred and heated at reflux for 1
20 hour. The excess thionyl chloride was removed by evaporation and by azeotropeing with toluene to give 7-benzyloxy-4-chloro-6-methoxyquinazoline hydrochloride (3.45g), which was subjected to an aqueous work-up with sodium hydroxide solution to give 7-benzyloxy-4-chloro-6-methoxyquinazoline.

25

Example 12

A mixture of 4-(5-benzyloxy-1-diethoxymethyl-oxindol-3-yl)-6-methoxy-7-(3-morpholinopropoxy)quinazoline (240mg, 0.38mmol) and TFA (8ml) was heated at reflux for 30 minutes. The TFA was removed by evaporation, the residue was triturated with ether, the
30 solid was collected by filtration, washed with ether and dried under vacuum. The orange solid

thus obtained was dissolved in a 1M hydrochloric acid solution (5ml, 5mmol) and heated at reflux for 40 minutes. After cooling, the reaction mixture was adjusted to pH2.3 with 6M sodium hydroxide and subjected to column chromatography on reverse phase silica gel (C18 10 μ) with a gradient of methanol in water (30 to 100%) as eluent. The fractions containing the compound were combined, the solution was adjusted to pH1.5 with concentrated hydrochloric acid and the solvents removed by evaporation. The residue was triturated with methanol/ether, the solid was collected by filtration, washed with ether and dried under vacuum at 40°C to give 4-(5-hydroxyoxindol-3-yl)-6-methoxy-7-(3-morpholinopropoxy)quinazoline as a hydrochloride salt (90mg, 45%) as an orange solid.

10 m.p. 230-245°C

¹H NMR Spectrum: (DMSO_d₆, TFA) 2.27-2.34(m, 2H); 3.11-3.18(m, 2H); 3.33(t, 2H); 3.53(d, 2H); 3.76(t, 2H); 3.89(s, 3H); 4.02(d, 2H); 4.31(t, 2H); 6.61(dd, 1H); 6.81(d, 1H); 7.15(d, 1H); 7.33(s, 1H); 7.74(s, 1H); 8.86(s, 1H)

MS - ESI: 451 [MH]⁺

15 Elemental analysis: Found C 52.9 H 6.2 N 9.9 Cl 9.5
C₂₄H₂₆N₄O₅ · 1.5H₂O · 1.5HCl Requires C 54.2 H 5.8 N 10.5 Cl 10.0%

The starting material was prepared as follows:

A mixture of 5-benzyloxyoxindole (1.88g, 7.9 mmol), (EP 0636608 A1), and triethyl orthoformate (180ml) was heated at reflux with stirring for 3 hours. The volatiles were removed by evaporation and the residue was purified by column flash chromatography using petroleum ether/ether (7/3) as eluent. Evaporation of the solvents lead to a crystalline material which was collected by filtration and dried under vacuum to give 5-benzyloxy-1-diethoxymethyloxindole (1.96g, 73%).

25 ¹H NMR Spectrum: (CDCl₃) 1.23(t, 6H); 3.51-3.58(m, 2H); 3.55(s, 2H); 3.68-3.76(m, 2H); 5.03(s, 2H); 6.22(s, 1H); 6.85(dd, 1H); 6.90(d, 1H); 7.30-7.44(m, 6H)

MS - ESI: 341 [MNa]⁺

Elemental analysis: Found C 70.0 H 4.2 N 7.0

C₂₀H₂₃NO₄ Requires C 70.4 H 4.1 N 6.8%

30 4-Chloro-6-methoxy-7-(2-methoxyethoxy)quinazoline (200mg, 0.59mmol),
(prepared as described for the starting material in Example 2), and sodium hydride (72mg,

- 73 -

3mmol, pre-washed with petroleum ether) were added successively to a solution of 5-benzyloxy-1-diethoxymethyloxindole (500mg, 1.47mmol) in THF (20ml) at ambient temperature. After stirring for 2 hours at ambient temperature, the volatiles were removed by evaporation and the residue was taken up in methylene chloride. The solution was washed
5 with saturated aqueous sodium chloride solution, dried (MgSO₄) and the solvent was removed by evaporation. The residue was purified by column flash chromatography using methylene chloride/ethyl acetate (1/1) followed by an increasingly polar mixture of methylene chloride/methanol (97/3 to 95/5) as eluent. The purified product was triturated with ether, collected by filtration, washed with ether and dried under vacuum to give 4-(5-benzyloxy-1-
10 diethoxymethyloxindol-3-yl)-6-methoxy-7-(3-morpholinopropoxy)quinazoline (340mg, 90%) as a yellow-orange solid.

¹H NMR Spectrum: (CDCl₃) 1.06(t, 6H); 2.25-2.31(m, 2H); 3.12-3.19(m, 2H); 3.36(t, 2H); 3.44(q, 4H); 3.55(d, 2H); 3.67(t, 3H); 3.86(s, 3H); 4.04(br d, 2H); 4.32(t, 2H); 5.10(s, 2H); 6.83(dd, 1H); 7.24(s, 1H); 7.33-7.36(br s, 1H); 7.40(t, 1H); 7.46(d, 1H); 7.58(br s, 1H);
15 7.95(d, 1H); 8.08(br s, 1H); 8.59(s, 1H); 9.36(s, 1H)

MS - ESI: 643 [MH]⁺

Example 13

A solution of 6-fluorooxindole (264mg, 1.75mmol), (prepared according to
20 Synthesis 1993, 51), in THF (3ml) was added dropwise under nitrogen to sodium hydride (42mg, 1.75mmol, pre-washed with hexane). After stirring the resulting mixture for 20 minutes at ambient temperature, 4-chloro-6,7-dimethoxyquinazoline (210mg, 0.93mmol), (prepared as described for the starting material in Example 1), was added as a solid, followed by DMF (4ml). The reaction mixture was then heated at 85°C for 1 hour, the solvent was
25 removed by evaporation and the residue partitioned between ether and water. The aqueous layer was separated and neutralised with 2M hydrochloric acid and the solid was collected by filtration, washed with ether and dried under vacuum to give 6,7-dimethoxy-4-(6-
fluorooxindol-3-yl)quinazoline (255mg, 81%) as a yellow solid.

m.p. 315°C (decomposition)

30 ¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 3.85(s, 3H); 3.98(s, 3H); 6.75-6.9(m, 2H); 7.24(s, 1H); 7.65(m, 1H); 7.75(s, 1H); 8.75(s, 1H)

- 74 -

MS: 340 [MH]⁺

Elemental analysis:	Found	C 62.4	H 4.4	N 12.2
C ₁₈ H ₁₄ N ₃ O ₃ F 0.23H ₂ O	Requires	C 63.0	H 4.2	N 12.2%

5 **Example 14**

A solution of 6-fluorooxindole (227mg, 1.5mmol), (prepared according to Synthesis 1993, 51), in THF (3ml) was added dropwise under nitrogen to sodium hydride (42mg, 1.75mmol, pre-washed with hexane). After stirring the resulting mixture for 20 minutes at ambient temperature, 4-chloro-6-methoxy-7-(3-morpholinopropoxy)quinazoline (169mg, 0.5mmol), (prepared as described for the starting material in Example 5), was added as a solid, followed by DMF (3ml). The mixture was then heated at 75°C for 1 hour, allowed to cool and the solvent removed by evaporation. The mixture was partitioned between water and ether. The aqueous layer was separated and adjusted to pH8 with 2M hydrochloric acid and then extracted with ethyl acetate. The organic layer was washed with brine, dried and volatiles removed by evaporation. The residue was triturated with ether and collected by filtration. This solid was dissolved in methylene chloride/methanol, a 5M solution of hydrogen chloride in isopropanol (0.5ml) was added and the volatiles removed by evaporation. The resulting solid was collected by filtration, washed with ether and dried under vacuum to give 4-(6-fluorooxindol-3-yl)-6-methoxy-7-(3-morpholinopropoxy)quinazoline hydrochloride (198mg, 81%) as a yellow solid.

m.p. 195-200°C

¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 2.25-2.4(m, 2H); 3.05-3.2(m, 2H); 3.25-3.35(m, 2H); 3.52(d, 2H); 3.72-3.85(m, 2H); 3.9(s, 3H); 4.05(d, 2H); 4.3(t, 2H); 6.8-6.9(m, 2H); 7.32(s, 1H); 7.62-7.7(m, 1H); 7.78(s, 1H); 8.7-8.8(m, 1H)

MS: 453[MH]⁺

Elemental analysis:	Found	C 54.1	H 5.7	N 10.6
C ₂₄ H ₂₃ N ₄ O ₄ F 0.8H ₂ O 1.8HCl	Requires	C 54.1	H 5.4	N 10.5%

Example 15

A solution of oxindole (319mg, 2.4mmol) in THF (3ml) was added dropwise under nitrogen to sodium hydride (58mg, 2.4mmol, pre-washed with hexane) and the mixture stirred

for 20 minutes at ambient temperature. A solution of 4-chloro-7-(2-methoxyethoxy)quinazoline (191mg, 0.8mmol) in DMF (3ml) was added and the mixture then heated at 70°C for 1 hour. The mixture was allowed to cool and the solvent removed by evaporation. The mixture was partitioned between water and ether. The aqueous phase was
5 separated and adjusted to pH7 with 2M hydrochloric acid. The resulting solid was collected by filtration and washed with water and ethanol. This solid was then suspended in ether, recollected by filtration, washed with ether and dried under vacuum to give 7-(2-methoxyethoxy)-4-(oxindol-3-yl)quinazoline (200mg, 75%).
m.p. 245-248°C

10 ¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 3.36(s, 3H); 3.75(t, 2H); 4.31(t, 2H); 6.92-7.05(m, 2H); 7.12-7.2(m, 2H); 7.3-7.4(dd, 1H); 7.65(d, 1H); 8.4(d, 1H); 8.8(s, 1H)
MS: 336 [MH]⁺

Elemental analysis:	Found	C 67.4	H 5.3	N 12.5
C ₁₉ H ₁₇ N ₃ O ₃ 0.07H ₂ O	Requires	C 67.8	H 5.1	N 12.5%

15

The starting material was prepared as follows:

A solution of 2-amino-4-fluorobenzoic acid (3g, 19.3mmol) in formamide (30ml) was heated at 150°C for 6 hours. The reaction mixture was poured onto ice/water 1/1 (250ml). The precipitated solid was collected by filtration, washed with water and dried to
20 give 7-fluoro-3,4-dihydroquinazolin-4-one (2.6g, 82%).

Sodium (400mg, 17mmol) was added carefully to 2-methoxyethanol (10ml) and the mixture heated at reflux for 30 minutes. 7-Fluoro-3,4-dihydroquinazolin-4-one (750mg, 4.57mmol) was added to the resulting solution and the mixture heated at reflux for 15 hours. The mixture was cooled and poured into water (250ml). The mixture was acidified to pH4
25 with concentrated hydrochloric acid. The resulting solid product was collected by filtration, washed with water and then with ether, and dried under vacuum to give 7-(2-methoxyethoxy)-3,4-dihydroquinazolin-4-one (580mg, 58%).

A solution of 7-(2-methoxyethoxy)-3,4-dihydroquinazolin-4-one (500mg, 2.2mmol) in thionyl chloride (15ml) and DMF (0.1ml) was heated at reflux for 3 hours. The volatiles
30 were removed by evaporation to give 4-chloro-7-(2-methoxyethoxy)quinazoline hydrochloride as a cream solid (520mg, 83%).

A suspension of 4-chloro-7-(2-methoxyethoxy)quinazoline hydrochloride (500mg, 1.8mmol) in a mixture of water (20ml) and ethyl acetate (20ml) was diluted with a saturated solution of sodium hydrogen carbonate. After stirring at ambient temperature for 15 minutes the solution was extracted with ethyl acetate. The organic layer was washed with brine, dried
 5 (MgSO₄) and evaporated to give 4-chloro-7-(2-methoxyethoxy)quinazoline (345mg, 80%).

Example 16

A solution of oxindole (266mg, 2.0mmol) in THF (2ml) was added dropwise under nitrogen to a stirred suspension of sodium hydride (80mg, of a 60% dispersion in mineral oil,
 10 2.0mmol) in THF (1ml). After stirring the resulting mixture for 15 minutes at ambient temperature, a solution of 4-chloro-6,7-dimethoxyquinazoline (449mg, 2.0mmol), (prepared as described for the starting material in Example 1), in THF (20ml) was added. The mixture was stirred at ambient temperature for 2 hours, quenched with water and the volatiles removed by evaporation. The resulting solid was purified by flash chromatography eluting with
 15 methylene chloride/methanol (19/1). The purified product was triturated with methanol, collected by filtration, washed with ether and dried under vacuum to give **6,7-dimethoxy-4-(oxindol-3-yl)quinazoline** (77mg, 12%).

¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 3.86(s, 3H); 3.98(s, 3H); 6.9-7.1(m, 4H); 7.22(s, 1H); 7.63(d, 1H); 7.79(s, 1H); 8.34(s, 1H)

20 MS: 322 [MH]⁺

Elemental analysis:	Found	C 66.6	H 4.4	N 12.7
---------------------	-------	--------	-------	--------

C ₁₈ H ₁₅ N ₃ O ₃ 0.1H ₂ O	Requires	C 66.9	H 4.7	N 13.0%
---	----------	--------	-------	---------

Example 17

25 A solution of 4-(7-benzyloxyoxindol-3-yl)-6-methoxy-7-(3-morpholinopropoxy)quinazoline (210mg, 0.39mmol) in TFA (15ml) and thioanisole (2.3ml, 19mmol) was stirred at ambient temperature for 20 hours. The volatiles were removed by evaporation and the residue was triturated with ether. The solid product was collected by filtration, washed with ether and dried under vacuum. The solid was dissolved in water (5ml)
 30 containing 2M hydrochloric acid (2 drops) and purified by reverse phase C18 chromatography eluting with water/methanol a (gradient from 80/20 to 30/70). The purified solid was

dissolved in methanol and 7.5M methanolic hydrogen chloride (0.2ml) was added. The volatiles were removed by evaporation, the solid collected and dried under vacuum at 45°C to give 4-(7-hydroxyoxindol-3-yl)-6-methoxy-7-(3-morpholinopropoxy)quinazoline hydrochloride (73mg, 35%).

- 5 ¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 2.2-2.35(m, 2H); 3.15(t, 2H); 3.34(t, 2H); 3.54(d, 2H); 3.7(t, 2H); 3.86(s, 3H); 4.01(d, 2H); 4.29(t, 2H); 6.73(d, 1H); 6.88(t, 1H); 7.17(d, 1H); 7.30(s, 1H); 7.80(s, 1H); 8.96(s, 1H)

MS - ESI: 451 [MH]⁺

Elemental analysis:		Found	C 54.1	H 5.8	N 10.5
10	C ₂₄ H ₂₆ N ₄ O ₅ · 0.3H ₂ O · 2HCl	Requires	C 54.5	H 5.5	N 10.6%

The starting material was prepared as follows:

- 7-Benzyloxy-3-methylthioxindole (610mg, 2.1mmol), (EP 8226), was dissolved in hot ethanol (60ml). The mixture was allowed to cool and Rainey Nickel (40ml volume,
15 prewashed with ethanol) was added and the mixture was stirred at ambient temperature for 1.5 hours. The insolubles were removed by filtration and the filter pad was thoroughly washed with ethanol. The solvent was removed from the filtrate by evaporation and the residue was purified by column chromatography eluting with ethyl acetate/petroleum ether (40/60) to give 7-benzyloxyoxindole (395mg, 71%).

- 20 ¹H NMR Spectrum: (CDCl₃) 3.55(s, 2H); 5.1(s, 2H); 6.86(d, 2H); 6.95(t, 1H); 7.35-7.45(m, 5H); 7.52(br s, 1H)

MS - ESI: 262 [MNa]⁺

Elemental analysis: Found C 75.5 H 5.8 N 5.8

C₁₅H₁₃NO₂ Requires C 75.3 H 5.5 N 5.8%

- 25 Sodium hydride (36mg, 1.5mmol) was added in portions to a solution of 7-benzyloxyoxindole (360mg, 1.5mmol) in THF (8ml), the mixture was stirred for 30 minutes at ambient temperature and a solution of 4-chloro-6-methoxy-7-(3-morpholinopropoxy)quinazoline (204mg, 0.6mmol), (prepared as described for the starting material in Example 5), in THF (4ml) and DMF (1.5ml) was added. The mixture was stirred
30 for 30 minutes at ambient temperature followed by 1.5 hours at 65°C. The THF was removed by evaporation, the mixture was partitioned between ethyl acetate and water and the aqueous

layer was adjusted to pH8 with 2M hydrochloric acid. The precipitated product was collected by filtration, washed with water, ethyl acetate, and ether, and dried under vacuum to give 4-(7-benzyloxyoxindol-3-yl)-6-methoxy-7-(3-morpholinopropoxy)quinazoline (190mg, 58%).

¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 2.2-2.35(m, 2H); 3.15(t, 2H); 3.34(t, 2H); 3.54(d, 2H); 3.71(t, 2H); 3.85(s, 3H); 4.05(d, 2H); 4.3(t, 2H); 5.26(s, 2H); 6.9-7.0(m, 2H); 7.22-7.38(m, 3H); 7.4(t, 2H); 7.59(d, 2H); 7.79(s, 1H); 8.8(s, 1H)

MS - ESI: 541 [MH]⁺

Elemental analysis:	Found	C 66.4	H 6.0	N 10.1
---------------------	-------	--------	-------	--------

C ₃₁ H ₃₂ N ₄ O ₅ 1H ₂ O	Requires	C 66.6	H 6.1	N 10.0%
---	----------	--------	-------	---------

Example 18

A solution of oxindole (200mg, 1.5mmol) in THF (3ml) was added dropwise under nitrogen to sodium hydride (60mg, 1.5mmol, prewashed with hexane) in THF (3ml). After stirring for 20 minutes at ambient temperature, a solution of 4-chloro-6-(3-morpholinopropoxy)-7-methoxyquinazoline (169mg, 0.5mmol) in DMF (3ml) was added. The mixture was heated at 75°C for 1 hour, allowed to cool and the volatiles removed by evaporation. The mixture was partitioned between water and ether. The aqueous layer was adjusted to pH8 with 2M hydrochloric acid. The resulting precipitate was collected by filtration, washed with water, and dried under vacuum. The solid was dissolved in methylene chloride/methanol and 3M ethereal hydrogen chloride (0.5ml) was added. The mixture was diluted with ether, the solid was collected by filtration and dried under vacuum to give 7-methoxy-6-(3-morpholinopropoxy)-4-(oxindol-3-yl)quinazoline hydrochloride (175mg, 69%).

¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 2.3(m, 2H); 3.1(t, 2H); 3.3(t, 2H); 3.5(d, 2H); 3.75(t, 2H); 3.99(s, 3H); 3.95-4.05(m, 2H); 4.15(t, 2H); 7.0(d, 1H); 7.05(t, 1H); 7.15(t, 1H); 7.33(s, 1H); 7.7(d, 1H); 7.8(s, 1H); 8.81(s, 1H)

MS - ESI: 435 [MH]⁺

Elemental analysis:	Found	C 56.0	H 6.1	N 10.8
---------------------	-------	--------	-------	--------

C ₂₄ H ₂₆ N ₄ O ₄ 1H ₂ O 1.8HCl	Requires	C 55.6	H 5.8	N 10.8%
--	----------	--------	-------	---------

The starting material was prepared as follows:

A suspension of 4-(3-chloro-4-fluoroanilino)-6-(3-morpholinopropoxy)-7-methoxyquinazoline (6.0g 13.4mmol), (WO 96 33980), in 6M hydrochloric acid (120ml) was
5 heated at reflux for 6 hours. The mixture was cooled to 0°C and carefully neutralised with cooling by addition of concentrated aqueous ammonia. The resulting precipitate was collected by filtration, washed with dilute aqueous ammonia and water and dried under vacuum to give 7-methoxy-6-(3-morpholinopropoxy)-3,4-dihydroquinazolin-4-one (4.2g).

¹H NMR Spectrum: (DMSO-d₆) 2.4(m, 6H); 3.59(t, 4H); 3.75(t, 2H); 3.90(s, 3H); 4.12(t, 2H);
10 7.12(s, 1H); 7.43(s, 1H); 7.98(s, 1H); 12.0(br s, 1H)

MS - ESI: 320 [MH]⁺

Elemental analysis:	Found	C 58.6	H 6.5	N 12.7
C ₁₆ H ₂₁ N ₃ O ₄ 0.5H ₂ O	Requires	C 58.5	H 6.7	N 12.8%

A solution of 7-methoxy-6-(3-morpholinopropoxy)-3,4-dihydroquinazolin-4-one
15 (990mg, 3.1mmol) in thionyl chloride (10ml) and DMF (0.1ml) was heated at 80°C for 1.5 hours. The mixture was allowed to cool, toluene was added and the solvent was removed by evaporation. The residue was partitioned between ethyl acetate and water and the aqueous layer was adjusted to pH7.5 with 2M sodium hydroxide solution. The organic layer was separated, washed with brine, dried (MgSO₄) and the solvent removed by evaporation. The
20 residue was purified by flash chromatography eluting with methylene chloride/methanol (1/9 followed by 95/5). The purified solid was triturated with hexane, collected by filtration and washed with ether to give 4-chloro-7-methoxy-6-(3-morpholinopropoxy)quinazoline (614mg, 58%).

¹H NMR Spectrum: (CDCl₃) 2.12(m, 2H); 2.50(br s, 4H); 2.59(t, 2H); 3.73(t, 4H); 4.05(s,
25 3H); 4.27(t, 2H); 7.33(s, 1H); 7.40(s, 1H); 8.86(s, 1H)

Example 19

A solution of 5,6-dimethoxyoxindole (107mg, 0.56mmol), (Chem Pharm Bull. 1971, 19, 1325-1327), in a mixture of THF (1ml) and DMF (1ml) was added dropwise under

nitrogen to a suspension of sodium hydride (22mg, 0.55mmol, prewashed with hexane) in THF (1ml). The mixture was stirred for 20 minutes at ambient temperature and a solution of 4-chloro-6-methoxy-7-(2-methoxyethoxy)quinazoline (50mg, 0.18mmol), (prepared as described for the starting material in Example 2), in THF (1ml) was added dropwise. The mixture was heated at 60°C for 1.5 hours, allowed to cool and partitioned between ethyl acetate and water. The aqueous layer was adjusted to pH7 with 1M hydrochloric acid. The organic layer was washed with brine, dried (MgSO₄), and volatiles removed by evaporation. The residue was purified by column flash chromatography eluting with methylene chloride/acetonitrile/methanol (60/38/2 followed by 60/35/5). The resulting solid was dissolved in methylene chloride/methanol and 3M ethereal hydrogen chloride was added. The mixture was concentrated by evaporation, the resulting precipitate was collected by filtration, washed with ether and dried under vacuum to give 4-(5,6-dimethoxyoxindol-3-yl)-6-methoxy-7-(2-methoxyethoxy)quinazoline hydrochloride (33mg, 51%).

¹HNMR Spectrum: (DMSO-d₆, CF₃CO₂D) 3.35(s, 3H); 3.67(s, 3H); 3.78(m, 2H); 3.82(s, 3H); 3.86(s, 3H); 4.3(m, 2H); 6.67(s, 1H); 7.20(s, 1H); 7.27(s, 1H); 7.67(s, 1H); 8.91(s, 1H)

MS - ESI: 426 [MH]⁺

Elemental analysis:	Found	C 55.4	H 5.5	N 8.6
C ₂₂ H ₂₃ N ₃ O ₆ · 0.7H ₂ O · HCl	Requires	C 55.7	H 5.4	N 8.9%

20 Example 20

A solution of 5-cyanooxindole (285mg, 1.8mmol), (Tet. Lett., 1987, 28, 4027), in THF (12ml) was added dropwise under nitrogen, to a suspension of sodium hydride (96mg, 2.4mmol, prewashed with hexane) in THF (5ml). The mixture was stirred for 20 minutes at ambient temperature, and a solution of 4-chloro-6-methoxy-7-(2-methoxyethoxy)quinazoline (162mg, 0.6mmol), (prepared as described for the starting material in Example 2), in DMF (0.5ml) in THF (3ml) was added dropwise. The mixture was stirred for 20 minutes at ambient temperature and then 1 hour at 65°C. The solvent was removed by evaporation and the residue was purified by column flash chromatography eluting with methylene chloride/methanol (97/3 followed by 96/4 and 90/10). The solid was collected by filtration, resuspended in methylene chloride and 2M ethereal hydrogen chloride was added. The solid

was collected by filtration, washed with ether and dried under vacuum at 60°C to give 4-(5-cyanooxindol-3-yl)-6-methoxy-7-(2-methoxyethoxy)quinazoline hydrochloride (92mg, 40%).

¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 3.36(s, 3H); 3.8(m, 2H); 3.90(s, 3H); 4.34(t, 2H);

5 7.09(d, 1H); 7.28(s, 1H); 7.46(d, 1H); 7.94(s, 1H); 8.1(s, 1H); 8.70(s, 1H)

MS - ESI: 391 [MH]⁺

Elemental analysis:	Found	C 62.6	H 4.7	N 14.0
---------------------	-------	--------	-------	--------

C ₂₁ H ₁₈ N ₄ O ₄ 0.1 HCl 0.4H ₂ O	Requires	C 62.9	H 4.7	N 14.0%
---	----------	--------	-------	---------

10 Example 21

A solution of 5-nitrooxindole (400mg, 2.2mmol) in THF (4ml) was added to sodium hydride (89mg, 3.7mmol, prewashed with THF) in THF (4ml). The mixture was stirred for 30 minutes at ambient temperature and a solution of 4-chloro-6-methoxy-7-(2-methoxyethoxy)quinazoline (200mg, 0.75mmol), (prepared as described for the starting material in Example 2), in THF (3ml) and DMF (1ml) was added dropwise. The mixture was stirred for 1 hour at ambient temperature, followed by 1.5 hours at 70°C. The solvent was removed by evaporation and the residue was partitioned between ethyl acetate and water. The aqueous layer was adjusted to pH7 with 2M hydrochloric acid. The precipitated product was collected by filtration, washed with ether and dried under vacuum at 60°C. The solid was dissolved in methanol (10ml) and 7.5M hydrogen chloride in methanol (2 ml) was added. The volatiles were removed by evaporation, the solid residue was triturated with ether, collected by filtration and dried under vacuum to give 6-methoxy-7-(2-methoxyethoxy)-4-(5-nitrooxindol-3-yl)-quinazoline hydrochloride (173mg, 57%).

¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 3.36(s, 3H); 3.79(t, 2H); 3.87(s, 3H); 4.35(t, 2H);

25 7.1(d, 1H); 7.31(s, 1H); 7.95(br s, 1H); 8.01(d, 1H); 8.56(br s, 1H); 8.66(s, 1H)

MS - ESI: 411 [MH]⁺

The starting material was prepared as follows:

Fuming nitric acid (1.58ml, 39.4mmol) was added dropwise to a solution of oxindole (5g, 37.5mmol) in concentrated sulphuric acid (25ml) cooled by an ice bath and

30

maintained below 5°C. The mixture was stirred for 30 minutes at 0°C and then poured into water. The precipitate was collected by filtration and thoroughly washed with water. The solid was suspended in 50% acetic acid (100ml) and heated at 90°C for 1 hour. The mixture was allowed to cool, the solid product was collected by filtration, washed with water and dried under vacuum over phosphorus pentoxide at 40°C to give 5-nitrooxindole (3.2g, 48%).

¹H NMR Spectrum: (CDCl₃, CD₃CO₂D) 3.68(s, 2H); 7.05(d, 1H); 8.16(s, 1H); 8.22(s, 1H)

MS - EI: 178 [M]⁺

Elemental analysis:	Found	C 53.6	H 3.5	N 15.7
---------------------	-------	--------	-------	--------

C ₈ H ₆ N ₂ O ₃	Requires	C 53.9	H 3.39	N 15.7%
---	----------	--------	--------	---------

Example 22

A solution of oxindole (164mg, 1.23mmol) in THF (3ml) was added dropwise under nitrogen, to sodium hydride (49mg, 1.23mmol, prewashed with hexane) in THF (3ml). The mixture was stirred for 30 minutes at ambient temperature and DMF (2ml) and then 4-chloro-7-(2-(imidazol-1-yl)ethoxy)-6-methoxyquinazoline (125mg, 0.41mmol) was added. The mixture was heated at 60°C for 30 minutes and the THF was removed by evaporation. The residue was partitioned between saturated aqueous ammonium chloride solution and ether. The organic layer was separated, dried (MgSO₄) and the solvent removed by evaporation. The solid was dissolved in methylene chloride (5ml), methanol (0.5ml) and isopropanol (3ml) and 5M isopropanolic hydrogen chloride (0.5ml) was added. The mixture was concentrated by evaporation to 2ml total volume and the resulting precipitated solid was collected by filtration, washed with isopropanol followed by ether and dried under vacuum to give 7-(2-(imidazol-1-yl)ethoxy)-6-methoxy-4-(oxindol-3-yl)quinazoline hydrochloride (110mg, 61%).

¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 3.85(s, 3H); 4.62(m, 2H); 4.75(m, 2H); 7.0-7.1(m, 2H); 7.16(t, 1H); 7.35(s, 1H); 7.68(d, 1H); 7.73(s, 1H); 7.77(s, 1H); 7.84(s, 1H); 8.83(s, 1H); 9.2(s, 1H)

MS - ESI: 402 [MH]⁺

Elemental analysis:	Found	C 55.7	H 5.3	N 13.1
---------------------	-------	--------	-------	--------

C ₂₂ H ₁₉ N ₃ O ₃ 1H ₂ O 1.6HCl	Requires	C 55.7	H 5.5	N 13.3%
--	----------	--------	-------	---------

0.8isopropanol

The starting material was prepared as follows:

Sodium hydride (1.44g of a 60% suspension in mineral oil, 36mmol) was added in portions over 20 minutes to a solution of 7-benzyloxy-6-methoxy-3,4-dihydroquinazolin-4-one (8.46g, 30mmol), (prepared as described for the starting material in Example 11), in DMF (70ml) and the mixture was stirred for 1.5 hours. Chloromethyl pivalate (5.65g, 37.5mmol) was added dropwise and the mixture stirred for 2 hours at ambient temperature. The mixture was diluted with ethyl acetate (100ml) and poured onto ice/water (400ml) and 2M hydrochloric acid (4ml). The organic layer was separated and the aqueous layer extracted with ethyl acetate, the combined extracts were washed with brine, dried (MgSO₄) and the solvent removed by evaporation. The residue was triturated with a mixture of ether and petroleum ether, the solid was collected by filtration and dried under vacuum to give 7-benzyloxy-6-methoxy-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (10g, 84%).

¹H NMR Spectrum: (DMSO-d₆) 1.11(s, 9H); 3.89(s, 3H); 5.3(s, 2H); 5.9(s, 2H); 7.27(s, 1H); 7.35(m, 1H); 7.47(t, 2H); 7.49(d, 2H); 7.51(s, 1H); 8.34(s, 1H)

A mixture of 7-benzyloxy-6-methoxy-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (7g, 17.7mmol) and 10% palladium-on-charcoal catalyst (700mg) in ethyl acetate (250ml), DMF (50ml), methanol (50ml) and acetic acid (0.7ml) was stirred under hydrogen at atmospheric pressure for 40 minutes. The catalyst was removed by filtration and the solvent removed from the filtrate by evaporation. The residue was triturated with ether, collected by filtration and dried under vacuum to give 7-hydroxy-6-methoxy-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (4.36g, 80%).

¹H NMR Spectrum: (DMSO-d₆) 1.1(s, 9H); 3.89(s, 3H); 5.89(s, 2H); 7.0(s, 1H); 7.48(s, 1H); 8.5(s, 1H)

Diethyl azodicarboxylate (435mg, 2.5mmol) was added dropwise to a suspension of 7-hydroxy-6-methoxy-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (612mg, 2mmol), 2-(imidazol-1-yl)ethanol (280mg, 2.5mmol), (J. Med. Chem. 1993, 25 4052-4060), and triphenylphosphine (655mg, 2.5mmol) in methylene chloride (10ml) at 5°C. The mixture was stirred for 10 minutes at 5°C and then 1 hour at ambient temperature. The mixture was poured directly on to a silica column and eluted with methylene chloride/methanol (95/5) to

give 7-(2-(imidazol-1-yl)ethoxy)-6-methoxy-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (640mg, 80%).

¹H NMR Spectrum: (CDCl₃) 1.19(s, 9H); 3.98(s, 3H); 4.34(m, 2H); 4.45(m, 2H); 5.94(s, 2H); 7.02(s, 1H); 7.07(s, 1H); 7.11(s, 1H); 7.64(s, 1H); 7.67(s, 1H); 8.17(s, 1H)

5 MS - ESI: 423 [MNa]⁺

Elemental Analysis:	Found	C 58.3	H 6.4	N 13.9
---------------------	-------	--------	-------	--------

C ₂₀ H ₂₄ N ₄ O ₅ · 0.7H ₂ O	Requires	C 58.2	H 6.2	N 13.6%
---	----------	--------	-------	---------

A solution of 7-(2-(imidazol-1-yl)ethoxy)-6-methoxy-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (640mg, 1.6mmol) in saturated methanolic ammonia (10ml) was stirred for 15 hours at ambient temperature. The volatiles were removed by evaporation, the solid was triturated with ether, collected by filtration and dried under vacuum to give 7-(2-(imidazol-1-yl)ethoxy)-6-methoxy-3,4-dihydroquinazolin-4-one (412mg, 90%).

¹H NMR Spectrum: (DMSO-d₆) 3.89(s, 3H); 4.4-4.5(m, 4H); 6.9(s, 1H); 7.16(s, 1H); 7.28(s, 1H); 7.47(s, 1H); 7.7(s, 1H); 7.99(s, 1H)

15 MS - ESI: 287 [MH]⁺

Elemental Analysis:	Found	C 57.8	H 5.2	N 19.3
---------------------	-------	--------	-------	--------

C ₁₄ H ₁₄ N ₄ O ₃ · 0.3H ₂ O	Requires	C 57.7	H 5.1	N 19.2%
---	----------	--------	-------	---------

A mixture of 7-(2-(imidazol-1-yl)ethoxy)-6-methoxy-3,4-dihydroquinazolin-4-one (412mg, 1.44mmol), thionyl chloride (5 ml) and DMF (0.2ml) was heated at reflux for 1 hour. The mixture was diluted with toluene and the volatiles were removed by evaporation. The residue was suspended in methylene chloride, cooled to 0°C and aqueous sodium hydrogen carbonate solution was added. The resulting precipitate was collected by filtration and dried under vacuum to give 4-chloro-7-(2-(imidazol-1-yl)ethoxy)-6-methoxyquinazoline (258mg, 59%).

25 ¹H NMR Spectrum: (DMSO-d₆) 4.01(s, 3H); 4.47(m, 2H); 4.53(m, 2H); 6.89(s, 1H); 7.27(s, 1H); 7.41(s, 1H); 7.49(s, 1H); 7.70(s, 1H); 8.88(s, 1H)

MS - ESI: 327 [MNa]⁺

Example 23

A solution of oxindole (232mg, 1.74mmol) in THF (3ml) was added dropwise under nitrogen, to sodium hydride (70mg, 1.74mmol, prewashed with hexane) in THF (3ml). The mixture was stirred for 30 minutes at ambient temperature and 4-chloro-6-methoxy-7-(4-pyridylmethoxy)quinazoline (175mg, 0.58mmol) and then DMF (2ml) was added. The mixture was heated at 60°C for 1 hour and the THF was removed by evaporation. The residue was partitioned between cooled 2M hydrochloric acid and ethyl acetate. The organic layer was extracted with 2M hydrochloric acid. The aqueous layers were combined, washed with ethyl acetate and ether, basified with aqueous sodium hydrogen carbonate and extracted with methylene chloride. The organic layer was washed with brine, dried (MgSO₄) and the solvent removed by evaporation. The residue was purified by chromatography on neutral aluminium oxide eluting with methylene chloride/methanol (90/10 followed by 80/20 and 75/25). The purified solid product was dissolved in methanol/methylene chloride and 5M methanolic hydrogen chloride (0.5ml) was added. The volatiles were removed by evaporation, the residue was resuspended in ether, collected by filtration, washed with ether and dried under vacuum to give 6-methoxy-4-(oxindol-3-yl)-7-(4-pyridylmethoxy)quinazoline hydrochloride (50mg, 21%).

¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 3.93(s, 3H); 5.75(s, 2H); 7-7.1(m, 2H); 7.18(t, 1H); 7.35(s, 1H); 7.75(d, 1H); 7.88(s, 1H); 8.17(d, 2H); 8.8(m, 1H); 9.05(d, 2H)

MS - ESI: 399 [MH]⁺

The starting material was prepared as follows:

Diethyl azodicarboxylate (348mg, 2mmol) was added dropwise to a suspension of 7-hydroxy-6-methoxy-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (457mg, 1.5mmol), (prepared as described for the starting material in Example 22), 4-pyridinemethanol (218mg, 2mmol), and triphenylphosphine (524mg, 2mmol) in methylene chloride (7ml) at 5°C. The mixture was stirred for 10 minutes at 5°C and then 1 hour at ambient temperature. The mixture was poured directly on to a silica column and eluted with methylene chloride/methanol (95/5) to give 6-methoxy-3-((pivaloyloxy)methyl)-7-(4-pyridylmethoxy)-3,4-dihydroquinazolin-4-one (510mg, 85%).

¹H NMR Spectrum: (DMSO-d₆) 1.11(s, 9H); 3.93(s, 3H); 5.38(s, 2H); 5.9(s, 2H); 7.23(s, 1H); 7.46(d, 2H); 7.55(s, 1H); 8.35(s, 1H); 8.60(d, 2H)

MS - ESI: 420 [MNa]⁺

	Elemental Analysis:	Found	C 63.2	H 6.0	N 10.6
5	C ₂₁ H ₂₃ N ₃ O ₅	Requires	C 63.5	H 5.8	N 10.6%

A solution of 6-methoxy-3-((pivaloyloxy)methyl)-7-(4-pyridylmethoxy)-3,4-dihydroquinazolin-4-one (500mg, 1.26mmol) in saturated methanolic ammonia (10ml) was stirred for 20 hours at ambient temperature. The volatiles were removed by evaporation, the solid residue was triturated with ether, collected by filtration and dried under vacuum to give

10 6-methoxy-7-(4-pyridylmethoxy)-3,4-dihydroquinazolin-4-one (330mg, 92%).

¹H NMR Spectrum: (DMSO-d₆) 3.91(s, 3H); 5.5(s, 2H); 7.19(s, 1H); 7.4-7.5(m, 3H); 7.97(s, 1H); 8.6(d, 2H)

MS - EI: 283 [M]⁺

	Elemental analysis:	Found	C 61.9	H 5.1	N 14.8
15	C ₁₅ H ₁₃ N ₃ O ₃ 0.4H ₂ O	Requires	C 62.0	H 4.8	N 14.5%

A suspension of 6-methoxy-7-(4-pyridylmethoxy)-3,4-dihydroquinazolin-4-one (300mg, 1.06mmol) in thionyl chloride (5ml) containing DMF (50μl) was heated at reflux for 1 hour. The mixture was allowed to cool, diluted with toluene and the volatiles were removed by evaporation. The residue was dissolved in methylene chloride and a cooled solution of

20 aqueous sodium hydrogen carbonate was added. The resulting precipitate was collected by filtration, washed with ether and dried under vacuum to give 4-chloro-6-methoxy-7-(4-pyridylmethoxy)quinazoline (175mg, 55%).

¹H NMR Spectrum: (DMSO-d₆) 4.05(s, 3H); 5.55(s, 2H); 7.47(s, 1H); 7.55(s, 1H); 7.64(d, 2H); 8.7(d, 2H); 8.89(s, 1H)

25 MS - ESI: 302 [MH]⁺

Example 24

Sodium hydride (62mg, 2.7mmol) was added to a solution of 1-diethoxymethyloxindole (317mg, 1.35mmol), (prepared as described for the starting material

30 in Example 2), in THF (10ml). The mixture was stirred for 10 minutes at ambient

temperature and 4-chloro-6-methoxy-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline (145mg, 0.45mmol) in THF (5ml) was added. The mixture was then stirred for 2 hours at ambient temperature, and was partitioned between methylene chloride and water. The organic layer was separated, washed with brine, dried (MgSO₄) and the solvent removed by evaporation.

- 5 The residue was purified by column chromatography eluting with methylene chloride/methanol (a gradient from 95/5 to 50/50). The purified product was triturated with ether, collected by filtration, washed with ether and dried under vacuum to give 4-(1-diethoxymethyloxindol-3-yl)-6-methoxy-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline (218mg, 94%).

- 10 ¹H NMR Spectrum: (CDCl₃, CD₃CO₂D) 1.26(t, 6H); 2.05-2.15(m, 4H); 2.4(m, 2H); 3.3-3.5(m, 6H); 3.6(q, 2H); 3.8(q, 2H); 3.87(s, 3H); 4.3(t, 2H); 6.47(s, 1H); 7.05(t, 1H); 7.15(t, 1H); 7.29(s, 1H); 7.63(d, 2H); 7.8(s, 1H); 8.3(s, 1H)
MS - ESI: 521 [MH]⁺

- 15 The starting material was prepared as follows:

- A mixture of 4-hydroxy-3-methoxybenzoic acid (8.4g, 50mmol), 3-(pyrrolidin-1-yl)propyl chloride (14.75g, 0.1mol), (prepared according to J. Am. Chem. Soc. 1955, 77, 2272), potassium carbonate (13.8g, 0.1mol) and potassium iodide (1.66g, 10mmol) in DMF (150ml) was stirred and heated at 100°C for 3 hours. The mixture was allowed to cool and
20 the insolubles were removed by filtration and the volatiles were removed from the filtrate by evaporation. The residue was dissolved in ethanol (75ml), 2M aqueous sodium hydroxide (75ml) was added and the mixture was heated at 90°C for 2 hours. The mixture was concentrated by evaporation, acidified with concentrated hydrochloric acid, washed with ether and then subjected to purification on a Diaion (trade mark of Mitsubishi) HP20SS resin
25 column, eluting with water and then with a gradient of methanol (0 to 25%) in dilute hydrochloric acid (pH2.2). The methanol was removed by evaporation and the aqueous residue was freeze dried to give 3-methoxy-4-(3-(pyrrolidin-1-yl)propoxy)benzoic acid hydrochloride (12.2g, 77%).

- ¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 2.2(m, 2H); 3.15(t, 2H); 3.3(t, 2H); 3.5(d, 2H);
30 3.7(t, 2H); 3.82(s, 3H); 4.05(d, 2H); 4.15(t, 2H); 7.07(d, 1H); 7.48(s, 1H); 7.59(d, 1H)
MS - EI: 279 [M]⁺

Fuming nitric acid (2.4ml, 57.9mmol) was added slowly at 0°C to a solution of 3-methoxy-4-(3-(pyrrolidin-1-yl)propoxy)benzoic acid (12.15g, 38.17mmol) in TFA (40ml). The cooling bath was removed and the reaction mixture stirred at ambient temperature for 1 hour. The TFA was removed by evaporation and ice/water was added to the residue and the solvent removed by evaporation. The solid residue was dissolved in dilute hydrochloric acid (pH2.2), poured onto a Diaion (trade mark of Mitsubishi) HP20SS resin column and eluted with methanol (gradient 0 to 50%) in water. Concentration of the fractions by evaporation gave a precipitate which was collected by filtration and dried under vacuum over phosphorus pentoxide to give 5-methoxy-2-nitro-4-(3-(pyrrolidin-1-yl)propoxy)benzoic acid hydrochloride (12.1g, 90%).

¹H NMR Spectrum: (DMSO-d₆, TFA) 1.8-1.9 (m, 2H); 2.0-2.1(m, 2H); 2.1-2.2(m, 2H); 3.0-3.1(m, 2H); 3.3(t, 2H); 3.6-3.7(m, 2H); 3.95(s, 3H); 4.25(t, 2H); 7.35(s, 1H); 7.62(s, 1H)

A solution of 5-methoxy-2-nitro-4-(3-(pyrrolidin-1-yl)propoxy)benzoic acid hydrochloride (9.63g, 24mmol) in thionyl chloride (20ml) and DMF (50μl) was heated at 45°C for 1.5 hours. The excess thionyl chloride was removed by evaporation and by azeotroping with toluene (x2). The resulting solid was suspended in THF (250ml) and methylene chloride (100ml) and ammonia was bubbled through the mixture for 30 minutes and the mixture stirred for a further 1.5 hours at ambient temperature. The volatiles were removed by evaporation, the residue was dissolved in water and applied to a Diaion (trade mark of Mitsubishi) HP20SS resin column and eluted with water/methanol (100/0 to 95/5). The solvent was removed by evaporation from the fractions containing product and the residue was dissolved in a minimum of methanol and the solution was diluted with ether. The resulting precipitate was collected by filtration, washed with ether and dried under vacuum to give 5-methoxy-2-nitro-4-(3-(pyrrolidin-1-yl)propoxy)benzamide (7.23g, 73%).

¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 1.85-1.95(m, 2H); 2-2.1(m, 2H); 2.15-2.25(m, 2H); 3.0-3.1(m, 2H); 3.31(t, 2H); 3.62(t, 2H); 3.93(s, 3H); 4.2(t, 2H); 7.16(s, 1H); 7.60(s, 1H)
MS - EI: 323 [M]⁺

Concentrated hydrochloric acid (5ml) was added to a suspension of 5-methoxy-2-nitro-4-(3-(pyrrolidin-1-yl)propoxy)benzamide (1.5g, 4.64mmol) in methanol (20ml) and the mixture was heated at 50°C to give a solution. Iron powder (1.3g, 23.2mmol) was added in portions and the reaction mixture was then heated at reflux for 1 hour. The mixture was

allowed to cool, the insolubles were removed by filtration through diatomaceous earth and the volatiles were removed from the filtrate by evaporation. The residue was purified on a Diaion (trade mark of Mitsubishi) HP20SS resin column, eluting with water and then with dilute hydrochloric acid (pH2). The fractions containing product were concentrated by evaporation and the resulting precipitate was collected by filtration and dried under vacuum over phosphorus pentoxide to give 2-amino-5-methoxy-4-(3-(pyrrolidin-1-yl)propoxy)benzamide hydrochloride (1.44g, 85%).

¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 1.9(br s, 2H); 2.05(br s, 2H); 2.2(br s, 2H); 3.05(br s, 2H); 3.3(t, 2H); 3.61(br s, 2H); 3.8(s, 3H); 4.11(t, 2H); 7.05(s, 1H); 7.53(s, 1H)

MS - EI: 293 [M]⁺

A mixture of 2-amino-5-methoxy-4-(3-(pyrrolidin-1-yl)propoxy)benzamide hydrochloride (5.92g, 16.2mmol) and Gold's reagent (3.5g, 21.4mmol) in dioxane (50ml) was heated at reflux for 5 hours. Acetic acid (0.7ml) and sodium acetate (1.33g) were added to the reaction mixture which was heated at reflux for a further 5 hours. The mixture was allowed to cool and the volatiles were removed by evaporation. The residue was dissolved in water, adjusted to pH8 with 2M aqueous sodium hydroxide solution and purified on a Diaion (trademark of Mitsubishi) HP20SS resin column eluting with methanol (gradient 0-50 %) in water. The fractions containing product were concentrated by evaporation and then freeze dried to give 4-hydroxy-6-methoxy-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline (4.55g, 83%).

¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 1.9(m, 2H); 2.0-2.1(m, 2H); 2.2-2.3(m, 2H); 3.05(m, 2H); 3.34(t, 2H); 3.6-3.7(br s, 2H); 3.94(s, 3H); 4.27(t, 2H); 7.31(s, 1H); 7.55(s, 1H); 9.02(s, 1H)

A mixture of 4-hydroxy-6-methoxy-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline (1.7g, 5mmol) and thionyl chloride (25ml) containing DMF (0.2ml) was heated at reflux for 3 hours. Excess thionyl chloride was removed by evaporation and by azeotropeing with toluene (x2). The residue was suspended in ether and 10 % aqueous solution of sodium hydrogen carbonate was added to the mixture. The organic layer was separated, dried (MgSO₄) and the solvent removed by evaporation to give 4-chloro-6-methoxy-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline (1.94g, quantitative).

¹H NMR Spectrum: (CDCl₃) 1.8(br s, 4H); 2.17(m, 2H); 2.6(br s, 4H); 2.7(t, 2H); 4.05(s, 3H); 4.3(t, 2H); 7.35(s, 1H); 7.38(s, 1H); 8.86(s, 1H)

MS - ESI: 322 [MH]⁺

Example 25

2M Hydrochloric acid (1.5ml, 3mmol) was added to a solution of 4-(1-
 5 diethoxymethyloxindol-3-yl)-6-methoxy-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline (151mg,
 0.29mmol), (prepared as described in Example 24), in ethanol (15ml) at 80°C and the mixture
 was stirred at 80°C for 1.5 hours. The reaction mixture was allowed to cool, the solvent was
 removed by evaporation and the residue was purified by reverse phase C18 chromatography
 eluting with a gradient of acetonitrile in water (0 - 40%). The acetonitrile was removed by
 10 evaporation from the fractions containing product and concentrated hydrochloric acid (0.1ml)
 was added to the aqueous phase. The solution was freeze dried to give **6-methoxy-4-**
(oxindol-3-yl)-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline hydrochloride (150mg, 100%).
¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 1.85-1.95(m, 2H); 2.0-2.1(m, 2H); 2.2-2.3(m, 2H);
 3.0-3.1(m, 2H); 3.3-3.4(t, 2H); 3.6-3.7(m, 2H); 3.87(s, 3H); 4.3(t, 2H); 7.0-7.1(m, 2H); 7.15-
 15 7.2(t, 1H); 7.33(s, 1H); 7.68(d, 1H); 7.8(s, 1H); 8.9(s, 1H)

MS - ESI: 419 [MH]⁺

Elemental analysis:	Found	C 53.9	H 6.4	N 10.7
C ₂₄ H ₂₆ N ₄ O ₃ 3H ₂ O 1.6HCl	Requires	C 54.3	H 6.3	N 10.7%

20 Example 26

A solution of 5-cyanooxindole (213mg, 1.35mmol), (Tet. Lett., 1987, 28, 4027), in
 DMF (3ml) was added dropwise to a suspension of sodium hydride (54mg, 1.35mmol,
 prewashed with hexane) in DMF (2ml). The mixture was stirred for 20 minutes at ambient
 temperature, 4-chloro-6-methoxy-7-(3-morpholinopropoxy)quinazoline (152mg, 0.45mmol),
 25 (prepared as described for the starting material in Example 5), was added and the mixture was
 heated at 70°C for 1 hour. The mixture was allowed to cool and then was diluted with water
 and adjusted to pH1 with 2M hydrochloric acid. The solution was washed with ether and then
 basified with aqueous sodium hydrogen carbonate solution. The resulting precipitate was
 collected by filtration, washed with water, ether and dried under vacuum. The solid was
 30 purified by flash chromatography eluting with methylene chloride/methanol (90/10 followed
 by 80/20). The purified solid was dissolved in methylene chloride/methanol and 5M

- 91 -

methanolic hydrogen chloride (0.5ml) was added. The solution was concentrated by evaporation to a total volume of 2ml and ether (15ml) was added. The resulting solid was collected by filtration, washed with ether and dried under vacuum to give **4-(5-cyanooxindol-3-yl)-6-methoxy-7-(3-morpholinopropoxy)quinazoline hydrochloride** (120mg, 50%).

5 ¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 2.32(m, 2H); 3.15(t, 2H); 3.35(t, 2H); 3.54(d, 2H); 3.75(t, 2H); 3.91(s, 3H); 4.05(d, 2H); 4.33(t, 2H); 7.10(d, 1H); 7.31(s, 1H); 7.47(d, 1H); 7.97(s, 1H); 8.06(s, 1H); 8.67(s, 1H)

MS - ESI: 460 [MH]⁺

Elemental analysis:	Found	C 54.9	H 5.4	N 12.9
10 C ₂₅ H ₂₃ N ₃ O ₄ 1.1H ₂ O 1.75HCl	Requires	C 55.3	H 5.4	N 12.9%

Example 27

A solution of 5-nitrooxindole (1.26g, 7.1mmol), (prepared as described for the starting material in Example 21), in THF (10ml) was added dropwise to sodium hydride
 15 (284mg, 7.1mmol, prewashed with THF) in THF (15ml). The mixture was stirred for 30 minutes at ambient temperature and a solution of 4-chloro-6-methoxy-7-(3-morpholinopropoxy)quinazoline (600mg, 1.8mmol, (prepared as described for the starting material in Example 5), in THF (10ml) and DMF (3ml) was added dropwise. The mixture was stirred for 30 minutes at ambient temperature, then heated at 75°C for 1.5 hours. DMF
 20 (4ml) was added and the mixture stirred for 12 hours at ambient temperature. The solvent was removed by evaporation, the residue was partitioned between ethyl acetate and water and the aqueous layer was adjusted to pH8 with 2M hydrochloric acid. The precipitated product was collected by filtration, washed with water and dried under vacuum. The solid was dissolved in a mixture of DMF and dilute hydrochloric acid (pH2) and was purified on a Diaion (trade
 25 mark of Mitsubishi) HP20SS resin column, eluting with a gradient of methanol and dilute hydrochloric acid (pH2). The purified solid was dissolved in methanol and 7M ethanolic hydrogen chloride (1.5ml) was added. The volatiles were removed by evaporation and the solid residue was triturated with a mixture of methanol and ether. The solid was collected by
 30 filtration and dried under vacuum to give **6-methoxy-7-(3-morpholinopropoxy)-4-(5-nitrooxindol-3-yl)-quinazoline hydrochloride** (275mg, 29%).

¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 1.9-2.1(m, 2H); 2.9(t, 2H); 3.06(t, 2H); 3.25(d, 2H); 3.41(t, 2H); 3.57(s, 3H); 3.72(d, 2H); 4.04(t, 2H); 6.8(d, 1H); 7.02(s, 1H); 7.65-7.75(m, 2H); 8.2(br s, 1H); 8.31(s, 1H)

MS - ESI: 480 [MH]⁺

5	Elemental analysis:	Found	C 50.5	H 5.5	N 11.9
	C ₂₄ H ₂₅ N ₃ O ₆ 2.5H ₂ O 1.3HCl	Requires	C 50.4	H 5.5	N 12.2%

Example 28

A solution of oxindole (259mg, 1.95mmol) in THF (3ml) was added dropwise to a suspension of sodium hydride (78mg, 1.95mmol, prewashed with hexane) in THF (3ml). The mixture was stirred for 30 minutes at ambient temperature and a solution of 4-chloro-6-methoxy-7-(2-(pyrrolidin-1-yl)ethoxy)quinazoline (200mg, 0.65mmol) in DMF (3ml) was added dropwise. The mixture was heated at 60°C for 30 minutes, the THF was removed by evaporation and the residue was partitioned between methylene chloride and water. The organic layer was separated, washed with brine, dried (MgSO₄) and the solvent removed by evaporation. The solid residue was purified by chromatography on neutral aluminium oxide eluting with methylene chloride/methanol (80/20). The purified solid was dissolved in methylene chloride containing methanol and 3M ethereal hydrogen chloride was added. The volatiles were removed by evaporation and the residue was suspended in ether. The solid was collected by filtration, washed with ether and dried under vacuum to give 6-methoxy-4-(oxindol-3-yl)-7-(2-(pyrrolidin-1-yl)ethoxy)quinazoline hydrochloride (55mg, 18%).

¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 1.85-1.95(m, 2H); 2.05-2.2(m, 2H); 3.15-3.3(m, 2H); 3.65-3.8(m, 4H); 3.88(s, 3H); 4.57(m, 2H); 7.0-7.1(m, 2H); 7.16(t, 1H); 7.38(s, 1H); 7.68(d, 1H); 7.81(s, 1H); 8.82(br s, 1H)

25 MS - ESI: 405 [MH]⁺

Elemental Analysis:	Found	C 56.1	H 5.9	N 11.0
C ₂₃ H ₂₄ N ₄ O ₃ 1H ₂ O 1.95HCl	Requires	C 56.0	H 5.7	N 11.3%

The starting material was prepared as follows:

Diethyl azodicarboxylate (566mg, 3.25mmol) was added dropwise to a solution of 7-hydroxy-6-methoxy-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (765mg, 2.5mmol), (prepared as described for the starting material in Example 22), 1-(2-hydroxyethyl)pyrrolidine (374mg, 3.25mmol) and triphenylphosphine (851mg, 3.25mmol) in methylene chloride (15ml). The mixture was stirred for 1 hour at ambient temperature and further triphenylphosphine (861mg, 3.25mmol) followed by diethyl azodicarboxylate (566mg, 3.25mmol) was added. The mixture was stirred for 1 hour at ambient temperature and the solvent was removed by evaporation. The residue was dissolved in ethyl acetate/ether and extracted with 2M hydrochloric acid. The aqueous layer was basified by addition of solid sodium hydrogen carbonate and extracted with methylene chloride. The combined extracts were dried (MgSO₄), and the solvent removed by evaporation. The resulting oil was purified by column chromatography on neutral aluminium oxide eluting with methylene chloride/methanol (90/10 followed by 80/20) to give 6-methoxy-3-((pivaloyloxy)methyl)-7-(2-(pyrrolidin-1-yl)ethoxy)-3,4-dihydroquinazolin-4-one (550mg, 55%).

¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 1.12(s, 9H); 1.75-1.85(m, 2H); 2-2.1(m, 2H); 3.15-3.25(m, 2H); 3.65-3.75(m, 4H); 3.93(s, 3H); 4.50(t, 2H); 5.93(s, 2H); 7.31(s, 1H); 7.57(s, 1H); 8.41(s, 1H)

MS - ESI: 404 [MH]⁺

A solution of 6-methoxy-3-((pivaloyloxy)methyl)-7-(2-(pyrrolidin-1-yl)ethoxy)-3,4-dihydroquinazolin-4-one (524mg, 1.3mmol) in methanol saturated with ammonia (10ml) was stirred for 20 hours at ambient temperature. The volatiles were removed by evaporation and the solid residue was triturated with ether, collected by filtration and dried under vacuum to give 6-methoxy-7-(2-(pyrrolidin-1-yl)ethoxy)-3,4-dihydroquinazolin-4-one (230mg, 61%).

¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 1.9-2.0(m, 2H); 2.05-2.2(m, 2H); 3.15-3.3(m, 2H); 3.6-3.8(m, 4H); 3.98(s, 3H); 4.53(t, 2H); 7.38(s, 1H); 7.61(s, 1H); 9.13(s, 1H)

A solution of 6-methoxy-7-(2-(pyrrolidin-1-yl)ethoxy)-3,4-dihydroquinazolin-4-one (228mg, 7.9mmol) in thionyl chloride (5ml) and DMF (0.2ml) was heated at reflux for 1 hour. Toluene was added and the volatiles were removed by evaporation. The residue was dissolved in methylene chloride and cooled aqueous sodium hydrogen carbonate solution was added. The organic layer was separated, washed with brine, dried (MgSO₄) and the solvent

removed by evaporation to give 4-chloro-6-methoxy-7-(2-(pyrrolidin-1-yl)ethoxy)quinazoline (200mg, 82%).

MS - [ESI]: 307 [M.]⁺

5 **Example 29**

A solution of 1-acetyloxindole (251mg, 1.6mmol), (J. Med. Chem. 1986, 1838), in DMF (3ml) was added to a suspension of sodium hydride (50mg, 2.4mmol) in DMF (10ml) and the mixture stirred for 10 minutes at ambient temperature. 4-Chloro-6-methoxy-7-(2-methoxyethoxy)quinazoline (108mg, 0.4mmol), (prepared as described for the starting material in Example 2), in DMF (3ml) was added and the mixture stirred for 1.5 hours at ambient temperature. The solvent was removed by evaporation and the residue was partitioned between methylene chloride and water. The organic layer was separated, washed with water, brine, dried (MgSO₄) and the solvent removed by evaporation. The residue was purified by column chromatography eluting with methylene chloride/ethyl acetate (70/30). The purified product was recrystallised from methylene chloride/ether, collected by filtration, washed with ether and dried under vacuum to give 4-(1-acetyloxindol-3-yl)-6-methoxy-7-(2-methoxyethoxy)quinazoline (77mg, 47%).

¹H NMR Spectrum: (CDCl₃) 2.82(s, 3H); 3.49(s, 3H); 3.87(s, 3H); 3.90(m, 2H); 4.36(m, 2H); 7.1-7.2(m, 2H); 7.22(s, 1H); 7.61(d, 1H); 7.76(s, 1H); 8.16(s, 1H); 8.37(d, 1H)

20 MS - EI: 407 [M.]⁺

Elemental analysis:	Found	C 64.5	H 5.4	N 10.4
C ₂₂ H ₂₁ N ₃ O ₅	Requires	C 64.9	H 5.2	N 10.3%

Example 30

25 A solution of 5-acetamidooxindole (212mg, 1.1mmol), (Zh. Obshch. Khim 1956, 26, 2019-2022), in DMF (2ml) was added dropwise to sodium hydride (89mg, 2.23mmol, prewashed with THF) in DMF (2ml). The mixture was stirred for 5 minutes at ambient temperature and a solution of 4-chloro-6-methoxy-7-(2-methoxyethoxy)quinazoline (100mg, 0.37mmol), (prepared as described for the starting material in Example 2), in DMF (2ml) was added. The mixture was stirred for 1.5 hours at ambient temperature, the solvent was removed by evaporation and the mixture was partitioned between methylene chloride and

30

water. The aqueous layer was adjusted to pH7 with 2M hydrochloric acid. The organic layer was separated, dried (MgSO_4) and the solvent was removed by evaporation. The residue was purified by chromatography eluting with methanol/methylene chloride (gradient from 5/95 to 15/85). The purified solid was dissolved in methylene chloride and 4M methanolic hydrogen chloride (4ml) was added. The volatiles were removed by evaporation, the solid residue was triturated with ether, collected by filtration, washed with ether and dried under vacuum at 60°C to give 4-(5-acetamidooxindol-3-yl)-6-methoxy-7-(2-methoxyethoxy)quinazoline hydrochloride (95mg, 56%).

¹H NMR Spectrum: (DMSO-d_6 , $\text{CF}_3\text{CO}_2\text{D}$) 2.01(s, 3H); 3.36(s, 3H); 3.78(t, 2H); 3.84(s, 3H); 4.33(t, 2H); 6.9(d, 1H); 7.3(s, 1H); 7.34(d, 1H); 7.68(s, 1H); 8.17(s, 1H); 8.9(s, 1H)

MS - EI: 422 [M]⁺

Elemental analysis:	Found	C 54.4	H 5.1	N 11.9
$\text{C}_{22}\text{H}_{22}\text{N}_4\text{O}_5 \cdot 1.3\text{H}_2\text{O} \cdot \text{HCl}$	Requires	C 54.8	H 5.3	N 11.6%

15 **Example 31**

A solution of 5-methylthiooxindole (668mg, 3.7mmol) in DMF (3ml) was added to a suspension of sodium hydride (149mg, 3.7mmol, prewashed with THF) in DMF (3ml). The mixture was stirred for 30 minutes at ambient temperature and a solution of 4-chloro-6-methoxy-7-(3-morpholinopropoxy)quinazoline (420mg, 1.2mmol), (prepared as described for the starting material in Example 5), in DMF (3ml) was added dropwise. The mixture was stirred for 1.5 hours at ambient temperature and the solvent was removed by evaporation. The residue was partitioned between water and ether and the aqueous layer was adjusted to pH8 with 2M hydrochloric acid. The precipitated product was collected by filtration, washed with water and dried under vacuum over phosphorus pentoxide. The solid was purified by column chromatography eluting with methylene chloride/methanol (95/5 followed by 90/10). The purified solid was dried under vacuum to give the product as the free base (450mg, 75%). The free base product (100mg) was dissolved in methylene chloride/methanol and a solution of 2.5M ethereal hydrogen chloride (1.5ml) was added. The volatiles were removed by evaporation to give 6-methoxy-4-(5-methylthiooxindol-3-yl)-7-(3-morpholinopropoxy)quinazoline hydrochloride (108mg).

- 96 -

¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 2.25-2.35(m, 2H); 2.42(s, 3H); 3.15(t, 2H); 3.35(t, 2H); 3.54(d, 2H); 3.75(t, 2H); 3.91(s, 3H); 4.05(d, 2H); 4.32(t, 2H); 7.01(d, 1H); 7.15(d, 1H); 7.33(s, 1H); 7.63(s, 1H); 7.79(s, 1H); 8.8(s, 1H)

MS - EI: 480 [M]⁺

5	Elemental analysis:	Found	C 53.3	H 5.8	N 9.7
	C ₂₅ H ₂₈ N ₄ O ₄ S 1.5H ₂ O 1.55HCl	Requires	C 53.2	H 5.8	N 9.9%

The starting material was prepared as follows:

10 A solution of methyl 5-fluoro-2-nitrophenylacetate (3g, 14mmol), (Synthesis 1993, 51), in DMSO (10ml) was added to a suspension of sodium thiomethoxide (1.08g, 15.5mmol) in anhydrous DMSO (10ml). The mixture was stirred for 10 minutes at ambient temperature and further sodium thiomethoxide (100mg, 1.5mmol) was added. The mixture was stirred for 15 minutes at ambient temperature and was then partitioned between ethyl acetate and water. The organic layer was separated, washed with water, brine, dried (MgSO₄) and the solvent
15 removed by evaporation. The residue was purified by column chromatography eluting with ethyl acetate/petroleum ether (15/85) to give methyl 5-methylthio-2-nitrophenylacetate (2.68g, 81%).

¹H NMR Spectrum: (CDCl₃) 2.54(s, 3H); 3.71(s, 3H); 4.01(s, 2H); 7.09(s, 1H); 7.23(d, 1H); 8.10(d, 1H)

20 MS - EI: 240 [M]⁺

Iron powder (2.36g, 42mmol) was added in portions to a solution of methyl 5-methylthio-2-nitrophenylacetate (2.55g, 10.6mmol) in acetic acid (30ml) and the mixture was heated at 100°C for 1 hour. The mixture was allowed to cool and the insolubles were removed by filtration and the filter pad was thoroughly washed with acetic acid and then ethyl
25 acetate. The volatiles were removed from the filtrate by evaporation and the residue was dissolved in ethyl acetate and was washed with 1M hydrochloric acid (50ml x3), water, brine and dried (MgSO₄). The solvent was removed by evaporation and the residue was purified by column chromatography eluting with ethyl acetate/petroleum ether (1/1) to give 5-methylthiooxindole (1.48g, 78%).

30 m.p. 120-122°C

¹H NMR Spectrum: (CDCl₃) 2.46(s, 3H); 3.53(s, 2H); 6.81(d, 1H); 7.15-7.30(m, 2H); 8.21(br s, 1H)

MS - EI: 179 [M]⁺

Elemental analysis:	Found	C 60.2	H 5.2	N 7.8
---------------------	-------	--------	-------	-------

5	C ₉ H ₉ NOS	Requires	C 60.3	H 5.1	N 7.8%
---	-----------------------------------	----------	--------	-------	--------

Example 32

3-Chloroperoxybenzoic acid (186mg, 0.78mmol) was added to a solution of 6-methoxy-4-(5-methylthiooxindol-3-yl)-7-(3-morpholinopropoxy)quinazoline (170mg, 10 0.35mmol), (prepared as described for the free base product in Example 31), in methylene chloride (40ml) and the mixture stirred for 2 hours at ambient temperature. The mixture was diluted with water (5ml) and ether (15ml) and the aqueous layer was adjusted to pH2 by addition of 1M hydrochloric acid. The aqueous layer was separated and applied to a reverse phase C18 column and eluted with a gradient of methanol/1% aqueous acetic acid (0/100 to 15 100/0). The purified product was dissolved in methanol and concentrated hydrochloric acid (3drops) was added. The volatiles were removed by evaporation, the solid residue was collected and dried under vacuum to give 6-methoxy-4-(5-methylsulphinyloxindol-3-yl)-7-(4-oxidomorpholinopropoxy)quinazoline hydrochloride (100mg, 60%).

¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 2.45(m, 2H); 2.67(s, 3H); 2.7-2.9(m, 4H); 3.94(s, 20 3H); 3.94-3.99(br s, 6H); 4.37(t, 2H); 7.13(d, 1H); 7.34(d, 1H); 7.36(s, 1H); 7.78(s, 1H); 8.0(s, 1H); 8.76(s, 1H)

MS - ESI: 513 [MH]⁺

Example 33

25 5-methylsulphonyloxindole (350mg, 1.66mmol) was added to a suspension of sodium hydride (66mg, 1.66mmol) in DMF (5ml) and the mixture was stirred for 30 minutes at ambient temperature. 4-Chloro-6-methoxy-7-(2-methoxyethoxy)quinazoline (148.5mg, 0.55mmol), (prepared as described for the starting material in Example 2), was added as a solid followed by DMF (1ml). The mixture was heated at 50°C for 1.5 hours and then 30 allowed to cool. The mixture was partitioned between water and ether and the aqueous layer was adjusted to pH7 with 2M hydrochloric acid. The precipitated product was collected by

filtration, washed with water, ether and dried under vacuum. The solid was adsorbed on silica and purified by column chromatography eluting with methylene chloride/methanol/acetonitrile (60/5/35). The purified solid was dissolved in methylene chloride/methanol and 7M ethanolic hydrogen chloride (1ml) was added. The volatiles were removed by evaporation, the solid residue triturated with ether, collected by filtration, washed with ether and dried under vacuum to give **6-methoxy-7-(2-methoxyethoxy)-4-(5-methylsulphonyloxindol-3-yl)quinazoline hydrochloride** (170mg, 70%).

¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 3.1(s, 3H); 3.36(s, 3H); 3.78(m, 2H); 3.93(s, 3H); 4.36(t, 2H); 7.18(d, 1H); 7.33(s, 1H); 7.64(d, 1H); 7.77(s, 1H); 8.13(s, 1H); 8.72(s, 1H)

MS - ESI: 444 [MH]⁺

Elemental analysis:	Found	C 51.0	H 4.9	N 8.8
C ₂₁ H ₁₂ N ₃ O ₆ S 1H ₂ O 0.85HCl	Requires	C 51.2	H 4.9	N 8.5%

The starting material was prepared as follows:

3-Chloroperoxybenzoic acid (919mg, 3.8mmol) was added to a solution of 5-methylthiooxindole (343mg, 1.9mmol), (prepared as described for the starting material in Example 31), in methylene chloride (100ml) cooled at 5°C and the mixture stirred for 2 hours at ambient temperature. Further 3-chloroperoxybenzoic acid (100mg) was added and the mixture stirred for a further 1 hour. The volatiles were removed by evaporation, the residue was triturated with ether, collected by filtration and dried under vacuum to give 5-methylsulphonyloxindole (363mg, 90%).

¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 3.14(s, 3H); 3.60(s, 2H); 7.01(d, 1H); 7.7-7.8(m, 2H)

Example 34

Sodium hydride (58mg, 2.4mmol) was added to solution of 5-(2-morpholinoacetyl)oxindole (312mg, 1.2mmol) and 4-chloro-6-methoxy-7-(2-methoxyethoxy)quinazoline (160mg, 0.6mmol), (prepared as described for the starting material in Example 2), in DMF (15ml). The mixture was stirred for 1.5 hours at ambient temperature and the volatiles were removed by evaporation. The residue was purified by column chromatography eluting with methylene chloride/methanol (96/4 followed by 94/6).

- 99 -

The purified solid product was triturated with ether, collected by filtration, washed with ether and dried under vacuum. The solid was dissolved in methylene chloride (10ml) and methanol (10ml) and 1M methanolic hydrogen chloride (2ml) was added. The volatiles were removed by evaporation and the solid residue was suspended in ether, collected by filtration, washed
 5 with ether and dried under vacuum to give 6-methoxy-7-(2-methoxyethoxy)-4-(5-(2-morpholinoacetyl)oxindol-3-yl)quinazoline hydrochloride (136mg, 39%).
¹H NMR Spectrum: (DMSO-d₆, CD₃CO₂D) 3.25(t, 2H); 3.36(s, 3H); 3.46(d, 2H); 3.78(m, 2H); 3.8-3.9(m, 2H); 3.86(s, 3H); 4.0(d, 2H); 4.35(m, 2H); 5.08(s, 2H); 7.15(d, 1H); 7.35(s, 1H); 7.78(d, 1H); 7.85(s, 1H); 8.31(s, 1H); 8.73(s, 1H)

10 MS - ESI: 493 [MH]⁺

Elemental analysis:	Found	C 53.5	H 5.8	N 9.7
C ₂₆ H ₂₈ N ₄ O ₆ 2H ₂ O 1.5HCl	Requires	C 53.5	H 5.8	N 9.6%

The starting material was prepared as follows:

15 Potassium carbonate (800mg, 5.8mmol) followed by potassium iodide (30mg, 0.2mmol) and morpholine (500μl, 5.7mmol) were added to a solution of 5-(2-chloroacetyl)oxindole (1.05g, 5mmol), (J. Med. Chem. 1991, 1860), in DMF (20ml) and the mixture was stirred for 1 hour at ambient temperature. The solvent was removed by evaporation and the residue was purified by reverse phase column chromatography on a
 20 Diaion (trade mark of Mitsubishi) HP20SS resin column, eluting with water/methanol (a gradient from 100/0 to 30/70). The purified product was then purified by chromatography on silica eluting with methylene chloride/methanol (95/5) to give 5-(2-morpholinoacetyl)oxindole (673mg, 51%).
¹H NMR Spectrum: (DMSO-d₆, CD₃CO₂D) 3.57(s, 2H); 5.06(s, 2H); 6.93(d, 1H);
 25 7.83(s, 1H); 7.9(d, 1H)

Example 35

Using a procedure analogous to that described in Example 49, 5-nitrooxindole (240mg, 1.4mmol), (prepared as described for the starting material in Example 21), was added
 30 to sodium hydride (80mg, 1.9mmol) in THF (15ml) and the solution was treated with 4-chloro-7-(2-methoxyethoxy)quinazoline (170mg, 0.7mmol), (prepared as described for the

starting material in Example 15), in DMF (5ml) to give, after work-up and purification, 7-(2-methoxyethoxy)-4-(5-nitrooxindol-3-yl)quinazoline (87mg, 32%).

¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 3.3(s, 3H); 3.7(t, 2H); 4.25(t, 2H); 7.0(d, 1H); 7.15(d, 1H); 7.22(dd, 1H); 7.95(dd, 1H); 8.5(d, 1H); 8.65(s, 1H); 8.75(s, 1H); 11.4(s, 1H)

5 MS - ESI: 381 [MH]⁺

Elemental analysis:	Found	C 57.9	H 4.1	N 14.7
---------------------	-------	--------	-------	--------

C ₁₉ H ₁₆ N ₄ O ₃ · 0.7H ₂ O	Requires	C 58.1	H 4.5	N 14.3%
---	----------	--------	-------	---------

Example 36

10 Using a procedure analogous to that described in Example 49, 5-cyanooxindole (250mg, 1.6mmol), (Tet. Lett., 1987, 28, 4027), was added to sodium hydride (100mg, 2.5mmol) in THF (15ml) and the solution was treated with 4-chloro-7-(2-methoxyethoxy)quinazoline (200mg, 0.84mmol), (prepared as described for the starting material in Example 15), in DMF (5ml) to give, after work-up and purification, 4-(5-cyanooxindol-3-yl)-7-(2-methoxyethoxy)quinazoline (160mg, 53%).

¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 3.3(s, 3H); 3.7(t, 2H); 4.3(t, 2H); 7.05(d, 1H); 7.15(d, 1H); 7.3(dd, 1H); 7.45(d, 1H); 7.95(s, 1H); 8.45(d, 1H); 8.8(s, 1H); 11.3(s, 1H)

MS - ESI: 359 [M-H]⁺

Elemental analysis:	Found	C 64.6	H 4.5	N 15.4
---------------------	-------	--------	-------	--------

20 C ₂₀ H ₁₆ N ₄ O ₃ · 0.6H ₂ O	Requires	C 64.7	H 4.7	N 15.1%
--	----------	--------	-------	---------

Example 37

25 Using a procedure analogous to that described in Example 26, oxindole (240mg, 1.8mmol) was added to sodium hydride (110mg, 2.75mmol) in THF (15ml) and the solution was treated with 4-chloro-7-(3-morpholinopropoxy)quinazoline (400mg, 1.3mmol) in DMF (5ml) to give, after work-up and purification, 7-(3-morpholinopropoxy)-4-(oxindol-3-yl)quinazoline (160mg, 30%).

¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 2.2(m, 2H); 3.1(t, 2H); 3.3(t, 2H); 3.5(d, 2H); 3.7(t, 2H); 4.0(m, 2H); 4.25(t, 2H); 6.92(d, 1H); 7.1(d, 1H); 7.15(d, 1H); 7.22(dd, 1H), 7.6(d, 1H); 8.4(d, 1H); 8.6(s, 1H), 10.05(br s, 1H), 10.9(br s, 1H)

MS - ESI: 405 [MH]⁺

5	Elemental analysis:	Found	C 65.9	H 5.9	N 13.9
	C ₂₃ H ₂₄ N ₄ O ₃ · 0.7H ₂ O	Requires	C 66.2	H 6.1	N 13.4%

The starting material was prepared as follows:

10 A solution of 2-amino-4-fluorobenzoic acid (3g, 19.3mmol) in formamide (30ml) was heated at 150°C for 6 hours. The reaction mixture was poured onto ice/water (1/1) (250ml). The precipitated solid was collected by filtration, washed with water and dried to give 7-fluoro-3,4-dihydroquinazolin-4-one (2.6g, 82%).

15 Sodium metal (4.4g, 191mmol) was added to benzyl alcohol (100ml) at ambient temperature and the mixture stirred for 30 minutes then heated at 80°C for 1 hour. The mixture was then cooled to 40°C and 7-fluoro-3,4-dihydroquinazolin-4-one (7.8g, 47mmol), was added. The reaction mixture was stirred at 130°C for 4 hours and allowed to cool to ambient temperature and stirred for 18 hours. The solution was quenched with water (800ml) and acidified to pH3 with concentrated hydrochloric acid. The resulting precipitate was collected by filtration, washed with water and ether and dried for 4 hours at 60°C under high
20 vacuum to give 7-benzyloxy-3,4-dihydroquinazolin-4-one (7.02g, 63 %).

7-Benzyloxy-3,4-dihydroquinazolin-4-one (7.0g, 27mmol) was suspended in dry DMF (50ml) and cooled to 5-10°C, then sodium hydride was added (1.22g of a 60 % suspension in mineral oil, 30mmol). The reaction mixture was allowed to return to ambient temperature and chloromethyl pivalate (4.75g, 31.5mmol) was added over 10 minutes and the
25 mixture stirred for 1 hour. The reaction mixture was quenched with water (250ml), the aqueous phase adjusted to pH5 and extracted with ether (300ml x3). The combined ether phases were washed with brine, dried (MgSO₄) and the solvent removed by evaporation. The solid residue was triturated with isohexane, collected by filtration and dried under vacuum to give 7-benzyloxy-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (9.1g, 90%).

7-benzyloxy-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (8.0g, 22mmol) was dissolved in TFA (40ml) and the mixture heated at reflux for 3 hours then allowed to cool to ambient temperature and stirred for 18 hours. The TFA was removed by evaporation and the residue resuspended in a mixture of ether and aqueous sodium hydrogen carbonate solution. The solid was collected by filtration, washed with water and ether and dried at 40°C for 3 hours under high vacuum to give 7-hydroxy-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (5.42g, 90%).

Morpholine (94g, 1.08mol) was added dropwise to a solution of 3-bromo-1-propanol (75g, 0.54mol) in toluene (750ml) and the reaction then heated at 80°C for 4 hours. The mixture was allowed to cool to ambient temperature and the precipitated solid was removed by filtration. The volatiles were removed from the filtrate and the resulting yellow oil was purified by distillation at 0.4-0.7 mmHg to give 4-(3-hydroxypropyl)morpholine (40g, 50%) as a colourless oil.

b.p. 68-70°C (~0.5mmHg)

¹H NMR Spectrum: (DMSO-d₆) 1.65-1.78(m, 2H); 2.50(t, 4H); 2.60(t, 2H); 3.68(t, 4H); 3.78(t, 2H); 4.90(br d, 1H)

7-Hydroxy-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (750mg, 2.7mmol) was suspended in methylene chloride (40ml) at 5°C and 4-(3-hydroxypropyl)morpholine (490mg, 3.4mmol) and triphenylphosphine (890mg, 3.4mmol) were added. The mixture was stirred for 5 minutes and diethyl azodicarboxylate (590mg, 3.4mmol) was added over 5 minutes at 5°C. The reaction mixture was stirred at 5°C for 30 minutes then at ambient temperature for 1 hour. The solution was then purified directly by column flash chromatography eluting with methylene chloride, and then ethyl acetate, acetonitrile/ethyl acetate (20/80), and acetonitrile/ethyl acetate/ammonia (50/50/0.5). The purified product was triturated with ether/isohexane and collected by filtration to give 7-(3-morpholinopropoxy)-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (745mg, 68%).

7-(3-Morpholinopropoxy)-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (680mg, 1.6mmol) was stirred in methanolic ammonia (20ml) at 40°C for 6 hours then for 18 hours at ambient temperature. The solvent was removed by evaporation, the residue was triturated with ether/isohexane and collected by filtration to give 7-(3-morpholinopropoxy)-3,4-dihydroquinazolin-4-one (450mg, 92%) as a white solid.

¹H NMR Spectrum: (DMSO-d₆) 1.9(quin, 2H); 2.35(t, 4H); 2.4(t, 2H); 3.55(t, 4H); 4.15(t, 2H); 7.05(m, 2H); 7.97(d, 1H); 8.02(s, 1H)

MS: 290 [MH]⁺

A mixture of 7-(3-morpholinopropoxy)-3,4-dihydroquinazolin-4-one (500mg, 1.7mmol), thionyl chloride (10ml) and DMF (0.1ml) was heated at reflux for 2 hours. The volatiles were removed by evaporation and the residue azeotroped with toluene. The crude product was partitioned between methylene chloride (50ml) and saturated aqueous sodium hydrogen carbonate solution (50ml) and the mixture stirred for 10 minutes. The organic phase was separated and the aqueous phase extracted with methylene chloride. The combined extracts were dried (MgSO₄) and the solvent removed by evaporation to give 4-chloro-7-(3-morpholinopropoxy)quinazoline (425mg, 80%).

Example 38

A solution of 5-cyanooxindole (284mg, 1.8mmol), (Tet. Lett., 1987, 28, 4027), in DMF (4ml) was added dropwise, to sodium hydride (72mg, 1.8mmol, prewashed with hexane) in DMF (4ml) under nitrogen. The mixture was stirred for 15 minutes at ambient temperature and 4-chloro-6-methoxy-7-(2-(pyrrolidin-1-yl)ethoxy)quinazoline (184mg, 0.6mmol), (prepared as described for the starting material in Example 28), was added as a solid. The mixture was heated at 60°C for 45 minutes and then solvent was removed by evaporation. Water was added to the residue and the mixture was adjusted to pH2 with 2M hydrochloric acid. The insolubles were removed by filtration and the filtrate was extracted with ether. The aqueous layer was adjusted to pH8 with 2M aqueous sodium hydroxide solution and was concentrated by evaporation to half initial volume. The precipitate was collected by filtration, washed with water and ether, and dried under vacuum. The solid was dissolved in methylene chloride/methanol and 2.9M ethereal hydrogen chloride was added. The volatiles were removed by evaporation, the solid residue was suspended in ether, collected by filtration, washed with ether and dried under vacuum to give 4-(5-cyanooxindol-3-yl)-6-methoxy-7-(2-(pyrrolidin-1-yl)ethoxy)quinazoline hydrochloride (110mg, 40%).

¹H NMR Spectrum: (DMSO-d₆, D₂O + NaOD) 1.7(br s, 4H); 2.5-2.6(m, 4H); 2.86(t, 2H); 3.80(s, 3H); 4.16(t, 2H); 6.65(d, 1H); 6.81(d, 1H); 6.89(s, 1H); 8.1(br s, 1H); 8.40(s, 1H); 9.14(br s, 1H)

MS - ESI: 430 [MH]⁺

Elemental analysis:	Found	C 55.0 H 5.4 N 13.1
C ₂₄ H ₂₃ N ₃ O ₃ 1.8H ₂ O 1.7HCl	Requires	C 55.0 H 5.4 N 13.3%

5 **Example 39**

A solution of 5-cyanooxindole (2.96g, 18.7mmol), (Tet. Lett., 1987, 28, 4027), in DMF (35ml) was added dropwise under nitrogen to sodium hydride (749mg, 18.7mmol, prewashed with hexane) in DMF (15ml). The mixture was stirred for 30 minutes at ambient temperature and 4-chloro-7-(2-(imidazol-1-yl)ethoxy)-6-methoxyquinazoline (1.98g, 18.7mmol), (prepared as described for the starting material in Example 22), was added as a solid. The mixture was heated at 55°C for 45 minutes, allowed to cool and poured into a mixture of methylene chloride (400ml) and saturated aqueous ammonium chloride solution (400ml). The resulting precipitate was collected by filtration, resuspended in water and stirred for 15 minutes at ambient temperature. The solid was collected by filtration, washed with water and then ether, and dried under vacuum to give a white solid. This solid was suspended in methylene chloride (50ml) and methanol (50ml) and 2.2M ethereal hydrogen chloride (6ml) was added. The mixture was stirred for 15 minutes at ambient temperature, the solid product was collected by filtration, washed with ether and dried under vacuum at 60°C to give 4-(5-cyanooxindol-3-yl)-7-(2-(imidazol-1-yl)ethoxy)-6-methoxyquinazoline hydrochloride (2g, 61%).

¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 3.9(s, 3H); 4.65(t, 2H); 4.76(t, 2H); 7.06(d, 1H); 7.3(s, 1H); 7.45(d, 1H); 7.74(s, 1H); 7.85(s, 1H); 8.01(br s, 1H); 8.08(br s, 1H); 8.58(s, 1H); 9.2(s, 1H)

MS - ESI: 427 [MH]⁺

25 Elemental analysis:	Found	C 52.8 H 4.0 N 15.9
C ₂₃ H ₁₈ N ₆ O ₃ 1.5H ₂ O 1.95HCl	Requires	C 52.7 H 4.4 N 16.0%

Example 40

A solution of 5-acetyloxindole (352mg, 2mmol), (EP 0155828 A2), in DMF (5ml) was added to a suspension of sodium hydride (80mg, 2mmol, prewashed with hexane) in DMF (1ml) and the mixture stirred for 30 minutes at ambient temperature. 4-Chloro-6-methoxy-7-(2-methoxyethoxy)quinazoline (180mg, 0.67mmol), (prepared as described for the starting material in Example 2), was added and the mixture was heated at 50°C for 1.5 hours. The mixture was partitioned between water and ether and the aqueous layer was adjusted to pH7 with 2M hydrochloric acid. The precipitate was collected by filtration, washed with water and dried under vacuum. The solid was dissolved in methylene chloride/methanol (30ml/30ml) and 7M ethanolic hydrogen chloride (1ml) was added. The mixture was left to stand for 15 minutes at ambient temperature, the resulting precipitate was collected by filtration, washed with ether and dried under vacuum to give **4-(5-acetyloxindol-3-yl)-6-methoxy-7-(2-methoxyethoxy)quinazoline hydrochloride** (195mg, 71%).

¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 2.52(s, 3H); 3.37(s, 3H); 3.80(t, 2H); 3.85(s, 3H); 4.35(t, 2H); 7.09(d, 1H); 7.33(s, 1H); 7.78(s, 1H); 7.88(d, 1H); 8.25(s, 1H); 8.85(s, 1H)

MS - ESI: 408 [MH]⁺

Elemental analysis:	Found	C 57.3	H 5.0	N 8.7
C ₂₂ H ₂₁ N ₃ O ₅ · 1.2H ₂ O · 0.95HCl	Requires	C 57.0	H 5.3	N 9.0%

Example 41

Using an analogous procedure to that described for Example 38, 5-cyanooxindole (284mg, 1.8mmol), (Tet. Lett., 1987, 28, 4027), was treated with sodium hydride (72mg, 1.8mmol) and 4-chloro-6-methoxy-7-(2-([N-methyl-N-(4-pyridyl)]amino)ethoxy)-3,4-dihydroquinazolin-4-one (207mg, 0.6mmol) to give **4-(5-cyanooxindol-3-yl)-6-methoxy-7-(2-([N-methyl-N-(4-pyridyl)]amino)ethoxy)quinazoline hydrochloride** (83mg, 28%).

¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 3.3(s, 3H); 3.8(s, 3H); 4.16(t, 2H); 4.48(t, 2H); 7.05(br s, 1H); 7.1(d, 1H); 7.27(s, 1H); 7.35(br s, 1H); 7.47(s, 1H); 7.90(s, 1H); 8.05(s, 1H); 8.2-8.35(br s, 2H); 8.69(s, 1H)

MS - ESI: 467 [MH]⁺

Elemental analysis:	Found	C 55.9	H 4.6	N 15.2
---------------------	-------	--------	-------	--------

C₂₆H₂₂N₆O₃ 1.6H₂O 1.7HCl

Requires

C 56.0 H 4.9 N 15.1%

The starting material was prepared as follows:

Diethyl azodicarboxylate (1.1ml, 7mmol) was added dropwise to a solution of 7-hydroxy-6-methoxy-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (1.7g, 5.55mmol),
 5 (prepared as described for the starting material in Example 22), 2-([N-methyl-N-(4-pyridyl)]amino)ethanol (1.06g, 7mmol), (EP 0359389 A1), and triphenylphosphine (1.8g, 7mmol) in methylene chloride (29ml). The mixture was stirred for 1 hour at ambient temperature, then purified directly by flash chromatography eluting with methylene
 10 chloride/methanol (90/10). The purified solid was triturated with ether and collected by filtration to give 6-methoxy-7-(2-([N-methyl-N-(4-pyridyl)]amino)ethoxy)-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (2.29g, 94%).

¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 1.12(s, 9H); 3.3(s, 3H); 3.84(s, 3H); 4.11(t, 2H); 4.42(t, 2H); 5.92(s, 2H); 7.0(br s, 1H); 7.19(s, 1H); 7.3(br s, 1H); 7.5(s, 1H); 8.27(d, 2H);

15 8.4(s, 1H)

MS - ESI: 411 [MH]⁺

Elemental analysis:

Found C 60.5 H 12.2 N 6.4

C₂₃H₂₈N₄O, 0.8H₂O

Requires C 60.7 H 12.3 N 6.5%

A solution of 6-methoxy-7-(2-([N-methyl-N-(4-pyridyl)]amino)ethoxy)-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (2.24g, 5mmol) in methanol saturated
 20 with ammonia (30ml) was stirred at ambient temperature for 17 hours. The solvent was removed by evaporation, the solid residue was resuspended in ether, collected by filtration, washed with ether and dried under vacuum to give 6-methoxy-7-(2-([N-methyl-N-(4-pyridyl)]amino)ethoxy)-3,4-dihydroquinazolin-4-one (1.45g, 95%).

25 ¹H NMR Spectrum: (DMSO-d₆) 3.04(s, 3H); 3.83(t, 2H); 3.85(s, 3H); 4.3(t, 2H); 6.7(s, 2H); 7.12(s, 1H); 7.4(s, 1H); 7.96(s, 1H); 8.1(d, 2H)

MS - ESI: 327 [MH]⁺

Elemental analysis:

Found C 59.0 H 5.9 N 16.5

C₁₇H₁₈N₄O, 1H₂O

Requires C 59.3 H 5.8 N 16.3%

30 A solution of 6-methoxy-7-(2-([N-methyl-N-(4-pyridyl)]amino)ethoxy)-3,4-dihydroquinazolin-4-one (1.5g, 4.6mmol) in thionyl chloride (18ml) and DMF (0.7ml) was

heated at reflux for 1 hour. The mixture was allowed to cool, toluene was added and the volatiles were removed by evaporation. The residue was dissolved in methylene chloride and aqueous sodium hydrogen carbonate solution was added. The organic layer was separated, washed with brine, dried (MgSO_4), and the volatiles removed by evaporation. The residue
5 was purified by flash chromatography eluting with methylene chloride/methanol (95/5 followed by 90/10) to give 4-chloro-6-methoxy-7-(2-([N-methyl-N-(4-pyridyl)]amino)ethoxy)-3,4-dihydroquinazolin-4-one (463mg, 30%).
 ^1H NMR Spectrum: (DMSO-d_6) 3.05(s, 3H); 3.89(t, 2H); 3.97(s, 3H); 4.4(t, 2H); 6.7(d, 2H); 7.39(s, 1H); 7.46(s, 1H); 8.11(d, 2H); 8.87(s, 1H)
10 MS - ESI: 345 $[\text{MH}]^+$

Example 42

5-Methylsulphonyloxindole (262mg, 1.24mmol), (prepared as described for the starting material in Example 33), was added to a suspension of sodium hydride (50mg,
15 1.24mmol) in DMF (5ml) and the mixture stirred for 30 minutes at ambient temperature. 4-Chloro-6-methoxy-7-(3-morpholinopropoxy)quinazoline (140mg, 0.41mmol), (prepared as described for the starting material in Example 5), was added and the mixture heated at 50°C for 1.5 hours. The mixture was allowed to cool and was then partitioned between water and ether and the aqueous layer was adjusted to pH8 with 2M hydrochloric acid. The organic
20 phase was separated and the aqueous phase was extracted with ethyl acetate. The organic extracts were combined and the volatiles removed by evaporation. The residue was dissolved in methylene chloride and the organic layer was washed with brine, dried (MgSO_4) and the solvent removed by evaporation. The residue was purified by column chromatography, eluting with methanol/acetonitrile/methylene chloride (10/30/60 followed by 15/25/60). The
25 purified solid was dissolved in methanol/methylene chloride and 7M ethanolic hydrogen chloride (1.5ml) was added. The mixture was stirred for 10 minutes at ambient temperature, the solvent was removed by evaporation. The residue was triturated with ether, collected by filtration, washed with ether and dried under vacuum to give 6-methoxy-4-(5-methylsulphonyloxindol-3-yl)-7-(3-morpholinopropoxy)quinazoline hydrochloride (95mg,
30 39%).

- 108 -

¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 2.3(t, 2H); 3.09(s, 3H); 3.18(t, 2H); 3.36(t, 2H); 3.54(d, 2H); 3.72(t, 2H); 3.94(s, 3H); 4.04(d, 2H); 4.33(t, 2H); 7.18(d, 1H); 7.34(s, 1H); 7.64(d, 1H); 7.79(s, 1H); 8.11(s, 1H); 8.68(s, 1H)

MS - ESI: 513 [MH]⁺

5	Elemental analysis:	Found	C 48.0	H 5.3	N 8.5
	C ₂₅ H ₂₈ N ₄ O ₆ S 2.2H ₂ O 1.85HCl	Requires	C 48.0	H 5.5	N 8.9%
	0.1methylene chloride				

Example 43

10 A solution of 5-acetyloxindole (272mg, 1.56mmol), (EP 0155828 A2), in DMF (1.5ml) was added dropwise to a suspension of sodium hydride (62mg, 1.56mmol, prewashed with hexane) in DMF (3ml) and the mixture stirred for 30 minutes at ambient temperature. 4-Chloro-6-methoxy-7-(3-morpholinopropoxy)quinazoline (175mg, 0.52mmol), (prepared as described for the starting material in Example 5), was added and the mixture was heated at

15 50°C for 1.5 hours. The mixture was poured into a mixture of ether (30ml), ethyl acetate (30ml) and water (50ml). The aqueous layer was separated and adjusted to pH7.8 with 2M hydrochloric acid and extracted with methylene chloride. The organic extracts were washed with water, brine, dried (MgSO₄) and the solvent removed by evaporation. The residue was purified by column chromatography eluting with methylene chloride/methanol (90/10). The

20 purified solid was dissolved in methylene chloride/methanol and 7M ethanolic hydrogen chloride (1.5ml) was added. The volatiles were removed by evaporation, the residue collected and dried under vacuum to give **4-(5-acetyloxindol-3-yl)-6-methoxy-7-(3-morpholinopropoxy)quinazoline hydrochloride** (135mg, 48%).

¹H NMR Spectrum: (DMSO-d₆, CD₃CO₂D) 2.33(m, 2H); 2.54(s, 3H); 3.15(t, 2H); 3.37(t, 2H); 3.54(d, 2H); 3.73(t, 2H); 3.85(s, 3H); 4.03(d, 2H); 4.33(t, 2H); 7.09(d, 1H); 7.35(s, 1H); 7.8(s, 1H); 7.87(d, 1H); 8.25(s, 1H); 8.8(s, 1H)

MS - ESI: 477 [MH]⁺

	Elemental analysis:	Found	C 56.2	H 5.9	N 10.1
	C ₂₆ H ₂₈ N ₄ O ₅ 0.6H ₂ O 1.8HCl	Requires	C 56.6	H 5.7	N 10.0%
30	0.08ethanol				

Example 44

Using a procedure analogous to that described in Example 26, 5-nitrooxindole (250mg, 1.4mmol), (prepared as described for the starting material in Example 21), was added to sodium hydride (110mg, 2.75mmol) in THF (15ml) and the solution was treated with 4-chloro-7-(3-morpholinopropoxy)quinazoline (400mg, 1.3mmol), (prepared as described for the starting material in Example 37), in DMF (5ml) to give, after work-up and purification, 7-(3-morpholinopropoxy)-4-(5-nitrooxindol-3-yl)quinazoline (110mg, 19%).

¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 2.2-2.3(m, 2H); 3.1(t, 2H); 3.3(t, 2H); 3.5(d, 2H); 3.7(t, 2H); 4.0(m, 2H); 4.25(t, 2H); 7.0(d, 1H); 7.1(d, 1H); 7.18(dd, 1H); 7.95(dd, 1H); 8.55(s, 1H), 8.7(m, 2H), 11.2(br s, 1H)

MS - ESI: 450 [MH]⁺

Elemental analysis:	Found	C 58.4	H 5.1	N 14.9
---------------------	-------	--------	-------	--------

C ₂₃ H ₂₃ N ₃ O ₅ 1.3H ₂ O	Requires	C 58.4	H 5.5	N 14.8%
---	----------	--------	-------	---------

15 **Example 45**

Using a procedure analogous to that described in Example 26, 5-cyanooxindole (250mg, 1.5mmol), (Tet. Lett., 1987, 28, 4027), was added to sodium hydride (110mg, 2.75mmol) in THF (15ml) and the solution was treated with 4-chloro-7-(3-morpholinopropoxy)quinazoline (400mg, 1.3mmol), (prepared as described for the starting material in Example 37), in DMF (5ml) to give, after work-up and purification, 4-(5-cyanooxindol-3-yl)-7-(3-morpholinopropoxy)quinazoline (110mg, 20%).

¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 2.2-2.3(m, 2H); 3.1(t, 2H); 3.3(t, 2H); 3.5(d, 2H); 3.65(t, 2H); 3.95(d, 2H); 4.2(t, 2H); 6.95(d, 1H); 7.1(d, 1H); 7.2(dd, 1H); 7.4(dd, 1H); 8.1(br s, 1H); 8.5(s, 1H), 8.6(m, 1H), 11.0(br s, 1H)

25 MS - ESI: 428 [M-H]⁺

Elemental analysis:	Found	C 65.2	H 5.5	N 15.8
---------------------	-------	--------	-------	--------

C ₂₄ H ₂₃ N ₃ O ₃ 0.6H ₂ O	Requires	C 65.5	H 5.5	N 15.9%
---	----------	--------	-------	---------

Example 46

- A solution of 5-methylsulphonyloxindole (303mg, 1.55mmol) in DMF (3.5ml) was added to a suspension of sodium hydride (62mg, 1.55mmol, prewashed with hexane) in DMF (1.5ml) and the mixture stirred for 30 minutes at ambient temperature. 4-Chloro-6-methoxy-7-(3-morpholinopropoxy)quinazoline (175mg, 0.51mmol), (prepared as described for the starting material in Example 5), was added and the mixture was stirred for 1.5 hours at 50°C and then partitioned between water and ether. The aqueous layer was adjusted to pH7-8 with 2M hydrochloric acid and the volatiles removed by evaporation. The residue was purified by column chromatography eluting with methylene chloride/methanol (9/1). The semi-purified solid was dissolved in dilute hydrochloric acid (pH2) and purified by reverse phase C18 chromatography eluting with methanol/dilute hydrochloric acid (pH2) (gradient from 0/100 to 30/70). The methanol was removed by evaporation, the aqueous residue was freeze dried to give 6-methoxy-4-(5-methylsulphonyloxindol-3-yl)-7-(3-morpholinopropoxy)quinazoline hydrochloride (80mg, 27%).
- ¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 2.3(m, 2H); 2.67(s, 3H); 3.16(t, 2H); 3.35(t, 2H); 3.54(d, 2H); 3.73(t, 2H); 3.9(s, 3H); 4.02(d, 2H); 4.33(t, 2H); 7.13(d, 1H); 7.33(t, 1H); 7.36(s, 1H); 7.78(s, 1H); 8.0(s, 1H); 8.77(s, 1H)
- MS - ESI: 497 [MH]⁺
- | | | | | |
|--|----------|--------|-------|--------|
| Elemental analysis: | Found | C 51.4 | H 5.9 | N 9.7 |
| C ₂₅ H ₂₈ N ₄ O ₅ S 0.8H ₂ O 2HCl | Requires | C 51.4 | H 5.5 | N 9.6% |

The starting material was prepared as follows:

- 3-Chloroperoxybenzoic acid (400mg, 1.7mmol) was added in portions to a solution of 5-methylthiooxindole (300mg, 1.7mmol), (prepared as described for the starting material in Example 31), cooled at 5°C and the mixture stirred for 1 hour at 5°C and then for 1 hour at ambient temperature. The solvent was removed by evaporation and the residue was triturated with ether. The precipitate was collected by filtration, washed with ether and dried under vacuum. The solid was purified by flash chromatography eluting with methylene chloride/acetonitrile/methanol (60/32/8). The purified product was triturated with ether, the solid was collected by filtration and dried under vacuum to give 5-methylsulphonyloxindole (355mg, 81%).

¹H NMR Spectrum: (DMSO-d₆) 2.7(s, 3H); 3.58(s, 2H); 6.98(d, 1H); 7.51(d, 1H); 7.55(s, 1H); 10.67(br s, 1H)

MS - EI: 195 [M]⁺

5 Example 47

- A solution of 5-methoxycarbonyloxindole (169mg, 0.88mmol), (Eur. J. Med. Chem. 1983, 18, 107-111), in DMF (2ml) was added dropwise under nitrogen to sodium hydride (35mg, 0.88mmol, prewashed with hexane) in DMF (1.5ml). The mixture was stirred for 20 minutes at ambient temperature and 4-chloro-6-methoxy-7-(3-morpholinopropoxy)quinazoline (100mg, 0.3mmol), (prepared as described for the starting material in Example 5), was added as a solid. The mixture was heated at 60°C for 45 minutes, allowed to cool and partitioned between ethyl acetate and water. The aqueous layer was adjusted to pH8. The organic layer was separated, washed with brine, dried (MgSO₄) and the solvent removed by evaporation. The residue was purified by column chromatography eluting with methylene chloride/methanol (92/8). The solid was dissolved in methylene chloride/methanol and 3M ethereal hydrogen chloride (1ml) was added. The volatiles were removed by evaporation, the solid residue suspended in ether, collected by filtration, washed with ether and dried under vacuum to give 4-(5-methoxycarbonyloxindol-3-yl)-6-methoxy-7-(3-morpholinopropoxy)quinazoline hydrochloride (55mg, 32%).
- ¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 2.25-2.4(m, 2H); 3.15(t, 2H); 3.35(t, 2H); 3.58(d, 2H); 3.75(t, 2H); 3.82(s, 3H); 3.90(s, 3H); 4.05(d, 2H); 4.35(t, 2H); 7.1(d, 1H); 7.35(s, 1H); 7.8(d, 1H); 7.82(s, 1H); 8.3(s, 1H); 8.8(s, 1H)

MS - ESI: 493 [MH]⁺

Elemental analysis:	Found	C 50.9 H 6.2 N 9.1
25 C ₂₆ H ₂₈ N ₄ O ₆ 3H ₂ O 1.85HCl	Requires	C 50.9 H 5.9 N 9.1%

Example 48

- A solution of 5-(2-morpholinoethylaminosulphonyl)oxindole (363mg, 1.1mmol) in DMF (3ml) was added dropwise to a suspension of sodium hydride (89mg, 2.2mmol, prewashed with hexane) and the mixture stirred for 30 minutes at ambient temperature. 4-

Chloro-6-methoxy-7-(2-methoxyethoxy)quinazoline (100mg, 0.37mmol), (prepared as described for the starting material in Example 2), was added and the mixture heated at 50°C for 30 minutes. DMF (5ml) and DMSO (1.5ml) were added and the mixture heated at 60°C for 45 minutes, the volatiles were removed by evaporation and the residue was partitioned
 5 between methylene chloride and saturated aqueous ammonium chloride solution. The organic layer was separated, washed with brine, dried (MgSO₄) and the solvent removed by evaporation. The residue was purified by column chromatography eluting with methylene chloride/acetonitrile/methanol (60/35/5 followed by 60/30/10). The purified product was triturated with ether, collected by filtration and dried under vacuum. The solid was dissolved
 10 in methylene chloride/methanol and 5M ethereal hydrogen chloride (1.5ml) was added. The mixture was stirred for 30 minutes at ambient temperature, the volatiles were removed by evaporation and the solid residue was triturated with ether, collected by filtration and dried under vacuum to give **6-methoxy-7-(2-methoxyethoxy)-4-(5-(2-morpholinoethylaminosulphonyl)oxindol-3-yl)quinazoline hydrochloride** (110mg, 46%).
 15 ¹H NMR Spectrum: (DMSO-d₆, CD₃CO₂D) 2.97(t, 2H); 3.06(t, 2H); 3.15(t, 2H); 3.36(s, 3H); 3.36(m, 2H); 3.67(t, 2H); 3.78(t, 2H); 3.9(m, 2H); 3.92(s, 3H); 4.35(t, 2H); 7.15(d, 1H); 7.33(s, 1H); 7.57(d, 1H); 7.76(s, 1H); 8.07(s, 1H); 8.69(s, 1H)
 MS - ESI: 558 [MH]⁺
 Elemental analysis: Found C 47.7 H 5.4 N 11.0
 20 C₂₆H₃₁N₃O₇S 1.1H₂O 2HCl Requires C 48.0 H 5.5 N 10.8%

The starting material was prepared as follows:

A solution of 5-chlorosulphonyloxindole (1g, 4.3mmol), (prepared as described for the starting material in Example 60), and 4-(2-aminoethyl)morpholine (620μl, 4.7mmol) in
 25 ethanol (20ml) was heated at 65°C for 1 hour. The solvent was removed by evaporation, the residue was partitioned between methylene chloride and water and the aqueous layer was adjusted to pH7-8 with 5% aqueous sodium hydrogen carbonate solution. The organic layer was separated, washed with brine, dried (MgSO₄) and the solvent removed by evaporation. The residue was purified by column chromatography eluting with methylene
 30 chloride/methanol (95/5) to give 5-(2-morpholinoethylaminosulphonyl)oxindole (750mg, 54%).

- 113 -

¹H NMR Spectrum: (CDCl₃) 2.31(t, 4H); 2.44(t, 2H); 3.01(t, 2H); 3.60(s, 2H); 3.64(t, 4H); 5.16(br s, 1H); 6.97(d, 1H); 7.74(s, 1H); 7.78(d, 1H); 8.1(br s, 1H)

Example 49

5 6-Trifluoromethyloxindole (270mg, 1.3mmol) was dissolved in dry, degassed DMF (5ml) under argon. Sodium hydride (63mg, 1.6mmol) was added and the suspension stirred for 30 minutes at ambient temperature. A solution of 4-chloro-6-methoxy-7-(2-methoxyethoxy)quinazoline (111mg, 0.332mmol), (prepared as described for the starting material in Example 2), in dry, degassed DMF (5ml) was added dropwise and the mixture
10 heated at 100°C for 2 hours. The mixture was allowed to cool and the DMF was removed by evaporation. The residue was purified by flash column chromatography eluting with methylene chloride/methanol (100/0 and then 15/1). The purified orange solid was triturated with ether, collected by filtration and dried under vacuum at 40°C to give 6-methoxy-7-(2-methoxyethoxy)-4-(6-trifluoromethyloxindol-3-yl)quinazoline (100mg, 52%).

15 ¹H NMR Spectrum: (CDCl₃) 3.32(s, 3H); 3.74(t, 2H); 3.84(s, 3H); 4.32(t, 2H); 7.2(s, 1H); 7.3(m, 2H); 7.68(s, 1H); 7.78(d, 1H); 8.82(s, 1H); 11.25(s, 1H)

MS - ESI: 434 [MH]⁺

Elemental analysis:	Found	C 56.2	H 4.0	N 9.2
C ₂₁ H ₁₈ N ₃ O ₄ F ₃ · 0.7H ₂ O	Requires	C 56.6	H 4.4	N 9.4%

20

The starting material was prepared as follows:

Dibenzyl malonate (55.6ml, 222mmol) was added dropwise to a suspension of sodium hydride (8.9g of a 60% suspension in mineral oil, 222mmol) in DMF under argon and the reaction mixture stirred at ambient temperature for 30 minutes. 2-Bromo-5-
25 trifluoromethyl-1-nitrobenzene (30g, 111mmol) was then added and the solution stirred for 18 hours. The reaction mixture was poured onto ice and extracted with ether (1200ml). The ether layer was washed with water, brine, dried (MgSO₄), and the solvent removed by evaporation. The resulting red oil was purified by flash chromatography eluting with methylene chloride/isohexane (1/1) to give dibenzyl (2-nitro-4-
30 trifluoromethyl)phenylmalonate as a yellow oil (30g, 57%) which crystallised on standing.

- 114 -

Lithium chloride (5.3g, 130mmol) and water (1.14g, 40mmol) were added to a solution of dibenzyl (2-nitro-4-trifluoromethyl)phenylmalonate (30g, 60mmol) in DMSO (500ml) under argon and the resulting solution was heated to 100°C for 4 hours. The reaction mixture was allowed to cool to ambient temperature then diluted with ethyl acetate (1000ml).

- 5 The organic phase was separated, washed with water, brine, dried (MgSO₄), and the solvent removed by evaporation. The resulting red oil was purified by flash chromatography eluting with ethyl acetate/isohexane (1/5) to give benzyl (2-nitro-4-trifluoromethyl)phenylacetate as a yellow solid (16.6g, 77%).

- 10 Tin(II)chloride dihydrate (50.8g, 230mmol) was added to a solution of benzyl (2-nitro-4-trifluoromethyl)phenylacetate (15.3g, 45mmol) in ethyl acetate (250ml), and the mixture heated at reflux for 2 hours under argon. Concentrated aqueous ammonia was added to give an aqueous phase of pH8. The mixture was stirred for 30 minutes and the insoluble material was removed by filtration through diatomaceous earth. The filtrate was purified by column chromatography eluting with isohexanes/ethyl acetate (3/2) to give 6-
- 15 trifluoromethyloxindole as a white solid (7.2g, 80%).

¹H NMR Spectrum: (CDCl₃) 3.6(s, 2H); 7.15(s, 1H); 7.3(m, 2H); 9.0(br s, 1H)

MS: 200 [M-H]⁺

Elemental analysis: Found C 53.5 H 3.0 N 7.1

C₉H₆NO F₃ Requires C 53.7 H 3.0 N 7.0%

20

Example 50

- A solution of 6-fluorooxindole (407mg, 2.7mmol), (Synthesis 1993, 51), in DMF (6ml) was added dropwise to sodium hydride (108mg, 2.7mmol, prewashed with hexane) in DMF (6ml). The mixture was stirred for 30 minutes at ambient temperature and
- 25 4-chloro-7-(2-(imidazol-1-yl)ethoxy)-6-methoxyquinazoline (274mg, 0.9mmol), (prepared as described for the starting material in Example 22), was added as a solid. The mixture was heated at 65°C for 30 minutes, allowed to cool and was partitioned between saturated aqueous ammonium chloride solution and methylene chloride. The organic layer was separated, dried (MgSO₄), and the solvent removed by evaporation. The residue was triturated with ether,
- 30 collected by filtration, washed with ether and dried under vacuum. The solid was dissolved in methylene chloride/methanol and 5M ethereal hydrogen chloride (4ml) was added. The

mixture was concentrated by evaporation to 5ml total volume and the resulting solid was collected by filtration, washed with ether and dried under vacuum to give **4-(6-fluorooxindol-3-yl)-7-(2-(imidazol-1-yl)ethoxy)-6-methoxyquinazoline hydrochloride** (175mg, 40%).

¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 3.86(s, 3H); 4.62(t, 2H); 4.76(t, 2H); 6.8-6.9(m, 2H); 7.34(s, 1H); 7.65-7.70(m, 1H); 7.73(s, 2H); 7.84(s, 1H); 8.79(s, 1H); 9.22(s, 1H)

MS - ESI: 420 [MH]⁺

Elemental analysis:	Found	C 52.2	H 4.2	N 14.0
---------------------	-------	--------	-------	--------

C ₂₂ H ₁₈ FN ₅ O ₃ 0.7H ₂ O 1.95HCl	Requires	C 52.5	H 4.3	N 13.9%
--	----------	--------	-------	---------

10 Example 51

Using an analogous procedure to that described in Example 50, 5-bromooxindole (318mg, 1.5mmol), (prepared according to J. Am. Chem. Soc. 1975, 67, 1656), was treated with sodium hydride (60mg, 1.5mmol) in DMF (4ml) and 4-chloro-6-methoxy-7-(3-morpholinopropoxy)quinazoline (169mg, 0.5mmol), (prepared as described for the starting material in Example 5), to give **4-(5-bromooxindol-3-yl)-6-methoxy-7-(3-morpholinopropoxy)quinazoline hydrochloride** (108mg, 35%).

¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 2.2-2.3(m, 2H); 3.15(t, 2H); 3.35(t, 2H); 3.54(d, 2H); 3.73(t, 2H); 3.93(s, 3H); 4.04(d, 2H); 4.33(t, 2H); 6.95(d, 1H); 7.25(d, 1H); 7.33(s, 1H); 7.76(s, 1H); 7.8(s, 1H); 8.73(s, 1H)

20 MS - ESI: 515 [MH]⁺

Elemental analysis:	Found	C 47.2	H 4.6	N 8.9
---------------------	-------	--------	-------	-------

C ₂₄ H ₂₅ N ₄ O ₄ Br 1.8H ₂ O 1.8HCl	Requires	C 47.1	H 5.0	N 9.2%
---	----------	--------	-------	--------

Example 52

25 Using a procedure analogous to that described in Example 26, 6-trifluoromethyloxindole (250mg, 1.25mmol), (prepared as described for the starting material in Example 49), was added to sodium hydride (100mg, 2.5mmol) in DMF (15ml) and the solution was reacted with 4-chloro-7-(3-morpholinopropoxy)quinazoline (300mg, 1.0mmol),

(prepared as described for the starting material in Example 37), in DMF (5ml) to give, after work-up and purification, 7-(3-morpholinopropoxy)-4-(6-trifluoromethyloxindol-3-yl)quinazoline (80mg, 16%).

¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 2.2(m, 2H); 3.1(m, 2H); 3.3(t, 2H); 3.5(d, 2H);
5 3.7(t, 2H); 4.0(d, 2H); 4.25(t, 2H); 7.1-7.2(m, 4H); 7.75(d, 1H); 8.4(d, 1H); 8.85(s, 1H),
11.2(br s, 1H)

MS - ESI: 473 [MH]⁺

Elemental analysis:	Found	C 59.3	H 4.7	N 11.7
---------------------	-------	--------	-------	--------

C ₂₄ H ₂₃ F ₃ N ₄ O ₃ · 0.6H ₂ O	Requires	C 59.6	H 5.1	N 11.6%
--	----------	--------	-------	---------

10

Example 53

Sodium hydride (53mg, 1.33mmol) was added to a solution of 6-trifluoromethyloxindole (200mg, 0.995mmol), (prepared as described for the starting material in Example 49), in dry, degassed DMF and the suspension stirred for 15 minutes at ambient
15 temperature. A solution of 4-chloro-6-methoxy-7-(3-morpholinopropoxy)quinazoline (111mg, 0.332mmol), (prepared as described for the starting material in Example 5), in a dry, degassed mixture of DMF (5ml) and THF (2ml) was added dropwise and the reaction mixture was heated at 60°C for 3 hours. The mixture was allowed to cool, the DMF was removed by evaporation, 1M hydrochloric acid was added until the aqueous phase was pH8 and ethyl
20 acetate was added to dissolve the resulting precipitate. The organic phase was separated and the aqueous phase re-extracted with ethyl acetate (50ml x3). The ethyl acetate fractions were combined, dried (MgSO₄) and the solvent removed by evaporation. The residue was purified by flash column chromatography eluting with methylene chloride/methanol (10/1). The purified product was triturated with boiling ether and collected by filtration to give 6-
25 methoxy-7-(3-morpholinopropoxy)-4-(6-trifluoromethyloxindol-3-yl)quinazoline (51mg, 31%).

¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 2.2(m, 2H), 3.1(m, 2H), 3.3(t, 2H), 3.5(m, 2H),
3.65(t, 2H), 3.82(s, 3H), 3.9(m, 2H), 4.28(t, 2H), 7.22(s, 1H), 7.3(m, 2H), 7.68(s, 1H), 7.74(d,
1H), 8.86(s, 1H), 11.54(s, 1H)

30 MS: 503 [MH]⁺

- 117 -

Elemental analysis:	Found	C 57.0	H 4.7	N 10.5
$C_{25}H_{23}N_4O_4F_3 \cdot 1.2H_2O$	Requires	C 57.1	H 5.3	N 10.7%

Example 54

- 5 A solution of 5-carbamoyloxindole (391mg, 2.2mmol) in DMF (5ml) was added dropwise to sodium hydride (89mg, 2.2mmol, prewashed with hexane) in DMF (3.5ml). The mixture was stirred for 20 minutes at ambient temperature and 4-chloro-6-methoxy-7-(3-morpholinopropoxy)quinazoline (250mg, 0.74mmol), (prepared as described for the starting material in Example 5), was added as a solid and the mixture was heated at 60°C for 1 hour.
- 10 The volatiles were removed by evaporation and the residue was partitioned between ethyl acetate and water. The aqueous layer was adjusted to pH8.5 with 2M hydrochloric acid. The precipitate was collected by filtration, washed with water and dried under vacuum. The solid was dissolved in methylene chloride/methanol and 3M ethereal hydrogen chloride (2ml) was added. The mixture was concentrated by evaporation and the resulting precipitate was
- 15 collected by filtration, washed with ether and dried under vacuum to give 4-(5-carbamoyloxindol-3-yl)-6-methoxy-7-(3-morpholinopropoxy)quinazoline hydrochloride (184mg, 45%).

¹H NMR Spectrum: (DMSO_d₆, CF₃CO₂D) 2.3-2.4(m, 2H); 3.16(t, 2H); 3.35(t, 2H); 3.54(d, 2H); 3.75(t, 2H); 3.92(s, 3H); 4.04(d, 2H); 4.32(t, 2H); 7.03(d, 1H); 7.36(s, 1H); 7.77(d, 1H);

20 7.8(s, 1H); 8.31(s, 1H); 8.88(s, 1H)

MS - ESI: 478 [MH]⁺

Elemental analysis:	Found	C 52.1	H 5.7	N 12.0
$C_{25}H_{27}N_5O_5 \cdot 2H_2O \cdot 1.7HCl$	Requires	C 52.2	H 5.7	N 12.2%

- 25 The starting material was prepared as follows:

A solution of 5-cyanooxindole (1.2g, 7.6mmol), (Tet. Let., 1987, 28, 4027), in 85% sulphuric acid (8ml) was heated at 80°C for 6 hours. The mixture was allowed to cool, poured into ice/water and the mixture was adjusted to pH5 with 2M sodium hydroxide

- 118 -

solution. The solid product was collected by filtration, washed with water, and dried under vacuum at 60°C over phosphorus pentoxide to give 5-carbamoyloxindole (1.02g, 76%).

¹H NMR Spectrum: (DMSO-d₆) 3.52(s, 2H); 6.83(d, 1H); 7.14(br s, 1H); 7.74(s, 1H); 7.74(d, 1H); 7.76(br s, 1H); 10.6(s, 1H)

5 MS - EI: 176 [M]⁺

Example 55

A solution of 5-(2-morpholinoethylaminosulphonyl)oxindole (364mg, 1.1mmol), (prepared as described for the starting material in Example 48), in DMF (3.5ml) was added dropwise to a suspension of sodium hydride (90mg, 2.2mmol, prewashed with hexane) in DMF (1.5ml). The mixture was stirred for 30 minutes at ambient temperature and 4-chloro-6,7-dimethoxyquinazoline (84mg, 0.37mmol), (prepared as described for the starting material in Example 1), was added as a solid followed by DMSO (2ml). The mixture was stirred for 45 minutes at 60°C, allowed to cool and partitioned between methylene chloride and saturated aqueous ammonium chloride solution. The organic layer was separated, washed with brine, dried (MgSO₄) and the solvent removed by evaporation. The residue was purified by column chromatography eluting with methylene chloride/methanol (95/5 followed by 92/8). The purified solid was triturated with ether, collected by filtration, washed with ether and dried under vacuum. The solid was dissolved in methylene chloride/methanol and 2M ethereal hydrogen chloride (1ml) was added. The volatiles were removed by evaporation, the solid residue triturated with ether, collected by filtration, washed with ether and dried under vacuum to give 6,7-dimethoxy-4-(5-(2-morpholinoethylaminosulphonyl)oxindol-3-yl)quinazoline hydrochloride (106mg, 50%).

¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 2.99(t, 2H); 3.1(t, 2H); 3.15(t, 2H); 3.37(d, 2H); 3.67(t, 2H); 3.92(s, 3H); 3.9(br s, 2H); 4.01(s, 3H); 7.15(d, 1H); 7.32(s, 1H); 7.56(d, 1H); 7.77(s, 1H); 8.08(s, 1H); 8.7(s, 1H)

MS - ESI: 514 [MH]⁺

Elemental analysis:	Found	C 48.7	H 5.1	N 11.6
C ₂₄ H ₂₇ N ₃ O ₆ S 1.2H ₂ O 1.5HCl	Requires	C 48.9	H 5.3	N 11.9%

Example 56

Using a procedure analogous to that described in Example 26, 6-cyanooxindole (250mg, 1.6mmol), (prepared as described for the starting material in Example 57), was added to sodium hydride (100mg, 2.5mmol) in DMF (15ml) and the solution was treated with 4-chloro-7-(3-morpholinopropoxy)quinazoline (300mg, 1.0mmol), (prepared as described for the starting material in Example 37), in DMF (5ml) to give, after work-up and purification, 4-(6-cyanooxindol-3-yl)-7-(3-morpholinopropoxy)quinazoline (90mg, 20%).

¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 2.2(m, 2H); 3.1(t, 2H); 3.3(t, 2H); 3.5(d, 2H); 3.6(t, 2H); 4.0(m, 2H); 4.25(t, 2H); 7.1-7.3(m, 4H); 7.75(d, 1H); 8.5(d, 1H); 8.55(s, 1H), 11.0(s, 1H)

MS - ESI: 429 [M-H]⁺

Elemental analysis:	Found	C 62.9	H 5.4	N 15.6
C ₂₄ H ₂₃ N ₃ O ₃ · 1.6H ₂ O	Requires	C 62.9	H 5.7	N 15.3%

Example 57

Sodium hydride (100mg, 2.6mmol) was added to a solution of 6-cyanooxindole (350mg, 2.2mmol) in dry degassed DMF (10ml) under argon and the suspension stirred for 30 minutes at ambient temperature. A solution of 4-chloro-6-methoxy-7-(2-methoxyethoxy)quinazoline (200mg, 0.74mmol), (prepared as described for the starting material in Example 2), in dry degassed DMF (2ml) was added dropwise and the reaction mixture was heated at 60°C for 2 hours. The mixture was allowed to cool and the DMF was removed by evaporation and the residue was partitioned between ethyl acetate (50ml) and water (50ml). The organic phase was separated and the aqueous phase was extracted with ethyl acetate (50ml x3). The organic extracts were combined, dried (MgSO₄) and the solvent removed by evaporation. The solid residue was purified by flash column chromatography eluting with methylene chloride/methanol/ammonia (99/1/1 and then 95/5/1). The purified solid was triturated in hot ether and collected by filtration and then triturated in hot methylene chloride/methanol (1/1), collected by filtration and dried to give 4-(6-cyanooxindol-3-yl)-6-methoxy-7-(2-methoxyethoxy)quinazoline (84mg, 29%).

- 120 -

¹H NMR Spectrum: (CDCl₃) 3.34(s, 3H); 3.74(t, 2H); 3.84(s, 3H); 4.52(t, 2H); 7.3(m, 3H); 7.7(m, 2H); 8.70(s, 1H); 11.32(s, 1H)

MS: 391 [MH]⁺

Elemental analysis: Found C 62.8 H 4.5 N 14.3

5 C₂₁H₁₈N₄O₄ 0.3H₂O Requires C 62.8 H 4.4 N 14.1%

The starting material was prepared as follows:

Diethyl malonate (44.6ml, 293mmol) was added dropwise over 1.5 hours to a suspension of sodium hydride (11.7g of a 60% suspension in mineral oil, 293mmol) in DMF
10 under argon and the reaction mixture was stirred at ambient temperature for 30 minutes. The resulting solution was added dropwise over 1 hour to a solution of 2-chloro-5-cyano-1-nitrobenzene (24.3g, 133mmol) in DMF (75ml) cooled with a dry ice/acetone bath. The temperature of the cooling bath was adjusted to prevent the freezing of the red reaction mixture. After the addition was completed, the reaction was left to rise to ambient
15 temperature over 2 hours. The resulting red solution was poured slowly into a stirred mixture of ice (100ml) and 1M hydrochloric acid (10ml). The resulting yellow precipitate was collected by filtration, washed with water until the filtrate was at pH7.0 and then dried under vacuum to give diethyl (4-cyano-2-nitrophenyl)malonate (39.6g, 97%).

Lithium chloride (10.7g, 255mmol) and water (2.3g, 127mmol) were added to a
20 solution of diethyl (4-cyano-2-nitrophenyl)malonate (39g, 127mmol) in DMSO (700ml) under argon and the solution was heated at 100°C for 4 hours. The reaction mixture was allowed to cool and then poured slowly into stirred ice water (1000ml). The resulting pale yellow precipitate was collected by filtration, washed with water (100ml x4) and dried under vacuum to give ethyl (4-cyano-2-nitrophenyl)acetate (27.1g, 91%).

25 Iron powder (11.5g, 205.2mmol) was added to a saturated solution of ethyl (4-cyano-2-nitrophenyl)acetate (12g, 51.3mmol) in hot glacial acetic acid and the reaction mixture was stirred at 100°C for 2 hours. The mixture was allowed to cool and poured into stirred ice water (1000ml). The aqueous mixture was extracted with ethyl acetate, the extracts were combined, dried (MgSO₄) and the solvent removed by evaporation. The solid residue
30 was recrystallised from ethyl acetate to give 6-cyanooxindole (6.94g, 85%).

¹H NMR Spectrum: (CDCl₃) 3.58(s, 2H), 7.1(s, 1H), 7.38(s, 2H)

MS: 157 [M-H]⁺

Example 58

A solution of 5-(2-morpholinoethylaminosulphonyl)oxindole (320mg, 0.98mmol),
 5 (prepared as described for the starting material in Example 48), in DMF (3.5ml) was added to
 a suspension of sodium hydride (39.4mg, 0.98mmol, prewashed with THF) in DMF (1.5ml).
 The mixture was stirred for 30 minutes at ambient temperature and 4-chloro-7-(2-(imidazol-1-
 yl)ethoxy)-6-methoxyquinazoline (100mg, 0.32mmol), (prepared as described for the starting
 material in Example 22), was added. The mixture was stirred for 1 hour at 60°C, allowed to
 10 cool and partitioned between methylene chloride and saturated aqueous ammonium chloride
 solution. The organic layer was separated, washed with brine, dried (MgSO₄) and the solvent
 removed by evaporation. The residue was purified by column chromatography eluting with
 methanol/methylene chloride (5/95 to 15/85). The purified product was triturated with ether,
 collected by filtration and dried under vacuum. The solid was dissolved in methylene
 15 chloride/methanol and 2.2M ethereal hydrogen chloride (1ml) was added. The volatiles were
 removed by evaporation, the solid residue was triturated with ether, collected by filtration and
 dried under vacuum to give 7-(2-(imidazol-1-yl)ethoxy)-6-methoxy-4-(5-(2-
morpholinoethylaminosulphonyl)oxindol-3-yl)quinazoline hydrochloride (153mg, 64%).
¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 2.94(t, 2H); 3.10(t, 2H); 3.15(t, 2H); 3.36(d, 2H);
 20 3.65(t, 2H); 3.91(s, 3H); 4.0(d, 2H); 4.65(t, 2H); 4.76(t, 2H); 7.12(d, 1H); 7.35(s, 1H); 7.55(d,
 1H); 7.74(s, 1H); 7.82(s, 1H); 7.84(s, 1H); 8.07(br s, 1H); 8.57(s, 1H); 9.21(s, 1H)
 MS - ESI: 594 [MH]⁺

Elemental analysis:	Found	C 46.0	H 4.9	N 13.3
C ₂₈ H ₃₁ N ₇ O ₆ S 1.6H ₂ O 2.85HCl	Requires	C 46.3	H 5.1	N 13.5%

25

Example 59

Sodium hydride (80mg, 2.07mmol) was added to a solution of 6-cyanooxindole
 (280mg, 1.78mmol), (prepared as described for the starting material in Example 57), in dry,
 degassed DMF (10ml) and the suspension stirred for 15 minutes at ambient temperature. A
 30 solution of 4-chloro-6-methoxy-7-(3-morpholinopropoxy)quinazoline (200mg, 0.59mmol),
 (prepared as described for the starting material in Example 5), in a dry, degassed mixture of

DMF (2ml) and THF (2ml) was added dropwise and the reaction mixture was heated at 60°C for 3 hours. The mixture was allowed to cool, the solvents were removed by evaporation and the residue was partitioned between ethyl acetate and water. The organic phase was separated and the aqueous phase extracted with ethyl acetate (50ml x3). The ethyl acetate fractions
 5 were combined, dried (MgSO₄), the solvent removed by evaporation and the residue was purified by flash column chromatography eluting with methylene chloride/methanol (100/13). The purified product was triturated in boiling ether and the solid product collected by filtration to give 4-(6-cyanooxindol-3-yl)-6-methoxy-7-(3-morpholinopropoxy)quinazoline (36mg, 13%) as an orange solid.

10 ¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 2.3(m, 2H); 3.1(m, 2H); 3.3(m, 2H); 3.5(m, 2H); 3.65(m, 2H); 3.85(s, 3H); 4.0(m, 2H); 4.3(m, 2H); 7.3(m, 3H); 7.7(m, 2H); 8.7(s, 1H); 11.18(s, 1H)

MS: 460 [MH]⁺

Elemental analysis:	Found	C 64.0	H 5.5	N 15.2
---------------------	-------	--------	-------	--------

15 C ₂₅ H ₂₃ N ₅ O ₄ 0.4H ₂ O	Requires	C 64.3	H 5.6	N 15.0%
--	----------	--------	-------	---------

Example 60

A solution of 5-aminosulphonyloxindole (355mg, 1.67mmol) in DMF (3.5ml) was added dropwise to sodium hydride (67mg, 1.67mmol, prewashed with THF) in DMF (5ml).

20 The mixture was stirred for 30 minutes at ambient temperature and a solution of 4-chloro-6-methoxy-7-(2-methoxyethoxy)quinazoline (150mg, 0.56mmol), (prepared as described for the starting material in Example 2), in DMF (3ml) was added. The mixture was heated at 65°C for 2.5 hours and then stirred overnight at ambient temperature. The mixture was partitioned between ethyl acetate and water, the aqueous layer was adjusted to pH7 with 1M hydrochloric
 25 acid and left standing for 2 hours. The resulting precipitate was collected by filtration, washed with water and ether, and dried under vacuum. The solid was dissolved in methylene chloride/methanol and 2.2M ethereal hydrogen chloride (1ml) was added. The volatiles were removed by evaporation, the solid residue was triturated with ether, collected by filtration and dried under vacuum to give 4-(5-aminosulphonyloxindol-3-yl)-6-methoxy-7-(2-
 30 methoxyethoxy)quinazoline hydrochloride (144mg, 60%).

- 123 -

¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 3.36(s, 3H); 3.77(t, 2H); 3.92(s, 3H); 4.35(t, 2H); 7.11(d, 1H); 7.33(s, 1H); 7.61(d, 1H); 7.71(s, 1H); 8.1(s, 1H); 8.76(s, 1H)

MS - ESI: 445 [MH]⁺

Elemental analysis:		Found	C 49.4	H 4.5	N 11.5
5	C ₂₀ H ₂₀ N ₄ O ₆ S 0.3H ₂ O 0.95HCl	Requires	C 49.6	H 4.5	N 11.6%

The starting material was prepared as follows:

Oxindole (2g, 15mmol) was added in portions to chlorosulphonic acid (5ml, 75mmol) cooled at 5°C. The mixture was stirred at ambient temperature for 10 minutes and then heated at 80°C for 30 minutes. The mixture was allowed to cool, and then poured onto ice. The precipitate was collected by filtration, washed with water, and dried over phosphorus pentoxide to give 5-chlorosulphonyloxindole (2.8g, 80%).

¹H NMR Spectrum: (DMSO-d₆) 3.46(s, 2H); 6.72(d, 1H); 7.43(m, 2H); 10.43(br s, 1H)

MS - EI: 232 [M]⁺

15 A solution of 5-chlorosulphonyloxindole (2g, 0.86mmol) in ethanol (30ml), methylene chloride (150ml) and 2.8M methanolic ammonia (6.2ml, 17mmol) was stirred overnight at ambient temperature. The volatiles were removed by evaporation and the solid residue was purified by column chromatography eluting with methanol/methylene chloride (15/85) to give 5-aminosulphonyloxindole (1.54g, 84%).

20 ¹H NMR Spectrum: (DMSO-d₆) 3.58(s, 2H); 6.92(d, 1H); 7.18(s, 2H); 7.6-7.7(m, 2H); 10.73(s, 1H)

Example 61

Using a procedure analogous to that described in Example 26, 6-phenyloxindole (300mg, 1.4mmol) was added to sodium hydride (90mg, 2.25mmol) in DMF (10ml) and the solution was treated with 4-chloro-7-(3-morpholinopropoxy)quinazoline (210mg, 0.73mmol), (prepared as described for the starting material in Example 37), in DMF (10ml) to give, after work-up and purification, 7-(3-morpholinopropoxy)-4-(6-phenyloxindol-3-yl)quinazoline (365mg, 28%).

- 124 -

¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 2.2(m, 2H); 3.1(m, 2H); 3.3(t, 2H); 3.5(d, 2H); 3.65(t, 2H); 4.0(m, 2H); 4.25(t, 2H); 7.1(m, 2H); 7.15(d, 1H); 7.2(dd, 1H); 7.55(d, 1H); 8.4(d, 1H); 8.6(s, 1H), 9.8(br s, 1H); 11.0(s, 1H)

MS - ESI: 483 [MH]⁺

5	Elemental analysis:	Found	C 55.2	H 4.5	N 11.3
	C ₂₃ H ₂₃ N ₄ O ₃ Br 1H ₂ O	Requires	C 55.5	H 5.0	N 11.3%

The starting material was prepared as follows:

6-Bromooxindole (800mg, 3.8mmol), (prepared as described for the starting material
10 in Example 62), phenylboronic acid (500mg, 4.1mmol) sodium hydrogen carbonate (saturated aqueous solution, 50ml) and 1,2-dimethoxyethane (100ml) were mixed together and the solution was degassed using alternatively vacuum and argon.

Tetrakis(triphenylphosphine)palladium(0) (50mg) was added and the solution degassed again before heating it to reflux for 8 hours. The reaction mixture was evaporated to dryness using
15 a rotary evaporator and the solid residue partitioned between water (75ml) and ethyl acetate (100ml). The aqueous phase was reextracted with more ethyl acetate (100ml x3), the organic phases were combined, washed with brine and passed directly through a short plug of silica to remove the baseline impurities. The solvent was evaporated and the solid was triturated in ether and filtered to give 6-phenyloxindole as a pink solid (375mg, 60%).

20 ¹H NMR Spectrum: (DMSO-d₆) 3.5(s, 2H); 7.0(s, 1H); 7.2(d, 1H); 7.25(d, 1H); 7.35(d, 1H); 7.45(t, 2H); 7.6(d, 2H); 10.42(br s, 1H)

MS - ESI: 208 [M-H]⁺

Example 62

25 Using a procedure analogous to that described in Example 26, 6-bromooxindole (1.16g, 5.5mmol) was added to sodium hydride (324mg, 8.6mmol) in DMF (20ml) and the solution was treated with 4-chloro-7-(3-morpholinopropoxy)quinazoline (790mg, 2.7mmol), (prepared as described for the starting material in Example 37), in DMF (10ml) to give, after work-up and purification, 4-(6-bromooxindol-3-yl)-7-(3-morpholinopropoxy)quinazoline
30 (365mg, 28%).

- 125 -

¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 2.2(m, 2H); 3.1(m, 2H); 3.3(t, 2H); 3.5(d, 2H); 3.65(t, 2H); 4.0(m, 2H); 4.25(t, 2H); 7.1(m, 2H); 7.15(d, 1H); 7.2(dd, 1H); 7.55(d, 1H); 8.4(d, 1H); 8.6(s, 1H), 9.8(br s, 1H); 11.0(s, 1H)

MS - ESI: 483 [MH]⁺

5	Elemental analysis:	Found	C 55.2	H 4.5	N 11.3
	C ₂₃ H ₂₃ N ₄ O ₃ Br 1H ₂ O	Requires	C 55.5	H 5.0	N 11.3%

The starting material was prepared as follows:

2,5-Dibromonitrobenzene (41.5g, 148mmol) was dissolved in anhydrous DMF (500ml) and sodium hydride (17.7g of a 60% suspension in mineral oil, 443mmol) was added under argon. Diethyl malonate (49.6g, 295mmol) was added dropwise over 10 minutes giving a strong exotherm (mixture rose to 60°C). The reaction mixture was maintained at 60°C for 6 hours then left to cool to ambient temperature and stirred for 18 hours. The resulting dark red solution was poured slowly into 2M hydrochloric acid (1500ml) and extracted with ether (500ml x4). The organic phases were combined, washed with brine, dried (MgSO₄) and the solvent removed by evaporation. The crude oil was purified by flash chromatography eluting with isohexanes/ethyl acetate (100/0, then 95/5 and finally 80/20) to give diethyl (4-bromo-2-nitrophenyl)malonate (51g, 96%).

Using a procedure analogous to that described for the starting material in Example 57, diethyl (4-bromo-2-nitrophenyl)malonate (35g, 97 mmol) was converted into ethyl (4-bromo-2-nitrophenyl)acetate (26g, 93%).

Using a procedure analogous to that described for the starting material in Example 57, ethyl (4-bromo-2-nitrophenyl)acetate (26g, 90mmol) was treated with iron in acetic acid. The product thus obtained was recrystallised from ether/isohexanes to give 6-bromooxindole (8.2g, 42%).

¹H NMR Spectrum: (DMSO-d₆) 3.45(s, 2H); 6.9(s, 1H); 7.1(m, 2H); 10.45(br s, 1H)

MS - ESI: 210 and 212 [M-H]⁺

Example 63

Using a procedure analogous to that described in Example 26, 6-(1-t-butoxycarbonylpyrrol-2-yl)oxindole (500mg, 1.7mmol) was added to sodium hydride (90mg, 2.25mmol) in THF (10ml) and the solution was treated with 4-chloro-6-methoxy-7-(3-morpholinopropoxy)quinazoline (250mg, 0.74mmol), (prepared as described for the starting material in Example 5), in DMF (2ml) to give, after work-up and purification, 4-(6-(1-t-butoxycarbonylpyrrol-2-yl)oxindol-3-yl)-6-methoxy-7-(3-morpholinopropoxy)quinazoline (185mg, 40%).

¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 1.3(s, 9H); 2.2(m, 2H); 3.1(t, 2H); 3.3(t, 2H); 3.5(d, 2H); 3.6(t, 2H); 3.8(s, 3H); 3.9(d, 2H); 4.25(t, 2H); 6.2(m, 2H); 6.95(s, 1H); 7.0(d, 1H); 7.25(m, 2H); 7.6(d, 1H); 7.7(s, 1H); 8.95(s, 1H); 9.7(br s, 1H); 11.2(s, 1H)

MS - ESI: 600 [MH]⁺

Elemental analysis:	Found	C 65.7	H 5.9	N 11.9
C ₃₃ H ₃₇ N ₅ O ₆ 0.4H ₂ O	Requires	C 65.3	H 6.3	N 11.5%

15

The starting material was prepared as follows:

Triethylamine (13ml, 93mmol), 4-(N,N-dimethylamino)pyridine (100mg), and di-t-butyl dicarbonate (18.1g, 83mmol) were added successively to a solution of pyrrole (5g, 74.6mmol) in acetonitrile (125ml). The reaction mixture was then heated at 60°C for 30 minutes. The reaction was allowed to cool and the solvent was removed by evaporation. The residue was purified by flash column chromatography eluting with isohexanes/ethyl acetate (100/0 and then 95/5) to give 1-t-butoxycarbonylpyrrole (8.74g, 70%).

n-Butyllithium (29ml of a 1.6M solution in hexane, 47.4mmol) was added dropwise over 10 minutes to a solution of 2,2,6,6-tetramethylpiperidine (6.7g, 47.4mmol) in THF (175ml) cooled to -78°C. A solution of 1-t-butoxycarbonylpyrrole (8.19g, 49mmol) in THF (25ml) was added and the mixture was maintained below -70°C. The reaction mixture was stirred for 1.5 hours at -78°C and tri-isopropyl borate (17g, 90mmol) was added. The reaction mixture was then allowed to warm to ambient temperature over 1 hour. The mixture was diluted with ether (500ml) and washed with 1M potassium hydrogen sulphate solution (100ml x3), 1M sodium hydrogen carbonate solution (100ml x2) and brine (100ml). The ether phase

30

- 127 -

was dried (MgSO_4) and the solvent removed by evaporation. The residue was purified by flash column chromatography eluting with isohexanes/ethyl acetate (a gradient from 100/0 to 50/50) to give the 1-t-butoxycarbonylpyrrol-2-ylboronic acid (4.05g, 43%).

1-t-Butoxycarbonylpyrrol-2-ylboronic acid (1.5g, 7.1mmol) and a saturated aqueous solution of sodium hydrogen carbonate (25ml) were added to a solution of 6-bromooxindole (1.1g, 5.2mmol), (prepared as described for the starting material in Example 62), in 1,2-dimethoxyethane (75ml). The reaction mixture was degassed and placed under argon and tetrakis(triphenylphosphine)palladium(0) (50mg) was added and the mixture heated at reflux for 6 hours. The organic solvent was removed by evaporation and the reaction mixture partitioned between ethyl acetate and water. The organic phase was separated and the aqueous phase was re-extracted twice with ethyl acetate. The organic phases were combined, washed with brine, dried (MgSO_4), and the solvent removed by evaporation. The residue was triturated with ether/isohexanes (1/1) and collected by filtration to give 6-(1-t-butoxycarbonylpyrrol-2-yl)oxindole (1.11g, 71%).

^1H NMR Spectrum: (DMSO-d_6) 1.25(s, 9H); 3.45(s, 2H); 6.2(m, 2H); 6.65(s, 1H); 6.85(d, 1H); 7.15(d, 1H); 7.25(s, 1H); 10.35(br s, 1H)

Example 64

4-(6-(1-t-Butoxycarbonylpyrrol-2-yl)oxindol-3-yl)-6-methoxy-7-(3-morpholinopropoxy)quinazoline (150mg, 0.25mmol), (prepared as described in Example 63), was dissolved in TFA (5ml) and triethylsilane (0.5ml) and the mixture stirred at ambient temperature for 2 hours. The volatiles were removed by evaporation and azeotroping with toluene. The residue suspended in methanol/methylene chloride and 1M ethereal hydrogen chloride (1.25ml) was added and the volatiles were removed by evaporation. The residue was triturated with t-butyl methyl ether and collected by filtration to give 4-(6-(pyrrol-2-yl)oxindol-3-yl)-6-methoxy-7-(3-morpholinopropoxy)quinazoline hydrochloride (132mg, 87%) as a dark purple solid.

^1H NMR Spectrum: (DMSO-d_6 , $\text{CF}_3\text{CO}_2\text{D}$) 2.3(m, 2H); 3.1(m, 2H); 3.25(m, 2H); 3.45(d, 2H); 3.8(m, 5H); 3.9(d, 2H); 4.25(t, 2H); 6.1(m, 1H); 6.4(m, 1H); 6.8(s, 1H); 7.2(s, 1H); 7.25(m, 2H); 7.6(d, 1H); 7.8(s, 1H); 8.55(s, 1H); 10.9(br s, 1H); 11.1(m, 1H); 11.3(br s, 1H)

MS - ESI: 500 [MH]⁺

Elemental analysis:	Found	C 55.7	H 5.9	N 11.4
C ₃₃ H ₃₇ N ₅ O ₆ · 1.7H ₂ O · 2HCl	Requires	C 55.8	H 5.8	N 11.6%

5 Example 65

A solution of oxindole (122mg, 0.91mmol) in DMF (1.5ml) was added to a suspension of sodium hydride (37mg, 0.91mmol, prewashed with hexane) in DMF (3.5ml). The mixture was stirred for 30 minutes at ambient temperature and 4-chloro-6-methoxy-7-(2-([N-methyl-N-(3-morpholinopropylsulphonyl)]amino)ethoxy)quinazoline (140mg, 0.3mmol) was added as a solid and the mixture stirred for 1.5 hours at ambient temperature. The mixture was poured onto a mixture of ether and water (50ml/50ml), the aqueous layer was adjusted to pH8 with 1M hydrochloric acid and was extracted with methylene chloride. The organic extract was washed with brine, dried (MgSO₄) and the solvent removed by evaporation. The residue was purified by column chromatography eluting with methylene chloride/methanol (95/5). The purified product was dissolved in methylene chloride (15ml) and methanol (2ml), and 2.2M ethereal hydrogen chloride (1ml) was added. The mixture was stirred for 30 minutes at ambient temperature and the volatiles were removed by evaporation. The residue was triturated with ether, collected by filtration and dried under vacuum at 70°C to give 6-methoxy-7-(2-([N-methyl-N-(3-morpholinopropylsulphonyl)]amino)ethoxy)-4-(oxindol-3-yl)quinazoline hydrochloride (100mg, 59%).

¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 2.1-2.2(m, 2H); 3.01(s, 3H); 3.15(t, 2H); 3.25(t, 2H); 3.33(t, 2H); 3.46(d, 2H); 3.7-3.8(m, 4H); 3.89(s, 3H); 3.99(d, 2H); 4.36(t, 2H); 7.0-7.1(m, 2H); 7.18(t, 1H); 7.35(s, 1H); 7.71(d, 1H); 7.79(s, 1H); 8.91(s, 1H)

MS - ESI: 556 [MH]⁺

The starting material was prepared as follows:

Diethyl azodicarboxylate (679mg, 3.9mmol) was added dropwise to a suspension of 7-hydroxy-6-methoxy-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (918mg, 3mmol), (prepared as described for the starting material in Example 22), triphenylphosphine (1g, 3.9mmol) and 2-([N-(t-butylcarbonyl)-N-methyl]amino)ethanol (682mg, 3.9mmol), (Synth. Commun. 1993, 23, 2443), in methylene chloride (20ml) and the mixture stirred for 1

hour at ambient temperature. Further 2-([N-(t-butylcarbonyl)-N-methyl]amino)ethanol (105mg, 0.6mmol), triphenylphosphine (786mg, 3mmol) and diethyl azodicarboxylate (522mg, 3mmol) were added and the mixture stirred for 30 minutes at ambient temperature. The mixture was concentrated to half volume by evaporation and purified by column chromatography eluting with methylene chloride/ether (7/3 increasing to 1/1) to give 6-methoxy-7-(2-([N-(t-butylcarbonyl)-N-methyl]amino)ethoxy)-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (1.3g, 98%).

¹H NMR Spectrum: (CDCl₃) 1.2(s, 9H); 1.45(s, 9H); 3.05(br s, 3H); 3.72(br s, 2H); 3.98(s, 3H); 4.25(br s, 2H); 5.95(s, 2H); 7.1(br s, 1H); 7.6(s, 1H); 8.2(s, 1H)

A solution of 6-methoxy-7-(2-([N-(t-butylcarbonyl)-N-methyl]amino)ethoxy)-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (1.39g, 3mmol) in methylene chloride (4ml) and TFA (4ml) was stirred at ambient temperature for 1 hour. Toluene was added, and the volatiles were removed by evaporation. The residue was triturated with ether and the resulting solid was collected by filtration. The solid was dissolved in water, sodium hydrogen carbonate was added and the aqueous mixture was extracted with methylene chloride. The organic extract was dried (MgSO₄) and the solvent removed by evaporation. The residue was triturated with ether and the solid was collected by filtration to give 6-methoxy-7-(2-(methylamino)ethoxy)-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (800mg, 73%).

¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 1.13(s, 9H); 2.72(s, 3H); 3.45(br s, 2H); 3.95(s, 3H); 4.5(t, 2H); 5.94(s, 2H); 7.31(s, 1H); 7.6(s, 1H); 8.47(s, 1H)

MS - ESI: 364 [MH]⁺

3-Morpholinopropane sulphonyl chloride (280mg, 1.1mmol), (WO 93 01181) followed by triethylamine (336μl, 2.4mmol) was added to a solution of 6-methoxy-7-(2-(methylamino)ethoxy)-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (350mg, 0.96mmol) in methylene chloride (15ml) and the mixture stirred for 30 minutes at ambient temperature. The mixture was partitioned between methylene chloride and water, the organic layer was separated, washed with brine, dried (MgSO₄) and the volatiles were removed by evaporation. The residue was purified by column chromatography eluting with ethyl acetate/methylene chloride (1/1) followed by methylene chloride/methanol (95/5). The purified product was triturated with ether, collected by filtration and dried under vacuum to

- 130 -

give 6-methoxy-7-(2-([N-methyl-N-(3-morpholinopropylsulphonyl)]amino)ethoxy)-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (320mg, 60%).

¹H NMR Spectrum: (DMSO-d₆) 1.11(s, 9H); 1.8-1.9(m, 2H); 2.25-2.35(m, 6H); 2.94(s, 3H); 3.19(t, 2H); 3.32(s, 3H); 3.51(t, 4H); 3.61(t, 2H); 3.9(s, 3H); 4.29(t, 2H); 5.9(s, 2H); 7.23(s, 1H); 7.52(s, 1H); 8.36(s, 1H)

MS - ESI: 577 [MNa]⁺

A solution of 6-methoxy-7-(2-([N-methyl-N-(3-morpholinopropylsulphonyl)]amino)ethoxy)-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (310mg, 0.55mmol) in saturated methanolic ammonia (15ml) was stirred at ambient temperature overnight. The volatiles were removed by evaporation, the solid residue was triturated with ether, collected by filtration and dried under vacuum to give 6-methoxy-7-(2-([N-methyl-N-(3-morpholinopropylsulphonyl)]amino)ethoxy)-3,4-dihydroquinazolin-4-one (245mg, 100%).

¹H NMR Spectrum: (DMSO-d₆) 1.8(t, 2H); 2.2-2.3(m, 6H); 2.94(s, 3H); 3.18(t, 2H); 3.51(t, 4H); 3.60(t, 2H); 3.88(s, 3H); 4.27(t, 2H); 7.18(s, 1H); 7.47(s, 1H); 7.99(s, 1H)

A solution of 6-methoxy-7-(2-([N-methyl-N-(3-morpholinopropylsulphonyl)]amino)ethoxy)-3,4-dihydroquinazolin-4-one (176mg, 0.4mmol) in thionyl chloride (4ml) containing DMF (100μl) was stirred at 45°C for 1 hour. The volatiles were removed by evaporation and by azeotrope with toluene. The residue was partitioned between methylene chloride and water, the aqueous phase was adjusted to pH8 with saturated aqueous sodium hydrogen carbonate solution. The organic layer was separated, washed with brine, dried (MgSO₄) and the solvent removed by evaporation. The residue was purified by column chromatography eluting with methanol/methylene chloride (a gradient from 3/100 to 10/90). The purified product was triturated with ether, collected by filtration and dried under vacuum to give 4-chloro-6-methoxy-7-(2-([N-methyl-N-(3-morpholinopropylsulphonyl)]amino)ethoxy)quinazoline (153mg, 71%).

¹H NMR Spectrum: (DMSO-d₆) 1.75-1.85(m, 2H); 2.25-2.35(m, 6H); 2.96(s, 3H); 3.2(t, 2H); 3.51(t, 4H); 3.65(t, 2H); 4.01(s, 3H); 4.4(t, 2H); 7.43(s, 1H); 7.54(s, 1H); 8.89(s, 1H)

Example 66

Using a procedure analogous to that described in Example 26, 6-(1-t-butoxycarbonylpyrrolidin-2-yl)oxindole (420mg, 1.4mmol) was added to sodium hydride (85mg, 2.2mmol) in THF (15ml) and the solution was treated with 4-chloro-6-methoxy-7-(3-morpholinopropoxy)quinazoline (235mg, 0.7mmol), (prepared as described for the starting material in Example 5), in DMF (2ml) to give, after work-up and purification, 4-(6-(1-t-butoxycarbonylpyrrolidin-2-yl)oxindol-3-yl)-6-methoxy-7-(3-morpholinopropoxy)quinazoline (135mg, 32%).

¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 1.1-1.4(m, 9H); 1.8(m, 3H); 2.3(m, 3H); 3.1(m, 2H); 3.4(t, 2H); 3.5(m, 4H); 3.8(m, 5H); 4.0(m, 2H); 4.3(t, 2H); 4.7-4.9(m, 1H); 6.9(m, 2H); 7.25(s, 1H); 7.6(d, 1H); 7.75(s, 1H); 8.95(s, 1H); 10.95(s, 1H)

MS - ESI: 604 [MH]⁺

Elemental analysis:	Found	C 64.6	H 6.9	N 11.0
C ₃₃ H ₄₁ N ₅ O ₆ 0.7H ₂ O	Requires	C 64.3	H 6.9	N 11.4%

15

The starting material was prepared as follows:-

A mixture of 6-(1-t-butoxycarbonylpyrrol-2-yl)oxindole (560mg, 1.9mmol), (prepared as described for the starting material in Example 63), and 10% palladium-on-charcoal catalyst (50mg) in ethyl acetate (30ml) was stirred under hydrogen at 1 atmosphere pressure for 3 days. The catalyst was removed by filtration through diatomaceous earth, the solvent was removed by evaporation and the residue triturated with ether/isohexanes (1/5) and collected by filtration to give 6-(1-t-butoxycarbonylpyrrolidin-2-yl)oxindole (480mg, 85%) as a pale yellow solid.

¹H NMR Spectrum: (DMSO-d₆) 1.25(br s, 9H); 1.75(m, 3H); 2.3(m, 1H); 3.4(s, 2H); 3.45(m, 2H); 4.75(br s, 1H); 6.6(s, 1H); 6.75(d, 1H); 7.1(d, 2H); 10.1(br s, 1H)

MS - ESI: 301 [M-H]⁺

Example 67

TFA (5ml) and triethylsilane (0.5g) were added to 4-(6-(1-t-butoxycarbonylpyrrolidin-2-yl)oxindol-3-yl)-6-methoxy-7-(3-

30

morpholinopropoxy)quinazoline (75mg, 0.12mmol), (prepared as described in Example 66), and the solution was stirred for 30 minutes. The volatiles were removed by evaporation and by azeotroping with toluene. The bright orange residue was dissolved in methanol and a solution of 1M ethereal hydrogen chloride was added. The resulting precipitate was collected
5 by filtration, washed with ether and dried at 60°C under vacuum for 2 hours to give 6-methoxy-7-(3-morpholinopropoxy)-4-(6-(pyrrolidin-2-yl)oxindol-3-yl)quinazoline hydrochloride (45mg, 63%).

¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 2.0(m, 3H); 2.3(m, 3H); 3.1(m, 2H); 3.3(m, 4H); 3.5(d, 2H); 3.75(t, 2H); 3.85(s, 3H); 4.0(d, 2H); 4.3(t, 2H); 4.5(m, 1H); 7.1(m, 2H); 7.3(s,
10 1H); 7.7(d, 1H); 7.75(s, 1H); 8.8(m, 2H); 11.2(s, 1H)

MS - ESI: 504 [MH]⁺

Elemental analysis:	Found	C 52.1	H 5.6	N 10.4
C ₂₈ H ₃₃ N ₅ O ₄ 0.7H ₂ O 3HCl	Requires	C 51.8	H 6.2	N 10.8%

15 Example 68

A solution of 5-aminosulphonyloxindole (283mg, 1.3mmol), (prepared as described for the starting material in Example 60), in DMF (3.5ml) was added dropwise to a suspension of sodium hydride (53mg, 1.3mmol, prewashed with hexane) in DMF (1.5ml). The mixture was stirred for 30 minutes at ambient temperature and a solution of 4-chloro-6-methoxy-7-(3-morpholinopropoxy)quinazoline (150mg, 0.44mmol), (prepared as described for the starting
20 material in Example 5), in DMF (4ml) was added. The mixture was stirred for 20 minutes at 65°C, DMSO (2ml) was added and heating was continued for 1 hour at 65°C. The mixture was allowed to cool, the volatiles were removed by evaporation and the residue was partitioned between ether and water. The aqueous layer was adjusted to pH7.8 with 1M
25 hydrochloric acid and the resulting precipitate was collected by filtration, washed with water and ether, and dried under vacuum. The solid was dissolved in methanol/methylene chloride (1/1) and 2.2M ethereal hydrogen chloride (1ml) was added. The mixture was stirred for 10 minutes at ambient temperature and the volatiles were removed by evaporation. The solid residue was triturated with ether, collected by filtration, washed with ether and dried under

vacuum to give **4-(5-aminosulphonyloxindol-3-yl)-6-methoxy-7-(3-morpholinopropoxy)quinazoline hydrochloride** (62mg, 24%).

¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 2.25-2.35(m, 2H); 3.15(t, 2H); 3.36(t, 2H); 3.55(d, 2H); 3.73(t, 2H); 3.93(s, 3H); 4.03(d, 2H); 4.33(t, 2H); 7.13(d, 1H); 7.35(s, 1H); 7.62(d, 1H);
 5 7.74(s, 1H); 8.1(s, 1H); 8.76(s, 1H)

MS - ESI: 514 [MH]⁺

Elemental analysis:	Found	C 49.2	H 5.2	N 11.8
---------------------	-------	--------	-------	--------

C ₂₄ H ₂₇ N ₃ O ₆ S 1.0H ₂ O 1.4HCl	Requires	C 49.5	H 5.3	N 12.0%
--	----------	--------	-------	---------

10 **Example 69**

A solution of 5-methylaminosulphonyloxindole (262mg, 1.16mmol) in DMF (3.5ml) was added dropwise to a suspension of sodium hydride (46mg, 1.16mmol, prewashed with hexane) in DMF (1.5ml). The mixture was stirred for 30 minutes at ambient temperature and 4-chloro-6-methoxy-7-(3-morpholinopropoxy)quinazoline (130mg, 0.38mmol), (prepared as
 15 described for the starting material in Example 5), was added as a solid. The mixture was stirred for 45 minutes at 60°C and was then partitioned between ether and water. The aqueous layer was separated and was adjusted to pH7-8 with 1M hydrochloric acid. The precipitate was collected by filtration, washed with water and ether, and dried under vacuum. The solid was suspended in methylene chloride/methanol (1/1) and 2.2M ethereal hydrogen chloride
 20 (1.5ml) was added. The mixture was stirred for 10 minutes at ambient temperature and the volatiles were removed by evaporation. The solid residue was triturated with ether, collected by filtration, washed with ether and dried under vacuum at 60°C to give **6-methoxy-4-(5-methylaminosulphonyloxindol-3-yl)-7-(3-morpholinopropoxy)quinazoline hydrochloride** (116mg, 53%).

25 ¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 2.29(s, 3H); 3.16(t, 2H); 3.33(t, 2H); 3.54(d, 2H); 3.74(t, 2H); 3.93(s, 3H); 4.02(d, 2H); 4.33(t, 2H); 7.14(d, 1H); 7.34(s, 1H); 7.52(d, 1H); 7.76(s, 1H); 8.03(s, 1H); 8.67(s, 1H)

MS - ESI: 528 [MH]⁺

Elemental analysis:	Found	C 50.3	H 5.4	N 11.3
---------------------	-------	--------	-------	--------

30 C ₂₅ H ₂₉ N ₃ O ₆ S 0.85H ₂ O 1.6HCl	Requires	C 49.9	H 5.4	N 11.6%
--	----------	--------	-------	---------

- 134 -

The starting material was prepared as follows:

Methylamine (2.7ml of a 2.9M solution in chloroform) was added to a suspension of 5-chlorosulphonyloxindole (900mg, 3.9mmol), (prepared as described for the starting material in Example 60), in a mixture of methylene chloride (20ml) and ethanol (20ml). The mixture was stirred for 15 minutes at ambient temperature, the volatiles were removed by evaporation. The solid residue was suspended in water, collected by filtration, washed with water, ether, and dried under vacuum over phosphorus pentoxide to give 5-methylaminosulphonyloxindole (880mg, 90%).

¹H NMR Spectrum: (DMSO_d₆) 2.37(d, 3H); 3.58(s, 2H); 6.96(d, 1H); 7.25(br s, 1H); 7.58(s, 1H); 7.60(d, 1H)

Example 70

A solution of 5-methylaminosulphonyloxindole (252mg, 1.1mmol), (prepared as described for the starting material in Example 69), in DMF (2.5ml) was added dropwise to a suspension of sodium hydride (45mg, 1.1mmol, prewashed with hexane) in DMF (1.5ml). The mixture was stirred for 30 minutes at ambient temperature and 4-chloro-6-methoxy-7-(2-methoxyethoxy)quinazoline (100mg, 0.37mmol), (prepared as described for the starting material in Example 2), was added as a solid. The mixture was then stirred for 45 minutes at 60°C and was then partitioned between ether and water. The aqueous layer was separated and was adjusted to pH7-8 with 1M hydrochloric acid. The precipitate was collected by filtration, washed with water and ether, and dried under vacuum. The solid was suspended in methylene chloride/methanol (1/1) and 2.2M ethereal hydrogen chloride (1.5ml) was added. The mixture was stirred for 10 minutes at ambient temperature and the volatiles were removed by evaporation. The solid residue was triturated with ether, collected by filtration, washed with ether and dried under vacuum at 60°C to give 6-methoxy-7-(2-methoxyethoxy)-4-(5-methylaminosulphonyloxindol-3-yl)quinazoline hydrochloride (123mg, 68%).

¹H NMR Spectrum: (DMSO_d₆, CF₃CO₂D) 2.29(s, 3H); 3.36(s, 3H); 3.79(t, 2H); 3.93(s, 3H); 4.35(t, 2H); 7.14(d, 1H); 7.32(s, 1H); 7.53(d, 1H); 7.73(s, 1H); 8.04(s, 1H); 8.73(s, 1H)

MS - ESI: 459 [MH]⁺

Elemental analysis:	Found	C 51.6	H 4.8	N 11.1
C ₂₁ H ₂₂ N ₄ O ₆ S 0.5H ₂ O 0.65HCl	Requires	C 51.4	H 4.9	N 11.4%

Example 71

A solution of oxindole (116mg, 0.87mmol) in DMF (1.5ml) was added to a suspension of sodium hydride (35mg, 0.87mmol, prewashed with hexane) in DMF (2.5ml).

- 5 The mixture was stirred for 30 minutes at ambient temperature and 4-chloro-6-methoxy-7-(2-([N-methyl-N-(1-methyl-1H-imidazol-4-ylsulphonyl)]amino)ethoxy)quinazoline (120mg, 0.29mmol) was added as a solid. The mixture was stirred for 10 minutes at ambient temperature followed by 20 minutes at 50°C. The mixture was allowed to cool, and was partitioned between ether and water. The aqueous layer was separated and adjusted to pH7.7
- 10 with 1M hydrochloric acid. The mixture was left to stand for 1.5 hours and the resulting precipitate was collected by filtration, washed with water, ether and dried under vacuum. The solid was purified by column chromatography eluting with methylene chloride/methanol (95/5 and then 90/10). The purified solid was dissolved in methylene chloride/methanol (2/1) and 2M ethereal hydrogen chloride (1.5ml) was added. The mixture was stirred for 10 minutes at
- 15 ambient temperature and the volatiles were removed by evaporation. The solid residue was triturated with ether, collected by filtration, washed with ether, dried under vacuum to give 6-methoxy-7-(2-([N-methyl-N-(1-methyl-1H-imidazol-4-ylsulphonyl)]amino)ethoxy)-4-(oxindol-3-yl)quinazoline hydrochloride (106mg, 65%).

- ¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 2.89(s, 3H); 3.56(t, 2H); 3.72(s, 3H); 3.86(s, 3H);
- 20 4.39(t, 2H); 7-7.1(m, 2H); 7.17(t, 1H); 7.3(s, 1H); 7.71(d, 2H); 7.76(s, 1H); 7.89(d, 1H); 8.91(s, 1H)

MS - ESI: 509 [MH]⁺

Elemental analysis:	Found	C 51.3	H 4.6	N 14.8
C ₂₄ H ₂₄ N ₆ O ₃ S 0.1H ₂ O 1.3HCl	Requires	C 51.7	H 4.6	N 15.0%

25

The starting material was prepared as follows:

- Using a procedure identical to that described for the synthesis of 6-methoxy-7-(2-([N-methyl-N-(3-morpholinopropylsulphonyl)]amino)ethoxy)-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one in Example 65, 6-methoxy-7-(2-(methylamino)ethoxy)-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (400mg, 1.1mmol), (prepared as
- 30 described for the starting material in Example 65), was treated with (1-methyl-1H-imidazol-4-

yl)sulphonyl chloride (219mg, 1.2mmol) and purified by column chromatography eluting with methanol/acetonitrile/methylene chloride (5/35/60) to give 6-methoxy-7-(2-([N-methyl-N-(1-methyl-1H-imidazol-4-ylsulphonyl)]amino)ethoxy)-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (507mg, 73%).

- 5 ¹H NMR Spectrum: (DMSO-d₆) 1.12(s, 9H); 2.84(s, 3H); 3.48(t, 2H); 3.69(s, 3H); 3.89(s, 3H); 4.31(t, 2H); 5.90(s, 2H); 7.19(s, 1H); 7.51(s, 1H); 7.79(s, 1H); 7.83(s, 1H); 8.36(s, 1H)

To a solution of 6-methoxy-7-(2-([N-methyl-N-(1-methyl-1H-imidazol-4-ylsulphonyl)]amino)ethoxy)-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (470mg, 0.93mmol) in methylene chloride (15ml) was added a saturated solution of methanolic ammonia (50ml). The mixture was stirred for 2 days at ambient temperature and the volatiles were removed by evaporation. The solid residue was triturated with ether, collected by filtration, washed with ether and dried under vacuum to give 6-methoxy-7-(2-([N-methyl-N-(1-methyl-1H-imidazol-4-ylsulphonyl)]amino)ethoxy)-3,4-dihydroquinazolin-4-one (300mg, 82%).

- 15 ¹H NMR Spectrum: (DMSO-d₆) 2.84(s, 3H); 3.48(t, 2H); 3.69(s, 3H); 3.87(s, 3H); 4.29(t, 2H); 7.14(s, 1H); 7.45(s, 1H); 7.79(s, 1H); 7.83(s, 1H); 7.98(s, 1H)

A solution of 6-methoxy-7-(2-([N-methyl-N-(1-methyl-1H-imidazol-4-ylsulphonyl)]amino)ethoxy)-3,4-dihydroquinazolin-4-one (250mg, 0.64mmol) in thionyl chloride (5ml) containing DMF (150μl) was heated at 50°C for 20 minutes. The mixture was allowed to cool, toluene was added and the volatiles were removed by evaporation. The residue was dissolved in methylene chloride and cooled to 5°C. Saturated aqueous sodium hydrogen carbonate solution was added until the aqueous layer maintained pH7.7 and the aqueous layer was extracted with methylene chloride. The combined organic extract was washed with brine, dried (MgSO₄) and the solvent removed by evaporation. The residue was triturated with ether, collected by filtration and dried under vacuum to give 4-chloro-6-methoxy-7-(2-([N-methyl-N-(1-methyl-1H-imidazol-4-ylsulphonyl)]amino)ethoxy)quinazoline (285mg, 91%).

- 25 ¹H NMR Spectrum: (DMSO-d₆) 2.86(s, 3H); 3.53(t, 2H); 3.68(s, 3H); 4.0(s, 3H); 4.42(t, 2H); 7.41(s, 1H); 7.49(s, 1H); 7.78(s, 1H); 7.83(s, 1H); 8.89(s, 1H)

Example 72

Using a procedure analogous to that described in Example 26, oxindole (0.32g, 2.4mmol) was added to sodium hydride (110mg, 2.8mmol) in DMF (15ml) and the solution was treated with 4-chloro-6-morpholinoquinazoline (200mg, 8mmol), (EP 0566226), in DMF (8ml). The product so obtained was treated with ethereal hydrogen chloride to give 4-(oxindol-3-yl)-6-morpholinoquinazoline hydrochloride (120mg, 43%).

¹H NMR Spectrum (DMSO-d₆, CF₃CO₂D) 3.2(m, 4H); 3.7(m, 4H); 7.0(m, 2H); 7.15(t, 1H); 7.6(m, 4H); 8.9(s, 1H), 11.2(s, 1H)

MS - ESI: 347 [MH]⁺

10	Elemental analysis:	Found	C 61.8	H 4.9	N 14.4
	C ₂₀ H ₁₈ N ₄ O ₂ 0.3H ₂ O 1HCl	Requires	C 61.9	H 5.1	N 14.4%

Example 73

A solution of 5-N,N-dimethylaminosulphonyloxindole (268mg, 1.1mmol) in DMF (3ml) was added dropwise to a suspension of sodium hydride (45mg, 1.1mmol, prewashed with hexane) in DMF (1.5ml). The mixture was stirred for 30 minutes at ambient temperature and 4-chloro-6-methoxy-7-(2-methoxyethoxy)quinazoline (100mg, 0.37mmol), (prepared as described for the starting material in Example 2), was added as a solid. The mixture was stirred for 10 minutes at ambient temperature and then 1 hour at 45°C. The mixture was allowed to cool and was partitioned between ether and water. The aqueous layer was separated, adjusted to pH7 with 1M hydrochloric acid and extracted with ethyl acetate. The organic extract was washed with brine, dried (MgSO₄) and the solvent removed by evaporation. The residue was purified by column chromatography eluting with methylene chloride/acetonitrile/methanol (60/36/4). The purified solid was collected, dried under vacuum and was dissolved in methylene chloride/methanol (3/1). 2.2M Ethereal hydrogen chloride (1ml) was added and the mixture stirred for 10 minutes at ambient temperature. The volatiles were removed by evaporation, the solid residue was triturated with ether, collected by filtration and dried under vacuum to give 4-(5-N,N-dimethylaminosulphonyloxindol-3-yl)-6-methoxy-7-(2-methoxyethoxy)quinazoline hydrochloride (75mg, 41%).

¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 2.50(s, 6H); 3.35(s, 3H); 3.77(t, 2H); 3.91(s, 3H); 4.35(t, 2H); 7.16(d, 1H); 7.30(s, 1H); 7.46(d, 1H); 7.81(br s, 1H); 7.98(br s, 1H); 8.69(s, 1H)

MS - EI: 472 [M]⁺

Elemental analysis:		Found	C 52.5	H 5.1	N 11.1
5	C ₂₂ H ₂₄ N ₄ O ₆ S 0.5H ₂ O 0.7HCl	Requires	C 52.1	H 5.1	N 11.0%

The starting material was prepared as follows:

Dimethylamine (430μl, 8.6mmol) was added to a solution of 5-chlorosulphonyloxindole (990mg, 4.3mmol), (prepared as described for the starting material in Example 60), in a mixture of methylene chloride (20ml), acetonitrile (20ml) and ethanol (20ml). The mixture was stirred for 1 hour at ambient temperature, dimethylamine (86μl) was added and stirring continued for 30 minutes at ambient temperature. The volatiles were removed by evaporation and the residue was purified by column chromatography eluting with methylene chloride/methanol 97/3 to give 5-N,N-dimethylaminosulphonyloxindole (315mg, 30%).

¹H NMR Spectrum: (DMSO-d₆) 2.57(s, 6H); 3.60(s, 2H); 7.01(d, 1H); 7.55(s, 1H); 7.59(s, 1H)

Example 74

A solution of oxindole (153mg, 1.15mmol) in DMF (3ml) was added to a suspension of sodium hydride (46mg, 1.15mmol, prewashed with hexane) in DMF (1ml) and the mixture was stirred for 30 minutes at ambient temperature. 4-Chloro-6-methoxy-7-(2-(2-methoxyethoxy)ethoxy)quinazoline (120mg, 0.38mmol) was added as a solid and the mixture was stirred for 10 minutes at ambient temperature and then for 45 minutes at 50°C. The mixture was allowed to cool and was partitioned between ether and water. The aqueous layer was separated and adjusted to pH7.5 with 1M hydrochloric acid. The resulting precipitate was collected by filtration washed with water followed by ether and dried over phosphorus pentoxide. The solid was dissolved in methylene chloride/methanol and 2.8M ethereal hydrogen chloride (1ml) was added. The volatiles were removed by evaporation, the solid residue was triturated with ether, collected by filtration and dried under vacuum to give 6-methoxy-7-(2-(2-methoxyethoxy)ethoxy)-4-(oxindol-3-yl)quinazoline hydrochloride (124mg, 73%).

¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 3.27(s, 3H); 3.49(m, 2H); 3.64(m, 2H); 3.87(br s, 5H); 4.32(m, 2H); 7.0-7.1(m, 2H); 7.18(t, 1H); 7.30(s, 1H); 7.7(d, 1H); 7.75(s, 1H); 8.9(s, 1H)
MS - ESI: 410 [MH]⁺

Elemental analysis:	Found	C 59.1	H 5.6	N 9.4
5 C ₂₂ H ₂₃ N ₃ O ₅ 0.4H ₂ O 0.75HCl	Requires	C 59.5	H 5.6	N 9.5%

The starting material was prepared as follows:

Diethyl azodicarboxylate (864μl, 5.5mmol) was added dropwise to a mixture of 7-hydroxy-6-methoxy-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (1.2g, 3.9mmol)
10 (prepared as described for the starting material in Example 22), triphenylphosphine (1.44g, 5.5mmol) and 2-(2-methoxyethoxy)ethanol (653μl, 5.5mmol) in methylene chloride (70ml) cooled at 0°C. The mixture was stirred for 1.5 hours at ambient temperature and the solvent was removed by evaporation. The residue was purified by column chromatography eluting with a mixture of ethyl acetate/methylene chloride (50/50 followed by 80/20). The purified
15 solid was suspended in ether, collected by filtration and dried under vacuum to give 6-methoxy-7-(2-(2-methoxyethoxy)ethoxy)-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (1.70g, 100%).

¹H NMR Spectrum: (DMSO-d₆) 1.13(s, 9 H); 3.26(s, 3H); 3.5(m, 2H); 3.65(m, 2H); 3.85(m, 2H); 3.91(s, 3H); 4.3(m, 2H); 5.9(s, 2H); 7.2(s, 1H); 7.5(s, 1H); 8.4(s, 1H)

20 Saturated methanolic ammonia (20ml) was added to a solution of 6-methoxy-7-(2-(2-methoxyethoxy)ethoxy)-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (2.26g, 5.5mmol) in a mixture of ethanol (40ml) and methylene chloride (15ml). The mixture was stirred for 24 hours at ambient temperature, and further methanolic ammonia (20ml) was added. The mixture was stirred for a further 24 hours at ambient temperature and the volatiles
25 were removed by evaporation. The residue was triturated with ether, collected by filtration, dried under vacuum to give 6-methoxy-7-(2-(2-methoxyethoxy)ethoxy)-3,4-dihydroquinazolin-4-one (975mg, 78%).

¹H NMR Spectrum: (DMSO-d₆) 3.25(s, 3H); 3.45(t, 2H); 3.6(t, 2H); 3.8(t, 2H); 3.9(s, 3H); 4.2(t, 2H); 7.15(s, 1H); 7.45(s, 1H); 8.0(s, 1H)

30 MS - EI: 294 [M]⁺

- 140 -

A solution of 6-methoxy-7-(2-(2-methoxyethoxy)ethoxy)-3,4-dihydroquinazolin-4-one (930mg, 3.16mmol) in thionyl chloride (15ml) and DMF (150µl) was heated at 60°C for 1.5 hours. The mixture was allowed to cool and the volatiles were removed by evaporation and by azeotrope with toluene. The residue was dissolved in methylene chloride and 5% aqueous sodium hydrogen carbonate solution was added until the aqueous layer was at pH8. The organic layer was separated, washed with brine, dried (MgSO₄) and the solvent removed by evaporation. The residue was purified by flash chromatography eluting with ethyl acetate to give 4-chloro-6-methoxy-7-(2-(2-methoxyethoxy)ethoxy)quinazoline (863mg, 87%).

¹H NMR Spectrum: (DMSO-d₆) 3.24(s, 3H); 3.47(m, 2H); 3.62(m, 2H); 3.84(t, 2H); 4.01(s, 3H); 4.25(t, 2H); 7.41(s, 1H); 7.49(s, 1H); 8.88(s, 1H)

Example 75

A solution of 5-aminosulphonyloxindole (265mg, 1.25mmol), (prepared as described for the starting material in Example 60), in DMF (2.5ml) was added to a suspension of sodium hydride (50mg, 1.25mmol, prewashed with pentane) in DMF (1.5ml). The mixture was stirred for 30 minutes at ambient temperature and 4-chloro-6-methoxy-7-(2-(2-methoxyethoxy)ethoxy)quinazoline (265mg, 1.25mmol), (prepared as described for the starting material in Example 74), was added followed by DMF (2ml) and DMSO (0.5ml). The mixture was heated at 60°C for 45 minutes, then allowed to cool and partitioned between ether and water. The aqueous layer was separated and adjusted to pH7 with 1M hydrochloric acid. The resulting precipitate was collected by filtration, washed with water and ether, and dried under vacuum. The solid was dissolved in methylene chloride/methanol and 2.8M ethereal hydrogen chloride (1ml) was added. The volatiles were removed by evaporation, the solid was triturated with ether, collected by filtration and dried under vacuum to give 4-(5-aminosulphonyloxindol-3-yl)-6-methoxy-7-(2-(2-methoxyethoxy)ethoxy)quinazoline hydrochloride (106mg, 50%).

¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 3.26(s, 3H); 3.49(m, 2H); 3.64(m, 2H); 3.86(br s, 2H); 3.91(s, 3H); 4.34(m, 2H); 7.10(d, 1H); 7.31(s, 1H); 7.59(d, 1H); 7.71(s, 1H); 8.09(s, 1H); 8.71(s, 1H)

MS - ESI: 489 [MH]⁺

Elemental analysis:

Found

C 51.4 H 4.9 N 10.8

- 141 -

$C_{22}H_{24}N_4O_7S \cdot 0.7H_2O \cdot 0.45HCl$ Requires C 51.1 H 5.03 N 10.8%

Example 76

A solution of 5-N,N-dimethylaminosulphonyloxindole (256mg, 1.07mmol),
 5 (prepared as described for the starting material in Example 73), in DMF (3ml) was added to a
 suspension of sodium hydride (43mg, 1.07mmol, prewashed with pentane) in DMF (2ml).
 The mixture was stirred for 30 minutes at ambient temperature and 4-chloro-6-methoxy-7-(3-
 morpholinopropoxy)quinazoline (120mg, 0.36mmol), (prepared as described for the starting
 material in Example 5), was added. The mixture was heated at 65°C for 1 hour, then allowed
 10 to cool and partitioned between ether and water. The aqueous layer was separated and was
 adjusted to pH8 with 1M hydrochloric acid. The aqueous layer was extracted with methylene
 chloride, the organic extracts were combined, washed with brine, dried ($MgSO_4$) and the
 solvent removed by evaporation. The residue was purified by column chromatography eluting
 with methylene chloride/methanol (94/6). The purified product was dissolved in methylene
 15 chloride/methanol (3/1) and 2.8M ethereal hydrogen chloride (1ml) was added. The volatiles
 were removed by evaporation, the residue was triturated with ether, collected by filtration and
 dried under vacuum to give 4-(5-N,N-dimethylaminosulphonyloxindol-3-yl)-6-methoxy-7-
 (3-morpholinopropoxy)quinazoline hydrochloride (109mg, 48%).

1H NMR Spectrum: ($DMSO-d_6$, CF_3CO_2D) 2.3-2.4(m, 2H); 2.51(s, 6H); 3.1-3.2(m, 2H); 3.3-
 20 3.4(m, 2H); 3.6(d, 2H); 3.73(t, 2H); 3.92(s, 3H); 4.05(d, 2H); 4.33(t, 2H); 7.17(d, 1H); 7.3(s,
 1H); 7.46(d, 1H); 7.83(br s, 1H); 7.98(s, 1H); 8.67(s, 1H)

MS - EI: 541 [M.]⁺

Elemental analysis:

Found C 50.3 H 5.9 N 10.9

$C_{26}H_{31}N_5O_6S \cdot 1.5H_2O \cdot 1.5HCl$

Requires C 50.4 H 5.9 N 11.0%

25 0.17ether

Example 77

The following illustrate representative pharmaceutical dosage forms containing the
 compound of formula I, or a pharmaceutically acceptable salt thereof (hereafter compound X),
 30 for therapeutic or prophylactic use in humans:

- 142 -

(a)	<u>Tablet I</u>	<u>mg/tablet</u>
	Compound X	100
	Lactose Ph.Eur.....	182.75
	Croscarmellose sodium	12.0
5	Maize starch paste (5% w/v paste)	2.25
	Magnesium stearate	3.0
(b)	<u>Tablet II</u>	<u>mg/tablet</u>
	Compound X	50
10	Lactose Ph.Eur.....	223.75
	Croscarmellose sodium	6.0
	Maize starch.....	15.0
	Polyvinylpyrrolidone (5% w/v paste).....	2.25
	Magnesium stearate	3.0
15		
(c)	<u>Tablet III</u>	<u>mg/tablet</u>
	Compound X	1.0
	Lactose Ph.Eur.....	93.25
	Croscarmellose sodium	4.0
20	Maize starch paste (5% w/v paste)	0.75
	Magnesium stearate	1.0
(d)	<u>Capsule</u>	<u>mg/capsule</u>
	Compound X	10
	Lactose Ph.Eur.....	488.5
25	Magnesium stearate	1.5
(e)	<u>Injection I</u>	<u>(50 mg/ml)</u>
	Compound X	5.0% w/v
	1N Sodium hydroxide solution.....	15.0% v/v
30	0.1N Hydrochloric acid	
	(to adjust pH to 7.6)	

- 143 -

Polyethylene glycol 400 4.5% w/v

Water for injection to 100%

(f) Injection II (10 mg/ml)

5 Compound X 1.0% w/v

Sodium phosphate BP 3.6% w/v

0.1N Sodium hydroxide solution..... 15.0% v/v

Water for injection to 100%

10 (g) Injection III (1 mg/ml, buffered to pH6)

Compound X 0.1% w/v

Sodium phosphate BP 2.26% w/v

Citric acid 0.38% w/v

Polyethylene glycol 400 3.5% w/v


15 Water for injection to 100%

Note

The above formulations may be obtained by conventional procedures well known in the pharmaceutical art. The tablets (a)-(c) may be enteric coated by conventional means, for example to provide a coating of cellulose acetate phthalate.

CLAIMS

1. A compound of the formula I:



(I)

[wherein:

R² represents hydroxy, halogeno, C₁₋₃alkyl, C₁₋₃alkoxy, C₁₋₃alkanoyloxy, trifluoromethyl, cyano, amino, nitro, C₂₋₄alkanoyl, C₁₋₄alkanoylamino, C₁₋₄alkoxycarbonyl, C₁₋₄alkylthio, C₁₋

15 ₄alkylsulphinyl, C₁₋₄alkylsulphonyl, carbamoyl, N-C₁₋₄alkylcarbamoyl, N,N-di(C₁₋
₄alkyl)carbamoyl, aminosulphonyl, N-C₁₋₄alkylaminosulphonyl, N,N-di(C₁₋

₄alkyl)aminosulphonyl, C₁₋₄alkylsulphonylamino, or a group R⁴X¹ wherein X¹ represents a direct bond, C₂₋₄alkanoyl, -CONR⁵R⁶-, -SO₂NR⁷R⁸- or -SO₂R⁹- (wherein R⁵ and R⁷, each independently represents hydrogen or C₁₋₂alkyl and R⁶, R⁸ and R⁹ each independently represents

20 C₁₋₄alkyl and wherein R⁴ is linked to R⁶, R⁸ or R⁹) and R⁴ represents phenyl or a 5 or 6-
membered heterocyclic group with one or two heteroatoms, selected independently from O, S
and N, which heterocyclic group may be saturated or unsaturated and which phenyl or
heterocyclic group may bear one or two substituents selected from hydroxy, halogeno, C₁₋
₃alkyl, C₁₋₃alkoxy, C₁₋₃alkanoyloxy, trifluoromethyl, cyano, amino, nitro and C₁₋

25 ₄alkoxycarbonyl;

n is an integer from 0 to 4;

R¹ represents hydrogen, C₁₋₄alkyl, C₁₋₄alkoxymethyl, di(C₁₋₄alkoxy)methyl or
C₁₋₄alkanoyl;

m is an integer from 0 to 4; and

30 R³ represents hydroxy, halogeno, nitro, trifluoromethyl, C₁₋₃alkyl, cyano, amino or R¹⁰X²
[wherein X² represents -O-, -CH₂-, -S-, -SO-, -SO₂-, -NR¹¹CO-, -CONR¹²-, -SO₂NR¹³-, -

NR¹⁴SO₂- or -NR¹⁵- (wherein R¹¹, R¹², R¹³, R¹⁴ and R¹⁵ each independently represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl), or X² represents a direct bond, and R¹⁰ is selected from one of the following seventeen groups:

- 1) hydrogen or C₁₋₃alkyl which may be unsubstituted or which may be substituted with one or more groups selected from hydroxy, fluoro and amino;
- 2) C₁₋₃alkylX³COR¹⁶ (wherein X³ represents -O- or -NR¹⁷- (in which R¹⁷ represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R¹⁶ represents C₁₋₃alkyl, -NR¹⁸R¹⁹- or -OR²⁰- (wherein R¹⁸, R¹⁹ and R²⁰ which may be the same or different each represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl));
- 3) C₁₋₃alkylX⁴R²¹ (wherein X⁴ represents -O-, -S-, -SO-, -SO₂-, -OCO-, -NR²²CO-, -CONR²³-, -SO₂NR²⁴-, -NR²⁵SO₂- or -NR²⁶- (wherein R²², R²³, R²⁴, R²⁵ and R²⁶ each independently represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R²¹ represents hydrogen, C₁₋₃alkyl, cyclopentyl, cyclohexyl or a 5 or 6 membered saturated heterocyclic group with one or two heteroatoms, selected independently from O, S and N, which C₁₋₃alkyl group may bear one or two substituents selected from oxo, hydroxy, halogeno and C₁₋₄alkoxy and which cyclic group may bear one or two substituents selected from oxo, hydroxy, halogeno, C₁₋₄alkyl, C₁₋₄hydroxyalkyl and C₁₋₄alkoxy);
- 4) C₁₋₃alkylX⁵C₁₋₃alkylX⁶R²⁷ (wherein X⁵ and X⁶ which may be the same or different are each -O-, -S-, -SO-, -SO₂-, -NR²⁸CO-, -CONR²⁹-, -SO₂NR³⁰-, -NR³¹SO₂- or -NR³²- (wherein R²⁸, R²⁹, R³⁰, R³¹ and R³² each independently represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R²⁷ represents hydrogen or C₁₋₃alkyl);
- 5) C₁₋₃alkylR³³ (wherein R³³ is a 5 or 6 membered saturated heterocyclic group with one or two heteroatoms, selected independently from O, S and N, which heterocyclic group may bear one or two substituents selected from oxo, hydroxy, halogeno, C₁₋₄alkyl, C₁₋₄hydroxyalkyl and C₁₋₄alkoxy);
- 6) C₂₋₃alkenylR³³ (wherein R³³ is as defined herein);
- 7) C₂₋₃alkynylR³³ (wherein R³³ is as defined herein);
- 8) R³⁴ (wherein R³⁴ represents a pyridone group, a phenyl group or a 5 or 6-membered aromatic heterocyclic group with 1 to 3 heteroatoms selected from O, N and S, which pyridone, phenyl or heterocyclic group may carry up to 5 substituents selected from hydroxy, halogeno, amino, C₁₋₄alkyl, C₁₋₄alkoxy, C₁₋₄hydroxyalkyl, C₁₋₄aminoalkyl, C₁₋₄alkylamino, C₁₋

- 146 -

- hydroxyalkoxy, carboxy, cyano, $-\text{CONR}^{35}\text{R}^{36}$ and $-\text{NR}^{37}\text{COR}^{38}$ (wherein R^{35} , R^{36} , R^{37} and R^{38} , which may be the same or different, each represents hydrogen, C_{1-4} alkyl or C_{1-3} alkoxy C_{2-3} alkyl));
- 9) C_{1-3} alkyl R^{34} (wherein R^{34} is as defined herein);
- 5 10) C_{2-3} alkenyl R^{34} (wherein R^{34} is as defined herein);
- 11) C_{2-3} alkynyl R^{34} (wherein R^{34} is as defined herein);
- 12) C_{1-3} alkyl X^7R^{34} (wherein X^7 represents $-\text{O}-$, $-\text{S}-$, $-\text{SO}-$, $-\text{SO}_2-$, $-\text{NR}^{39}\text{CO}-$, $-\text{CONR}^{40}-$, $-\text{SO}_2\text{NR}^{41}-$, $-\text{NR}^{42}\text{SO}_2-$ or $-\text{NR}^{43}-$ (wherein R^{39} , R^{40} , R^{41} , R^{42} and R^{43} each independently represents hydrogen, C_{1-3} alkyl or C_{1-3} alkoxy C_{2-3} alkyl) and R^{34} is as defined herein);
- 10 13) C_{2-3} alkenyl X^8R^{34} (wherein X^8 represents $-\text{O}-$, $-\text{S}-$, $-\text{SO}-$, $-\text{SO}_2-$, $-\text{NR}^{44}\text{CO}-$, $-\text{CONR}^{45}-$, $-\text{SO}_2\text{NR}^{46}-$, $-\text{NR}^{47}\text{SO}_2-$ or $-\text{NR}^{48}-$ (wherein R^{44} , R^{45} , R^{46} , R^{47} and R^{48} each independently represents hydrogen, C_{1-3} alkyl or C_{1-3} alkoxy C_{2-3} alkyl) and R^{34} is as defined herein);
- 14) C_{2-3} alkynyl X^9R^{34} (wherein X^9 represents $-\text{O}-$, $-\text{S}-$, $-\text{SO}-$, $-\text{SO}_2-$, $-\text{NR}^{49}\text{CO}-$, $-\text{CONR}^{50}-$, $-\text{SO}_2\text{NR}^{51}-$, $-\text{NR}^{52}\text{SO}_2-$ or $-\text{NR}^{53}-$ (wherein R^{49} , R^{50} , R^{51} , R^{52} and R^{53} each independently represents hydrogen, C_{1-3} alkyl or C_{1-3} alkoxy C_{2-3} alkyl) and R^{34} is as defined herein);
- 15 15) C_{1-3} alkyl $\text{X}^{10}\text{C}_{1-3}$ alkyl R^{34} (wherein X^{10} represents $-\text{O}-$, $-\text{S}-$, $-\text{SO}-$, $-\text{SO}_2-$, $-\text{NR}^{54}\text{CO}-$, $-\text{CONR}^{55}-$, $-\text{SO}_2\text{NR}^{56}-$, $-\text{NR}^{57}\text{SO}_2-$ or $-\text{NR}^{58}-$ (wherein R^{54} , R^{55} , R^{56} , R^{57} and R^{58} each independently represents hydrogen, C_{1-3} alkyl or C_{1-3} alkoxy C_{2-3} alkyl) and R^{34} is as defined herein);
- 20 16) R^{33} (wherein R^{33} is as defined herein); and
- 17) C_{1-3} alkyl $\text{X}^{10}\text{C}_{1-3}$ alkyl R^{33} (wherein X^{10} and R^{33} are as defined herein)];
- and salts thereof.

2. A compound as claimed in claim 1 wherein R^1 is hydrogen.

25

3. A compound as claimed in claim 1 or claim 2 wherein R^2 is halogeno, trifluoromethyl, cyano, nitro, C_{2-3} alkanoyl, C_{1-3} alkoxycarbonyl, C_{1-3} alkylsulphinyl, C_{1-3} alkylsulphonyl, carbamoyl, $\text{N}-\text{C}_{1-3}$ alkylcarbamoyl, $\text{N,N}-\text{di}(\text{C}_{1-3}\text{alkyl})\text{carbamoyl}$, aminosulphonyl, $\text{N}-\text{C}_{1-3}$ alkylaminosulphonyl, $\text{N,N}-\text{di}(\text{C}_{1-3}\text{alkyl})\text{aminosulphonyl}$, or a group
- 30 R^4X^1 (wherein X^1 is as defined in claim 1 and R^4 represents phenyl or a 5 or 6-membered heterocyclic group with one or two heteroatoms, selected independently from O, S and N, of

which at least one is N, which heterocyclic group may be saturated or unsaturated and which phenyl or heterocyclic group may bear one or two substituents selected from hydroxy, halogeno, C₁₋₂alkyl, C₁₋₂alkoxy, C₁₋₂alkanoyloxy, trifluoromethyl, cyano, amino, nitro and C₁₋₂alkoxycarbonyl).

5

4. A compound as claimed in any one of the preceding claims wherein R³ is hydroxy, halogeno, nitro, trifluoromethyl, C₁₋₃alkyl, cyano, amino or R¹⁰X² [wherein X² is as defined in claim 1 and R¹⁰ is selected from one of the following fifteen groups:


- 1) C₁₋₃alkyl which may be unsubstituted or substituted with one or more fluorine atoms, or C₂₋₃alkyl which may be unsubstituted or substituted with one or two groups selected from hydroxy and amino;
- 2) 2-(3,3-dimethylureido)ethyl, 3-(3,3-dimethylureido)propyl, 2-(3-methylureido)ethyl, 3-(3-methylureido)propyl, 2-ureidoethyl, 3-ureidopropyl, 2-(N,N-dimethylcarbamoyloxy)ethyl, 3-(N,N-dimethylcarbamoyloxy)propyl, 2-(N-methylcarbamoyloxy)ethyl, 3-(N-methylcarbamoyloxy)propyl, 2-(carbamoyloxy)ethyl, 3-(carbamoyloxy)propyl;
- 3) C₂₋₃alkylX⁴R²¹ (wherein X⁴ is as defined in claim 1 and R²¹ is a group selected from C₁₋₂alkyl, cyclopentyl, cyclohexyl, pyrrolidinyl and piperidinyl which group is linked to X⁴ through a carbon atom and which C₁₋₂alkyl group may bear one or two substituents selected from hydroxy, halogeno and C₁₋₂alkoxy and which cyclopentyl, cyclohexyl, pyrrolidinyl or piperidinyl group may carry one substituent selected from oxo, hydroxy, halogeno, C₁₋₂alkyl, C₁₋₂hydroxyalkyl and C₁₋₂alkoxy);
- 4) C₂₋₃alkylX⁵C₂₋₃alkylX⁶R²⁷ (wherein X⁵ and X⁶ are as defined in claim 1 and R²⁷ represents hydrogen or C₁₋₂alkyl);
- 5) C₁₋₂alkylR⁶² (wherein R⁶² is a group selected from pyrrolidinyl, piperazinyl, piperidinyl, 1,3-dioxolan-2-yl, 1,3-dioxan-2-yl, 1,3-dithiolan-2-yl and 1,3-dithian-2-yl, which group is linked to C₁₋₂alkyl through a carbon atom and which group may carry one substituent selected from oxo, hydroxy, halogeno, C₁₋₂alkyl, C₁₋₂hydroxyalkyl and C₁₋₂alkoxy) or C₂₋₃alkylR⁶³ (wherein R⁶³ is a group selected from morpholino, thiomorpholino, piperidino, piperazin-1-yl and pyrrolidin-1-yl which group may carry one substituent selected from oxo, hydroxy, halogeno, C₁₋₂alkyl, C₁₋₂hydroxyalkyl and C₁₋₂alkoxy);
- 6) R³⁴ (wherein R³⁴ is as defined in claim 1);

- 7) C_{1-4} alkyl R^{34} (wherein R^{34} is as defined in claim 1);
 8) 1- R^{34} but-2-en-4-yl (wherein R^{34} is as defined in claim 1);
 9) 1- R^{34} but-2-yn-4-yl (wherein R^{34} is as defined in claim 1);
 10) C_{1-3} alkyl X^7R^{34} (wherein X^7 and R^{34} are as defined in claim 1);
 11) 1-($R^{34}X^8$)but-2-en-4-yl (wherein X^8 and R^{34} are as defined in claim 1);
 12) 1-($R^{34}X^9$)but-2-yn-4-yl (wherein X^9 and R^{34} are as defined in claim 1);
 13) ethyl X^{10} methyl R^{34} (wherein X^{10} and R^{34} are as defined in claim 1);
 14) R^{33} (wherein R^{33} is as defined in claim 1); and
 15) ethyl X^{10} methyl R^{33} (wherein X^{10} and R^{33} are as defined in claim 1)].

10

5. A compound as claimed in claim 1 of the formula Ia:

15



20

(Ia)

[wherein:

- R^1 , R^2 and n are as defined in claim 1;
 25 R^{3a} represents hydrogen, hydroxy, methoxy, amino, nitro or halogeno;
 R^{4a} represents hydrogen, hydroxy, halogeno, cyano, nitro, trifluoromethyl, C_{1-3} alkyl, C_{1-3} alkoxy, C_{1-3} alkylthio, $-NR^{6a}R^{7a}$ (wherein R^{6a} and R^{7a} , which may be the same or different, each represents hydrogen or C_{1-3} alkyl) or $R^{8a}(CH_2)_zX^{2a}$ (wherein R^{8a} is a 5 or 6 membered saturated heterocyclic group with one or two heteroatoms, selected independently from O, S
 30 and N, which heterocyclic group may bear one or two substituents selected from oxo, hydroxy, halogeno, C_{1-4} alkyl, C_{1-4} hydroxyalkyl and C_{1-4} alkoxy, z is an integer from 0 to 4 and

X^{2a} represents a direct bond, -O-, -CH₂-, -S-, -SO-, -SO₂-, -NR^{9a}CO-, -CONR^{10a}-, -SO₂NR^{11a}-, -NR^{12a}SO₂- or -NR^{13a}- (wherein R^{9a}, R^{10a}, R^{11a}, R^{12a} and R^{13a} each independently represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl);

X^{1a} represents -O-, -CH₂-, -S-, -SO-, -SO₂-, -NR^{14a}CO-, -CONR^{15a}-, -SO₂NR^{16a}-, -NR^{17a}SO₂- or -NR^{18a}- (wherein R^{14a}, R^{15a}, R^{16a}, R^{17a} and R^{18a} each independently represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) or X^{1a} represents a direct bond;

R^{5a} is selected from one of the following seventeen groups:

- 1) hydrogen or C₁₋₃alkyl which may be unsubstituted or which may be substituted with one or more groups selected from hydroxy, fluoro and amino;
- 10 2) C₁₋₃alkylX^{3a}COR^{19a} (wherein X^{3a} represents -O- or -NR^{20a}- (in which R^{20a} represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R^{19a} represents C₁₋₃alkyl, -NR^{21a}R^{22a}- or -OR^{23a}- (wherein R^{21a}, R^{22a} and R^{23a} which may be the same or different each represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl));
- 15 3) C₁₋₃alkylX^{4a}R^{24a} (wherein X^{4a} represents -O-, -S-, -SO-, -SO₂-, -OCO-, -NR^{25a}CO-, -CONR^{26a}-, -SO₂NR^{27a}-, -NR^{28a}SO₂- or -NR^{29a}- (wherein R^{25a}, R^{26a}, R^{27a}, R^{28a} and R^{29a} each independently represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R^{24a} represents hydrogen, C₁₋₃alkyl, cyclopentyl, cyclohexyl or a 5 or 6 membered saturated heterocyclic group with one or two heteroatoms, selected independently from O, S and N, which C₁₋₃alkyl group may bear one or two substituents selected from oxo, hydroxy, halogeno and C₁₋₄alkoxy and which cyclic group may bear one or two substituents selected from oxo, hydroxy, halogeno, C₁₋₄alkyl, C₁₋₄hydroxyalkyl and C₁₋₄alkoxy);
- 20 4) C₁₋₃alkylX^{5a}C₁₋₃alkylX^{6a}R^{30a} (wherein X^{5a} and X^{6a} which may be the same or different are each -O-, -S-, -SO-, -SO₂-, -NR^{31a}CO-, -CONR^{32a}-, -SO₂NR^{33a}-, -NR^{34a}SO₂- or -NR^{35a}- (wherein R^{31a}, R^{32a}, R^{33a}, R^{34a} and R^{35a} each independently represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R^{30a} represents hydrogen or C₁₋₃alkyl);
- 25 5) C₁₋₃alkylR^{36a} (wherein R^{36a} is a 5 or 6 membered saturated heterocyclic group with one or two heteroatoms, selected independently from O, S and N, which heterocyclic group may bear one or two substituents selected from oxo, hydroxy, halogeno, C₁₋₄alkyl, C₁₋₄hydroxyalkyl and C₁₋₄alkoxy);
- 30 6) C₂₋₃alkenylR^{36a} (wherein R^{36a} is as defined herein);
- 7) C₂₋₃alkynylR^{36a} (wherein R^{36a} is as defined herein);

- 8) R^{37a} (wherein R^{37a} represents a pyridone group, a phenyl group or a 5 or 6-membered aromatic heterocyclic group with 1 to 3 heteroatoms selected from O, N and S, which pyridone, phenyl or heterocyclic group may carry up to 5 substituents selected from hydroxy, halogeno, amino, C₁₋₄alkyl, C₁₋₄alkoxy, C₁₋₄hydroxyalkyl, C₁₋₄aminoalkyl, C₁₋₄alkylamino, C₁₋₄hydroxyalkoxy, carboxy, cyano, -CONR^{38a}R^{39a} and -NR^{40a}COR^{41a} (wherein R^{38a}, R^{39a}, R^{40a} and R^{41a}, which may be the same or different, each represents hydrogen, C₁₋₄alkyl or C₁₋₃alkoxyC₂₋₃alkyl));
- 9) C₁₋₃alkylR^{37a} (wherein R^{37a} is as defined herein);
- 10) C₂₋₃alkenylR^{37a} (wherein R^{37a} is as defined herein);
- 11) C₂₋₃alkynylR^{37a} (wherein R^{37a} is as defined herein);
- 12) C₁₋₃alkylX^{7a}R^{37a} (wherein X^{7a} represents -O-, -S-, -SO-, -SO₂-, -NR^{42a}CO-, -CONR^{43a}-, -SO₂NR^{44a}-, -NR^{45a}SO₂- or -NR^{46a}- (wherein R^{42a}, R^{43a}, R^{44a}, R^{45a} and R^{46a} each independently represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R^{37a} is as defined herein);
- 13) C₂₋₃alkenylX^{8a}R^{37a} (wherein X^{8a} represents -O-, -S-, -SO-, -SO₂-, -NR^{47a}CO-, -CONR^{48a}-, -SO₂NR^{49a}-, -NR^{50a}SO₂- or -NR^{51a}- (wherein R^{47a}, R^{48a}, R^{49a}, R^{50a} and R^{51a} each independently represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R^{37a} is as defined herein);
- 14) C₂₋₃alkynylX^{9a}R^{37a} (wherein X^{9a} represents -O-, -S-, -SO-, -SO₂-, -NR^{52a}CO-, -CONR^{53a}-, -SO₂NR^{54a}-, -NR^{55a}SO₂- or -NR^{56a}- (wherein R^{52a}, R^{53a}, R^{54a}, R^{55a} and R^{56a} each independently represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R^{37a} is as defined herein);
- 15) C₁₋₃alkylX^{10a}C₁₋₃alkylR^{37a} (wherein X^{10a} represents -O-, -S-, -SO-, -SO₂-, -NR^{57a}CO-, -CONR^{58a}-, -SO₂NR^{59a}-, -NR^{60a}SO₂- or -NR^{61a}- (wherein R^{57a}, R^{58a}, R^{59a}, R^{60a} and R^{61a} each independently represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R^{37a} is as defined herein);
- 16) R^{36a} (wherein R^{36a} is as defined herein); and
- 17) C₁₋₃alkylX^{10a}C₁₋₃alkylR^{36a} (wherein X^{10a} and R^{36a} are as defined herein));
- and salts thereof.

6. A compound as claimed in claim 5 wherein R¹ is hydrogen.

30 7. A compound as claimed in claim 5 or claim 6 wherein R² is halogeno, trifluoromethyl, cyano, nitro, C₁₋₃alkanoyl, C₁₋₃alkoxycarbonyl, C₁₋₃alkylsulphinyl, C₁₋

,alkylsulphonyl, carbamoyl, N-C₁₋₃alkylcarbamoyl, N,N-di(C₁₋₃alkyl)carbamoyl, aminosulphonyl, N-C₁₋₃alkylaminosulphonyl, N,N-di(C₁₋₃alkyl)aminosulphonyl, or a group R⁴X¹ (wherein X¹ is as defined in claim 1 and R⁴ represents phenyl or a 5 or 6-membered heterocyclic group with one or two heteroatoms, selected independently from O, S and N, of which at least one is N, which heterocyclic group may be saturated or unsaturated and which phenyl or heterocyclic group may bear one or two substituents selected from hydroxy, halogeno, C₁₋₂alkyl, C₁₋₂alkoxy, C₁₋₂alkanoyloxy, trifluoromethyl, cyano, amino, nitro and C₁₋₂alkoxycarbonyl).

8. A compound as claimed in any one of claims 5 to 7 wherein R^{3a} is hydrogen.

9. A compound as claimed in any one of claims 5 to 8 wherein R^{4a} is hydrogen, hydroxy, cyano, nitro, trifluoromethyl, methyl, ethyl, methoxy, ethoxy, or R^{8a}(CH₂)_{za}X^{2a} (wherein R^{8a} is a group selected from pyrrolidinyl, piperazinyl, piperidinyl, morpholino and thiomorpholino which group may carry one or two substituents selected from oxo, hydroxy, halogeno, C₁₋₂alkyl, C₁₋₂hydroxyalkyl and C₁₋₂alkoxy, X^{2a} is -O-, -S-, -NR^{9a}CO-, -NR^{12a}SO₂- (wherein R^{9a} and R^{12a} each independently represents hydrogen or C₁₋₂alkyl) or NH and za is an integer from 1 to 3).


10. A compound as claimed in any one of claims 5 to 9 wherein X^{1a} is -O-, -S-, -NR^{14a}CO-, -NR^{17a}SO₂- (wherein R^{14a} and R^{17a} each independently represents hydrogen or C₁₋₂alkyl) or NH.

11. A compound as claimed in any one of claims 5 to 10 wherein R^{5a} is selected from one of the following fifteen groups:

- 1) C₁₋₃alkyl which may be unsubstituted or substituted with one or more fluorine atoms, or C₂₋₃alkyl which may be unsubstituted or substituted with one or two groups selected from hydroxy and amino;
- 2) 2-(3,3-dimethylureido)ethyl, 3-(3,3-dimethylureido)propyl, 2-(3-methylureido)ethyl, 3-(3-methylureido)propyl, 2-ureidoethyl, 3-ureidopropyl, 2-(N,N-dimethylcarbamoyloxy)ethyl, 3-

- (N,N-dimethylcarbamoyloxy)propyl, 2-(N-methylcarbamoyloxy)ethyl, 3-(N-methylcarbamoyloxy)propyl, 2-(carbamoyloxy)ethyl, 3-(carbamoyloxy)propyl;
- 3) $C_{2,3}\text{alkyl}X^{4a}R^{24a}$ (wherein X^{4a} is as defined in claim 5 and R^{24a} is a group selected from C_1 , C_2 alkyl, cyclopentyl, cyclohexyl, pyrrolidinyl and piperidinyl which group is linked to X^{4a} through a carbon atom and which $C_{1,2}$ alkyl group may bear one or two substituents selected from hydroxy, halogeno and $C_{1,2}$ alkoxy and which cyclopentyl, cyclohexyl, pyrrolidinyl or piperidinyl group may carry one substituent selected from oxo, hydroxy, halogeno, $C_{1,2}$ alkyl, $C_{1,2}$ hydroxyalkyl and $C_{1,2}$ alkoxy);
- 4) $C_{2,3}\text{alkyl}X^{5a}C_{2,3}\text{alkyl}X^{6a}R^{30a}$ (wherein X^{5a} and X^{6a} are as defined in claim 5 and R^{30a} represents hydrogen or $C_{1,2}$ alkyl);
- 5) $C_{1,2}\text{alkyl}R^{65a}$ (wherein R^{65a} is a group selected from pyrrolidinyl, piperazinyl, piperidinyl, 1,3-dioxolan-2-yl, 1,3-dioxan-2-yl, 1,3-dithiolan-2-yl and 1,3-dithian-2-yl, which group is linked to $C_{1,2}$ alkyl through a carbon atom and which group may carry one substituent selected from oxo, hydroxy, halogeno, $C_{1,2}$ alkyl, $C_{1,2}$ hydroxyalkyl and $C_{1,2}$ alkoxy) or $C_{2,3}\text{alkyl}R^{66a}$ (wherein R^{66a} is a group selected from morpholino, thiomorpholino, piperidino, piperazin-1-yl and pyrrolidin-1-yl which group may carry one substituent selected from oxo, hydroxy, halogeno, $C_{1,2}$ alkyl, $C_{1,2}$ hydroxyalkyl and $C_{1,2}$ alkoxy);
- 6) R^{37a} (wherein R^{37a} is as defined in claim 5);
- 7) $C_{1,4}\text{alkyl}R^{37a}$ (wherein R^{37a} is as defined in claim 5);
- 8) 1- R^{37a} but-2-en-4-yl (wherein R^{37a} is as defined in claim 5);
- 9) 1- R^{37a} but-2-yn-4-yl (wherein R^{37a} is as defined in claim 5);
- 10) $C_{1,3}\text{alkyl}X^{7a}R^{37a}$ (wherein X^{7a} and R^{37a} are as defined in claim 5);
- 11) 1-($R^{37a}X^{8a}$)but-2-en-4-yl (wherein X^{8a} and R^{37a} are as defined in claim 5);
- 12) 1-($R^{37a}X^{9a}$)but-2-yn-4-yl (wherein X^{9a} and R^{37a} are as defined in claim 5);
- 13) ethyl X^{10a} methyl R^{37a} (wherein X^{10a} and R^{37a} are as defined in claim 5);
- 14) R^{36a} (wherein R^{36a} is as defined in claim 5); and
- 15) ethyl X^{10a} methyl R^{36a} (wherein X^{10a} and R^{36a} are as defined in claim 5).

12. A compound as claimed in claim 5 of the formula Ib:



(Ib)

[wherein:

R^{1b} represents hydrogen;

R^{2b} represents halogeno, trifluoromethyl, cyano, nitro, C_{2-3} alkanoyl, C_{1-3} alkoxycarbonyl, C_{1-3} alkylsulphinyl, C_{1-3} alkylsulphonyl, carbamoyl, N - C_{1-3} alkylcarbamoyl, N,N -di(C_{1-3} alkyl)carbamoyl, aminosulphonyl, N - C_{1-3} alkylaminosulphonyl, N,N -di(C_{1-3} alkyl)aminosulphonyl, or a group $R^{6b}X^{2b}$ (wherein X^{2b} represents a direct bond, C_{2-4} alkanoyl, -CONR^{7b}R^{8b}-, -SO₂NR^{9b}R^{10b}- or -SO₂R^{11b}- (wherein R^{7b} and R^{9b} , each independently represents hydrogen and R^{8b} , R^{10b} and R^{11b} each independently represents C_{1-3} alkyl and wherein R^{6b} is linked to R^{8b} , R^{10b} or R^{11b}) and R^{6b} represents pyrrolidinyl, piperazinyl, piperidinyl,

morpholino, thiomorpholino, imidazolyl or triazolyl which heterocyclic group is linked to X^{2b} via a nitrogen atom and which heterocyclic group may bear one or two substituents selected from hydroxy, halogeno, C_{1-2} alkyl, C_{1-2} alkoxy, C_{1-2} alkanoyloxy, trifluoromethyl, cyano, amino, nitro and C_{1-4} alkoxycarbonyl);

nb is an integer from 0 to 2;

R^{3b} represents hydrogen;

R^{4b} represents hydrogen, hydroxy, cyano, nitro, trifluoromethyl, methyl, methoxy or a group $R^{12b}(CH_2)_zX^{3b}$ (wherein R^{12b} represents a group selected from pyrrolidinyl, piperazinyl, piperidinyl, morpholino and thiomorpholino which group may carry one or two substituents selected from oxo, hydroxy, halogeno, C_{1-2} alkyl, C_{1-2} hydroxyalkyl and C_{1-2} alkoxy,

z is an integer from 1 to 3 and X^{3b} represents -O- or -NR^{13b}CO- (wherein R^{13b} represents hydrogen or C_{1-2} alkyl));

X^{1b} represents -O- or -NR^{14b}CO- (wherein R^{14b} represents hydrogen or C₁₋₂alkyl); and R^{5b} represents methyl, ethyl, trifluoromethyl, 2,2,2-trifluoroethyl, 2-hydroxyethyl, 3-hydroxypropyl, 2-methoxyethyl, 3-methoxypropyl, 2-(methylsulphinyl)ethyl, 2-(methylsulphonyl)ethyl, 2-(N,N-dimethylsulphamoyl)ethyl, 2-(N-methylsulphamoyl)ethyl, 3-sulphamoyl, 2-(N,N-dimethylamino)ethyl, 3-(N,N-dimethylamino)propyl, 2-morpholinoethyl, 3-morpholinopropyl, 2-piperidinoethyl, 3-piperidinopropyl, 2-(piperazin-1-yl)ethyl, 3-(piperazin-1-yl)propyl, 2-(pyrrolidin-1-yl)ethyl, 3-(pyrrolidin-1-yl)propyl, (1,3-dioxolan-2-yl)methyl, 2-(1,3-dioxolan-2-yl)ethyl, 2-(2-methoxyethylamino)ethyl, 2-(2-hydroxyethylamino)ethyl, 3-(2-methoxyethylamino)propyl, 3-(2-hydroxyethylamino)propyl, 2-methylthiazol-4-ylmethyl, 2-acetamidothiazol-4-ylmethyl, 1-methylimidazol-2-ylmethyl, 2-(imidazol-1-yl)ethyl, 2-(1,2,3-triazol-1-yl)ethyl, 2-(1,2,3-triazol-2-yl)ethyl, 2-(1,2,4-triazol-1-yl)ethyl, 2-(1,2,4-triazol-4-yl)ethyl, 4-pyridylmethyl, 2-(4-pyridyl)ethyl, 3-(4-pyridyl)propyl 2-(4-pyridyloxy)ethyl, 2-(4-pyridylamino)ethyl, 2-(4-oxo-1,4-dihydro-1-pyridyl)ethyl, 2-thiomorpholinoethyl, 3-thiomorpholinopropyl, 2-(2-methoxyethoxy)ethyl, 2-(4-methylpiperazin-1-yl)ethyl, 3-(4-methylpiperazin-1-yl)propyl, 3-(methylsulphinyl)propyl, 3-(methylsulphonyl)propyl, 2-(2-methylimidazol-1-yl)ethyl, 2-(5-methyl-1,2,4-triazol-1-yl)ethyl, morpholino, 2-((N-(1-methylimidazol-4-ylsulphonyl)-N-methyl)amino)ethyl, 2-((N-(3-morpholinopropylsulphonyl)-N-methyl)amino)ethyl, 2-((N-methyl-N-4-pyridyl)amino)ethyl or 3-(4-oxidomorpholino)propyl;]

20 and salts thereof.

13. A compound as claimed in claim 1 selected from:

- 6,7-dimethoxy-4-(6-fluorooxindol-3-yl)quinazoline,
- 6,7-dimethoxy-4-(oxindol-3-yl)quinazoline,
- 25 7-(2-methoxyethoxy)-4-(oxindol-3-yl)quinazoline,
- 4-(5-bromooxindol-3-yl)-6-methoxy-7-(2-methoxyethoxy)quinazoline,
- 4-(5-hydroxyoxindol-3-yl)-6-methoxy-7-(3-morpholinopropoxy)quinazoline,
- 6-methoxy-7-(3-morpholinopropoxy)-4-(6-trifluoromethyloxindol-3-yl)quinazoline,
- 4-(5-aminosulphonyloxindol-3-yl)-6-methoxy-7-(2-methoxyethoxy)quinazoline,
- 30 4-(5-cyanooxindol-3-yl)-7-(3-morpholinopropoxy)quinazoline,
- 6-methoxy-7-(2-(2-methoxyethoxy)ethoxy)-4-(oxindol-3-yl)quinazoline and

4-(5-aminosulphonyloxindol-3-yl)-6-methoxy-7-(2-(2-methoxyethoxy)ethoxy)quinazoline and salts thereof.

14. A compound as claimed in claim 1 selected from:

- 5 4-(5-fluorooxindol-3-yl)-6-methoxy-7-(3-morpholinopropoxy)quinazoline,
4-(5-fluorooxindol-3-yl)-6-methoxy-7-(2-methoxyethoxy)quinazoline,
4-(5-cyanooxindol-3-yl)-7-(2-(imidazol-1-yl)ethoxy)-6-methoxyquinazoline and
7-(3-morpholinopropoxy)-4-(6-trifluoromethyloxindol-3-yl)quinazoline
and salts thereof.

10

15. A compound as claimed in claim 1 selected from:


- 4-(6-fluorooxindol-3-yl)-6-methoxy-7-(3-morpholinopropoxy)quinazoline,
6-methoxy-7-(2-methoxyethoxy)-4-(oxindol-3-yl)quinazoline and
6-methoxy-7-(3-morpholinopropoxy)-4-(oxindol-3-yl)quinazoline
15 and salts thereof.

16. A compound as claimed in any one of the preceding claims in the form of a pharmaceutically acceptable salt.

20 17. A process for the preparation of a compound of formula I or salt thereof (as defined in claim 1) which comprises:

(a) the reaction of a compound of the formula III:

25




30

(III)

(wherein R^3 and m are as defined in claim 1 and L^1 is a displaceable moiety), with a compound of the formula IV:

5




(IV)

10 (wherein R^1 , R^2 and n are as defined in claim 1) whereby to obtain compounds of the formula I and salts thereof;

(b) the deprotection of a compound of formula V:

15




(V)

20

(wherein R^2 , R^3 , n and m are as defined in claim 1 and P^1 represents a protecting group);

(c) for the preparation of compounds of formula I and salts thereof wherein R^1 is hydrogen, the reduction and cyclisation of a compound of formula VI:

- 157 -




(VI)

(wherein R^2 , R^3 , m and n are as defined in claim 1 and Y represents cyano, carboxy or

5 C_{1-4} alkoxycarbonyl);

(d) for the preparation of compounds of formula I and salts thereof wherein at least one of the R^2 groups is hydroxy the deprotection of a compound of formula VII:




(VII)

(wherein R^1 , R^2 , R^3 , n and m are as defined in claim 1, P^2 represents a phenolic hydroxy protecting group and $p1$ is an integer from 1 to 4 equal to the number of protected hydroxy groups and such that $n-p1$ is equal to the number of R^2 substituents which are not protected hydroxy);

(e) for the preparation of those compounds of formula I and salts thereof wherein at least one R^3 is $R^{10}X^2$ wherein R^{10} is as defined in claim 1 and X^2 is $-O-$, $-S-$, $-SO_2-$, $-CONR^{12}-$, $-SO_2NR^{13}-$ or $-NR^{15}-$ (wherein R^{12} , R^{13} and R^{15} each independently represents hydrogen, C_{1-3} alkyl or C_{1-3} alkoxy C_{2-3} alkyl), the reaction, of a compound of the formula VIII:

- 158 -




(VIII)

- 5 (wherein R^1 , R^2 , R^3 , n and X^2 are as defined in claim 1 and s is an integer from 0 to 3) with a compound of formula IX:



(IX)

- 10 (wherein R^{10} is as defined in claim 1 and L^1 is as herein defined);
 (f) for the preparation of compounds of formula I and salts thereof wherein at least one R^3 is $R^{10}X^2$ wherein R^{10} is as defined in claim 1 and X^2 is -O-, -S-, or -NR¹⁵- (wherein R^{15} represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl), the reaction of a compound of the formula X:




15

(X)

with a compound of the formula XI:

20




(XI)

(wherein L^1 and s are as defined herein and R^1 , R^2 , R^3 , R^{10} , n , and X^2 are all as defined in claim 1);

- (g) for the preparation of compounds of formula I and salts thereof wherein at least one R^3 is $R^{10}X^2$ wherein X^2 is as defined in claim 1 and R^{10} is $C_{1,5}$ alkyl R^{65} , [wherein R^{65} is selected from one of the following six groups:
 - 1) $X^{11}C_{1,3}$ alkyl (wherein X^{11} represents -O-, -S-, -SO₂-, -NR⁶⁶CO- or -NR⁶⁷SO₂- (wherein R^{66} and R^{67} which may be the same or different are each hydrogen, $C_{1,3}$ alkyl or $C_{1,3}$ alkoxy $C_{2,3}$ alkyl) ;
 - 10 2) NR⁶⁸R⁶⁹ (wherein R^{68} and R^{69} which may be the same or different are each hydrogen, $C_{1,3}$ alkyl or $C_{1,3}$ alkoxy $C_{2,3}$ alkyl);
 - 3) $X^{12}C_{1,3}$ alkyl X^6R^{27} (wherein X^{12} represents -O-, -S-, -SO₂-, -NR⁷⁰CO-, -NR⁷¹SO₂- or NR⁷² (wherein R^{70} , R^{71} , and R^{72} which may be the same or different are each hydrogen, $C_{1,3}$ alkyl or $C_{1,3}$ alkoxy $C_{2,3}$ alkyl) and X^6 and R^{27} are as defined in claim 1);
 - 15 4) R^{60} (wherein R^{60} is a 5 or 6 membered saturated heterocyclic group with one or two heteroatoms of which one is N and the other is selected independently from O, S and N, which heterocyclic group is linked to $C_{2,3}$ alkyl through a nitrogen atom and which heterocyclic group may bear one or two substituents selected from oxo, hydroxy, halogeno, $C_{1,4}$ alkyl, $C_{1,4}$ hydroxyalkyl and $C_{1,4}$ alkoxy);
 - 20 5) $X^{13}R^{34}$ (wherein X^{13} represents -O-, -S-, -SO₂-, -NR⁷³CO-, -NR⁷⁴SO₂-, or -NR⁷⁵- (wherein R^{73} , R^{74} , and R^{75} which may be the same or different are each hydrogen, $C_{1,3}$ alkyl or $C_{1,3}$ alkoxy $C_{2,3}$ alkyl) and R^{34} is as defined in claim 1); and
 - 6) $X^{14}C_{1,3}$ alkyl R^{34} (wherein X^{14} represents -O-, -S-, -SO₂-, -NR⁷⁶CO-, -NR⁷⁷SO₂- or -NR⁷⁸- (wherein R^{76} , R^{77} and R^{78} each independently represents hydrogen, $C_{1,3}$ alkyl or $C_{1,3}$ alkoxy $C_{2,3}$ alkyl) and R^{34} is as defined in claim 1);],
- 25 reacting a compound of the formula XII:

- 160 -



(XII)

(wherein L^1 and s are as herein defined and X^2 , R^1 , R^2 , R^3 and n are as defined in claim 1) with
 5 a compound of the formula XIII:



(XIII)

(wherein R^{65} is as defined herein) to give a compound of formula I or salt thereof;

10 (h) for the preparation of those compounds of formula I and salts thereof wherein one or more of the substituents $(R^3)_m$ is represented by $-NR^{79}R^{80}$ -, where one or both of R^{79} and R^{80} are C_{1-3} alkyl, the reaction of a compound of formula I wherein the substituent $(R^3)_m$ is an amino group and an alkylating agent;

for the preparation of compounds of formula I and salts thereof wherein one or more of the

15 substituents R^2 or R^3 is an amino group the reduction of a corresponding compound of formula I wherein the substituent(s) at the corresponding position(s) of the quinazoline and/or benz ring of the oxindole group is/are a nitro group(s);

and when a salt of a compound of formula I is required, reaction of the compound obtained with an acid or base whereby to obtain the desired salt.

20

18. A pharmaceutical composition which comprises as active ingredient a compound of formula I as defined in claim 1 or a pharmaceutically acceptable salt thereof in association with a pharmaceutically acceptable excipient or carrier.

19. A method for producing an antiangiogenic and/or vascular permeability reducing effect in a warm-blooded animal in need of such treatment which comprises administering to said animal an effective amount of a compound of formula I as defined in claim 1 or a pharmaceutically acceptable salt thereof.

5

20. A compound as claimed in any one of claims 1 to 16 or a pharmaceutically acceptable salt thereof for use as a medicament.

10

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 97/01211

A. CLASSIFICATION OF SUBJECT MATTER
 IPC 6 C07D403/04 C07D401/14 A61K31/505

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 95 15758 A (RHONE-POULENC) 15 June 1995 see page 1 - page 13; claims	1,16-20
A	WO 95 23141 A (PFIZER) 31 August 1995 see page 63; claims	1,16-20

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

12 August 1997

Date of mailing of the international search report

18.08.97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax (+31-70) 340-3016

Authorized officer

Francois, J

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/GB 97/01211

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9515758 A	15-06-95	US 5480883 A AU 1305095 A US 5646153 A	02-01-96 27-06-95 08-07-97
WO 9523141 A	31-08-95	AU 2972795 A CA 2183655 A CN 1141633 A EP 0746554 A FI 963283 A HU 76291 A JP 9501953 T NO 963506 A PL 315941 A	11-09-95 31-08-95 29-01-97 11-12-96 22-08-96 28-07-97 25-02-97 22-10-96 09-12-96